



## **Tobacco Consumption During Pregnancy and its Association with Psychosocial Child Development**

***PATRICIA BRENNAN, PhD***

*Emory University, USA*

*(Published online March 27, 2003)  
(2<sup>nd</sup> revised edition February 18, 2011)*

### ***Topic***

*Tobacco and pregnancy*

### **Introduction**

There is a growing body of literature suggesting that maternal tobacco consumption during pregnancy is associated with negative child behavioural outcomes.<sup>1-5</sup> Given the fact that more than half of the women who smoke cigarettes will continue to do so during pregnancy,<sup>6</sup> these tobacco-behavioural linkages may have far-reaching implications for development and mental health in children.

### **Subject**

Studies have linked maternal tobacco consumption during pregnancy to a number of adverse medical outcomes. For example, prenatal maternal smoking has been associated with low birthweight, chronic *ischemia*, *hypoxia*, hypertonicity, increased tremors, and increased startle response in infants.<sup>7-9</sup> It has also been suggested that the deleterious effects of maternal prenatal smoking on child development may extend to the psychosocial realm as well. This report reviews the evidence for the connection between maternal tobacco consumption and psychosocial child development and discusses relevant implications for interventions and public health policy.

### **Problems**

The study of maternal tobacco consumption and its effect on child behavioural or psychosocial outcomes is fraught with methodological problems. The most serious methodological concern is the difficulty in establishing a causal connection due to the potential role of confounds in this process.<sup>10</sup> Research in this area is quasi-experimental. For obvious ethical reasons, women are not randomly assigned to smoke or not to smoke during pregnancy. However, the factors identified with women who do smoke typically differ from those of women who do not in a number of different areas, including genetic background, childhood history of antisocial behaviour, socioeconomic status, mental health, personality traits, parenting styles, and exposure to stressful life events. These factors, in turn, are associated with greater risk for problematic child psychosocial development. However, no single study has been able to control for all of these potential confounds. Another methodological concern is the common use of retrospective rather

than prospective reports of maternal smoking. The recent finding that maternal self-reports of smoking during pregnancy are highly correlated with direct biochemical measures<sup>11</sup> alleviates concerns about the exclusive use of self-reports of maternal smoking in this research area.

### Research Context

Maternal prenatal smoking and its relationship to child psychosocial outcome has been examined in both cross-sectional and longitudinal studies, and in both clinical and community samples. Animal studies have examined the effect of nicotine exposure on behavioural outcomes and brain functioning and neurobiological deficits have been suggested as a likely mediator for negative behavioural outcomes.<sup>12</sup> Two recent human studies support this contention, noting associations between maternal prenatal smoking and decreased *frontal lobe* volumes in infants,<sup>13</sup> and a thinning of the *cerebral cortex* in adolescents.<sup>14</sup> Nevertheless, there remains disagreement in the field concerning the relative importance of the direct effects of prenatal tobacco smoking, its associated familial background factors, and potentially moderating environmental or genetic vulnerabilities in the prediction of negative child behavioural outcomes.

### Key Research Questions

The key research questions in this area are as follows:

- a) Is maternal tobacco consumption during pregnancy associated with deleterious behavioural outcomes in youth? And, if so:
- b) Can potential methodological confound account for this association? And, if not:
- c) Are these risks specific to particular behavioural outcomes?
- d) Are these risks moderated by other factors including genetics, the gender of the child, or the environmental context of development?

### Recent Research Results

Maternal prenatal smoking has been associated with increased risk for Attention Deficit Hyperactivity Disorder, oppositional behaviour, aggression, conduct disorder, and substance abuse outcomes in youth.<sup>1,2,5,15,16</sup> A majority of studies suggest that statistical control for a range of potential confounds, including parental criminality, maternal mental health, parenting behaviour, socioeconomic status, prenatal exposure to drugs and alcohol, and other perinatal complications, does not change the general pattern of results.<sup>1-5,15,16</sup> However, a few recent studies have found that the maternal prenatal smoking child outcome relationship is no longer significant when maternal background characteristics (e.g., childhood history of conduct disorder) and mother-child relationship qualities are taken into account.<sup>17-19</sup> Although evidence from twin studies suggests that the relationship between maternal smoking during pregnancy and child behaviour problems cannot be entirely accounted for by genetic influences,<sup>17,20</sup> Recent studies using innovative design strategies have suggested that genetic or familial background factors may be essential components of the prenatal smoking and child externalizing behaviour association.<sup>21,22</sup>

The noted maternal prenatal smoking child behaviour outcome association appears to be specific to externalizing or acting out behaviours; there does not appear to be an

association between maternal prenatal smoking and increased risk for internalizing problems such as depression.<sup>10</sup> In addition, gender appears to moderate the effects of maternal prenatal smoking on child psychosocial outcomes. Specifically, results are stronger for males in terms of the outcome of conduct disorder, and stronger for females in terms of the outcome of substance abuse.<sup>16,23</sup>

Family and socioeconomic context has been shown to moderate the effect of maternal prenatal smoking on child outcomes.<sup>1,24,25</sup> Recent gene by environment interaction studies also suggest that several distinct genetic *polymorphisms*, (including one that effects the metabolism of smoking-related carcinogens), may moderate the association between maternal prenatal smoking and child externalizing behaviour.<sup>26-28</sup>

Maternal genetic profiles have also been associated with the reduction or spontaneous quitting of smoking during pregnancy.<sup>29</sup> Further study of maternal psychosocial and genetic characteristics associated with cessation of smoking during pregnancy is needed to more effectively design intervention programs focused on pregnant women.

### Conclusions

There are several possible mechanisms or explanations for the noted relationship between maternal prenatal smoking and behavioural problems in offspring. One possible explanation is that prenatal exposure to this *teratogen* increases the risk for child externalizing problems, but only in genetically or otherwise environmentally vulnerable individuals. Alternatively, maternal prenatal smoking may serve as a marker for other environmental effects that increase the risk of externalizing problems in children. For example, maternal prenatal smoking may be an indicator of a passive, neglectful parenting style. It may not reflect abuse or overt parental hostility but rather a subtle disruption in the parent-child relationship that is difficult to measure via questionnaires or short-term observations, but that nevertheless increases the risk for externalizing problems in children. An additional possibility is that maternal cigarette smoking may set off a chain of transactional biological and environmental factors that act together to increase risk for deleterious child development. Our understanding of this transactional process is rudimentary at this time.

### Implications

Not all children whose mothers smoked during pregnancy will manifest deleterious behavioural or developmental outcomes. Future studies aimed at assessing the potentially moderating risk and protective factors in this process would be useful in designing effective prevention and intervention programs. A public health approach calls for prevention and intervention strategies designed to reduce the known risk factors for these deleterious psychosocial outcomes in children. Maternal prenatal smoking is a relatively modifiable perinatal risk factor. Smoking cessation programs for pregnant women,<sup>30</sup> (even low-intensity counselling interventions by general practitioners<sup>31</sup>), have been found to reduce or eliminate maternal smoking during pregnancy. An examination of the behavioural profiles of the children whose mothers successfully completed such programs would help provide important experimental evidence concerning the potentially causal role of maternal prenatal smoking on child behaviour problems.

**To learn more on this topic, consult the following sections of the Encyclopedia:**

- [How important is it?](#)
- [What do we know?](#)
- [What can be done?](#)
- [According to experts](#)
- [Key messages](#)

## REFERENCES

1. Wakschlag LS, Henry DB, Blair RJ, Dukic V, Burns J, Pickett KE. Unpacking the association: Individual differences in the relation of prenatal exposure to cigarettes and disruptive behavior phenotypes. *Neurotoxicology and Teratology* 2011;33(1):145-54.
2. Hutchinson J, Pickett KE, Green J, Wakschlag LS. Smoking in pregnancy and disruptive behaviour in 3-year-old boys and girls: an analysis of the UK Millennium Cohort Study. *Journal of Epidemiological and Community Health* 2010;64(1):82-8.
3. Stene-Larsen K, Borge AIH, Vollrath ME. Maternal smoking in pregnancy and externalizing behavior in 18-month-old children: results from a population-based prospective study. *Journal of the American Academy of Child and Adolescent Psychiatry* 2009;48(3):283-289.
4. Robinson M, Oddy WH, Li J, Kendall GE, de Klerk NH, Silburn SR, Zubrick SR, Newnham JP, Stanley FJ, Mattes E. Pre- and postnatal influences on preschool mental health: a large-scale cohort study. *Journal of Child Psychology and Psychiatry* 2008;49(10):1118-1128.
5. Linnet KM, Dalsgaard S, Obel C, Wisborg K, Henriksen TB, Rodriquez A, Kotimaa A, Moilanen I, Thomsen PH, Olsen J, Jarvelin M. Maternal lifestyle factors in pregnancy risk of attention deficit hyperactivity disorder and associated behaviors: Review of the current evidence. *American Journal of Psychiatry* 2003;160(6):1028-1040.
6. Centers for Disease Control and Prevention. National Center for Chronic Disease Prevention and Health Promotion. *Preventing smoking and exposure to secondhand smoke before, during, and after pregnancy*. Atlanta, GA: National Center for Chronic Disease Prevention and Health Promotion. Centers for Disease Control and Prevention; 2007.
7. Fried PA. Prenatal exposure to tobacco and marijuana: effects during pregnancy, infancy, and early childhood. *Clinical Obstetrics and Gynecology* 1993;36(2): 319-337.
8. Fried PA, Watkinson B, Dillon RF, Dulberg CS. Neonatal neurological status in a low-risk population after prenatal exposure to cigarettes, marijuana and alcohol. *Journal of Developmental and Behavioral Pediatrics* 1987;8(6):318-326.

9. Pollack H, Lantz PM, Frohna JG. Maternal smoking and adverse birth outcomes among singletons and twins. *American Journal of Public Health* 2000;90(3):395-400.
10. Brion MJ, Victora C, Matijasevich A, Horta B, Anselmi L, Steer C, Menezes AM, Lawlor DA, Davey Smith G. Maternal smoking and child psychological problems: Disentangling causal and noncausal effects. *Pediatrics* 2010;126(1):e57-65.
11. McDonald SD, Perkins SL, Walker MC. Correlation between self-reported smoking status and serum cotinine during pregnancy. *Addictive Behaviors* 2005;30(4):853-857.
12. Fung YK. Postnatal behavioral effects of maternal nicotine exposure in rats. *Journal of Pharmacy and Pharmacology* 1988;40(12):870-872.
13. Ekblad M, Korkeila J, Parkkola R, Lapinleimu H, Haataja L, Lehtonen L, PIPARI Study Group. Maternal smoking during pregnancy and regional brain volumes in preterm infants. *Journal of Pediatrics* 2010;156(2):185-190.
14. Toro R, Leonard G, Lerner JV, Lerner RM, Perron M, Pike GB, Richer L, Veillette S, Pausova Z, Paus T. Prenatal exposure to maternal cigarette smoking and the adolescent cerebral cortex. *Neuropsychopharmacology* 2008;33(5):1019-1027.
15. Day NL, Richardson GA, Goldschmidt L, Cornelius MD. Effects of prenatal tobacco exposure on preschoolers' behavior. *Journal of Developmental and Behavioral Pediatrics* 2000;21(3):180-188.
16. Weissman MM, Warner V, Wickramaratne PJ, Kandel DB. Maternal smoking during pregnancy and psychopathology in offspring followed to adulthood. *Journal of the American Academy of Child and Adolescent Psychiatry* 1999;38(7):892-899.
17. Maughan B, Taylor A, Caspi A, Moffitt TE. Prenatal smoking and early childhood conduct problems: Testing genetic and environmental explanations of the association. *Archives of General Psychiatry* 2004;61(8):836-843.
18. Roza SJ, Verhulst FC, Jaddoe VW, Steegers EA, Mackenbach JP, Hofman A, Tiemeier H. Maternal smoking during pregnancy and child behaviour problems: the Generation R Study. *International Journal of Epidemiology* 2009; 38(3):680-689.
19. Lavigne JV, Hopkins J, Gouze KR, Bryant FB, Lebailly SA, Binns HJ, Lavigne PM. Is smoking during pregnancy a risk factor for psychopathology in young children? A methodological caveat and report on preschoolers. *Journal of Pediatric Psychology* 2011;36(1):10-24.
20. Thapar A, Fowler T, Rice F, Scourfield J, van den Bree M, Thomas H, Harold G, Hay D. Maternal smoking during pregnancy and attention deficit hyperactivity disorder symptoms in offspring. *American Journal of Psychiatry* 2003;160(11):1985-1989.
21. D'Onofrio BM, Van Hulle CA, Waldman ID, Rodgers JL, Harden KP, Rathouz PJ, Lahey BB. Smoking during pregnancy and offspring externalizing problems: An exploration of genetic and environmental confounds. *Developmental Psychopathology* 2008;20(1):139-164.

22. Thapar A, Rice F, Hay D, Boivin J, Langley K, van den Bree M, Rutter M, Harold G. Prenatal smoking might not cause Attention-Deficit/Hyperactivity Disorder: Evidence from a novel design. *Biological Psychiatry* 2009;66(8):722-727.
23. Fergusson DM, Woodward LJ, Horwood LJ. Maternal smoking during pregnancy and psychiatric adjustment in late adolescence. *Archives of General Psychiatry* 1998;55(8):721-727.
24. Wakschlag LS, Hans SL. Maternal smoking during pregnancy and conduct problems in high-risk youth: A developmental framework. *Development and Psychopathology* 2002;14(2):351-369.
25. Huijbregts Huijbregts SC, Séguin JR, Zoccolillo M, Boivin M, Tremblay RE. Maternal prenatal smoking, parental antisocial behavior, and early childhood physical aggression. *Development and Psychopathology* 2008;20:437-453.
26. Neuman RJ, Lobos E, Reich W, Henderson CA, Sun LW, Todd RD. Prenatal smoking exposure and dopaminergic genotypes interact to cause a severe ADHD subtype. *Biological Psychiatry* 2007;61(12):1320-1328.
27. Hsieh CJ, Jeng SF, Su YN, Liao HF, Hsieh WS, Wu KY, Chen PC. CYP1A1 modifies the effect of maternal exposure to environmental tobacco smoke on child behavior. *Nicotine and Tobacco Research* 2010;12(11):1108-1117.
28. Wakschlag LS, Kistner EO, Pine DS, Biesecker G, Pickett KE, Skol AD, Dukic V, Blair RJ, Leventhal BL, Cox NJ, Burns JL, Kasza KE, Wright RJ, Cook EH Jr. Interaction of prenatal exposure to cigarettes and MAOA genotype in pathways to youth antisocial behavior. *Molecular Psychiatry* 2010;15(9):928-937.
29. Chen X, Woodcroft KJ. Polymorphisms in metabolic genes CYP1A1 and GSTM1 and changes in maternal smoking during pregnancy. *Nicotine and Tobacco Research* 2009;11(3):225-33.
30. Ford RPK, Cowan SF, Schluter PJ, Richardson AK, Wells JE. Smoke Change for changing smoking in pregnancy. *New Zealand Medical Journal* 2001;114(1128):107-110.
31. Melvin CL, Dolan-Mullen P, Windsor RA, Whiteside HP, Goldenberg RL. Recommended cessation counselling for pregnant women who smoke: a review of the evidence. *Tobacco Control* 2000;9(suppl. 3):80-84.

To cite this document:

Brennan P. Tobacco consumption during pregnancy and its association with psychosocial child development. 2<sup>nd</sup> rev ed. Wakschlag LS, topic ed. In: Tremblay RE, Boivin M, Peters RDeV, Barr RG, eds. *Encyclopedia on Early Childhood Development* [online]. Montreal, Quebec: Centre of Excellence for Early Childhood Development; 2011:1-7. Available at: <http://www.child-encyclopedia.com/documents/BrennanANGxp3.pdf>. Accessed [insert date].

Copyright © 2003-2011

**This article is funded by the Centre of Excellence for Early Childhood Development (CEECD) and the Strategic Knowledge Cluster on ECD (SKC-ECD)**



STRATEGIC KNOWLEDGE  
CLUSTER ON EARLY  
*child development*