



Services and Programs Proven to be Effective in Managing Pediatric Sleep Disturbances and Disorders, and Their Impact on the Social and Emotional Development of Young Children

**KARYN G FRANCE, PhD, Dip Clin Psych &
NEVILLE M BLAMPIED, MSc, FNZPsS**

University of Canterbury, NEW ZEALAND

(Published online March 11, 2004)

Topic

Sleeping behaviour

Introduction

During the preschool years, dramatic changes take place in a typical child's sleep, while enormous changes in physical, linguistic, cognitive and social development occur that profoundly alter both waking activities and sleep regulation. Establishing sleep habits that meet a child's individual needs and are adapted to her/his culture and family circumstances is vital to individual and family well-being. Within a broad range of individual, familial and cultural variations,¹ by the end of the preschool period a child who is a "good" sleeper will have a regular but not ritualized, emotionally and socially positive pre-bed routine, free of resistance and coercion. The child will be put to bed awake, without difficulty, by a variety of caregivers, and sleep independently wherever appropriate to family culture and circumstances. Sleep onset will be rapid, both initially and after later wakings, without crying, calling-out or adult attention, unless the child is ill or needs care, so that sleep is of age-appropriate duration and quality.^{2,3}

Achieving this outcome requires continuous, dynamic, learned adjustments affecting every aspect of sleep and involving all parents, caregivers, siblings and other family members. This is influenced by child temperament; parental adjustment, resources and practices; maternal health and well-being; and family/community circumstances.³ Careful assessment of family circumstances and environment and child development is necessary in diagnosing pediatric sleep disturbances (PSDs)^{4,5} Sleep may be measured by parental diaries (e.g. France & Hudson⁶), activity monitoring,⁷ infra-red video recording⁸ and clinic-based, multi-channel physiological recording (polysomnography).⁹

Subject

PSDs, a common reason for attending family health services,^{10,11} may be broadly differentiated into a psycho-social group focused on parent-child interaction and a group

(henceforth referred to as the bio-maturational group) in which atypical biological, especially neural, maturation appears to be critical.^{12,13}

Psycho-social PSDs include:

- a) Problems of bed resistance and sleep location. Children may resist/delay being prepared for and placed into bed with tantrums, escape/avoidance and demands for pre-bed rituals. Also, they may sleep often in locations other than where parents desire (e.g. co-sleeping with parents or siblings) because the child moves or is moved from their own cot or bed to stop or prevent them crying and to obtain sleep;
- b) Problems of sleep-onset delay and recurrent night waking. The infant or child needs parental attendance and attention to go to sleep initially or to resume sleep after later wakings;
- c) Fears and anxieties associated with bedtime, night-time and sleep.

Bio-maturational PSDs include:

- a) Parasomnias. These are undesirable behaviours occurring during sleep or sleep-wake transitions, including sleepwalking/talking, sleep terrors and rhythmic movement disorders, such as head banging and body rocking and also nocturnal enuresis (bedwetting);
- b) Circadian rhythm disorders, in which the individual's sleep-wake phases are not in synchrony with those of the family or community.

Psycho-social PSDs commonly co-occur, and may affect 15 to 35% of families.^{14,15} Bio-maturational PSDs are much less common, chronically affecting 1 to 3% of families,¹⁶ but children with parasomnias also frequently exhibit psycho-social PSDs.¹⁶ Little evidence links PSDs to family demographic variables, but more boys than girls may be affected by parasomnias.¹⁷

Obstructive sleep apnea (noisy breathing and profuse sweating) and other breathing difficulties are primarily problems of airway functioning and respiratory control during sleep.¹⁸ Any infant or child with symptoms of sleep apnea or anoxia (lack of oxygen) needs urgent medical evaluation. Some infants experience episodes of anoxia while asleep, often resulting in death in infants less than 12 months old (Sudden Infant Death Syndrome - SIDS). Risk of SIDS is reduced by placing infants on their back to sleep,¹⁹ by breastfeeding and by avoiding exposure to cigarette smoke and co-sleeping.²⁰

Problems

PSDs predict sleep disturbances and behavioural difficulties later in childhood^{21,22} and potentially throughout life, and sleep quality is linked to intellectual, emotional and social development.²³ If chronic or severe, PSDs are stressful for the child, siblings and parents, contributing to attachment difficulties, disruptions of learning, depression, family conflict and marital breakdown,^{11,24} and to overmedication with prescription and non-prescription drugs.¹⁴

Research Context

Considerable research has investigated the developmental neurophysiology of sleep from infancy onwards. Over the first few months of life, sleep is coordinated into a day-night pattern and consolidated. Cycles of rapid-eye-movement sleep (REM) and non-REM sleep shift from rapid cycling and 1:1 distribution at birth to a 1:2 distribution at eight months, and deep, non-REM sleep (associated with parasomnias) predominates early in sleep, while REM (associated with awakenings, dreams and nightmares) occurs more later.¹⁷ Research into factors predictive of PSDs reveals associations with first-born status, colic, difficult infant temperament, maternal depression and insecure adult attachment, and diversity in parenting strategies (see France & Blampied³ for a review). There is more treatment research for psycho-social than for bio-maturational PSDs, and this has shifted from mostly case studies to well-controlled investigations. Some treatments have achieved empirically validated status as *well-established*, *probably efficacious* [i.e. effective] or *promising*,²⁵⁻²⁹ using criteria from Chambless and Hollon.³⁰

Key Research Questions

Research has focused on how to facilitate the development of infants' ability to self-soothe so that sleep initiation is under child- rather than other-related cues. Understanding the behavioural trap by which parent-child interactions shape and maintain sleep disruptions has stimulated development of behavioural treatments and adaptations thereof, with concerns as to their effectiveness, acceptability, impact on attachment, adjustment and family well-being, and cultural appropriateness.

Recent Research Results

Parent education, at about birth or later, on infant sleep management^{31,32} and in regulating breastfeeding to optimize night sleep duration³³ facilitates sleep development and may prevent PSDs from developing.³⁴

Systematically structuring pre-bed routines using quiet, pleasant activities and praise for compliance (termed *Positive Routines*) reduces pre-bed tantrums and resistance.³⁵ Crying and calling out etc. during initial settling time or following later night wakings is reduced or eliminated by a range of interventions [variously called *Extinction*, *Graduated Extinction*, and *(Graduated) Planned Ignoring*; see Mindell²⁷]. All involve the immediate or progressive (graduated) delay/withdrawal of parental attention for sleep-disruptive behaviour, thereby (in principle) removing the reinforcer for the behaviour, a process termed behavioural extinction.² In older, more verbal children this can be supplemented by adding shaping and positive reinforcement (praise, tangible rewards) for achieving appropriate sleep,^{36,37} while in infants, modifying the withdrawal of adult attention by adding *Parental Presence*, in which the parent lies near the child but does not interact with them until the child goes to sleep,⁷ reduces distress to low levels. Positive Routines may be supplemented by adjusting bedtime later or earlier depending on sleep latency (*Bedtime Fading*) and by removal from bed and being kept awake when not sleeping (*Response Cost*).³⁸ Combining reducing doses of a sedative drug with planned ignoring also reduces distress,³⁹ but sedative drugs used alone have at best short-term effects.^{26,40}

Parents need to be carefully prepared for any intervention, supported during it⁴¹ and warned of the possibility of both initial brief increases in the frequency or intensity of behaviour following the removal of reinforcers (post-extinction response bursts) that may exacerbate sleep disturbance briefly upon treatment initiation,⁴⁰ and the possibility of spontaneous recovery of PSD following illness or changes in routine.⁴¹ Whether unmodified or modified, procedures involving withdrawing parental attention are largely non-stressful for parents and positive for the family⁴²⁻⁴⁴ and, importantly, have no reported adverse effects on child well-being or development.⁴⁵

Night-time fears/anxieties are reduced by treatments involving relaxation, modelling coping, positive thoughts/imagery and positive rewards for “bravery.”⁴⁶

There is comparatively little controlled research into treatments for bio-maturational PSDs (see Owens, France, & Wiggs²⁸ for a review). Scheduled awakenings, in which parents use baseline information to predict the time of a parasomnia event and wake the child 15 to 30 minutes beforehand, have successfully treated sleepwalking and sleep terrors.⁴⁷ Waking (via a urine alarm) is also an effective treatment for nocturnal enuresis,⁴⁸ although this is generally used only for older children. Infants and children with chronic illness, disabilities and special needs may experience high rates of PSDs, but there is little systematic research on treatment for such children.⁴⁹

Conclusions

The neuro-development of sleep and its importance to development are relatively well understood. PSDs are systematically described and diagnosed, and the psycho-social versus bio-maturational distinction is well established; however, the causes of and risk-factors for PSDs are less well specified. Development of good sleep habits in the first year of life depends on the infant learning to self-soothe and on the parents avoiding inadvertently reinforcing sleep-disruptive behaviours. Teaching parents how to structure their bedtime and sleep-related interactions with their infant/child so that self-soothing occurs and sleep-disruptive behaviour is not reinforced may prevent as well as treat PSDs. These treatments may be tailored, by gradual adjustment of parental attention, parental presence and/or brief use of sedatives, so as to reduce stress, apprehension and infant distress. Effective interventions promote family well-being and do not adversely affect child development. More research is needed into bio-maturational PSDs, into services for families facing chronic child illness and disabilities and into cultural factors.

Implications

- Staff working in pediatric/family services settings need regular training in best practice for the diagnosis and treatment of PSDs.
- PSDs need to be understood and treated within an ecological perspective for the child and the family.
- Parental and staff expectations that interventions will be stressful or distressing, or that they will have long-term ill-effects on the child or the family, can be countered by substantial evidence to the contrary where well-designed and properly supported interventions are used.

SLEEPING BEHAVIOUR

- Untreated, chronic PSDs, especially if severe and/or disruptive, have the potential for long-term negative consequences for the child and his/her family and should be treated promptly and effectively.
- While interventions (other than for children who are ill, disabled or have special needs) are typically brief, parents need good preparation for and support during the critical time.
- Interventions that employ best-practice procedures should have relatively rapid positive effects and these should be maintained long-term.
- Research needs to focus on improving and extending preventive interventions; matching treatments to families; improving the range and quality of services for children who are ill, disabled or have special needs; and assessing long-term impacts on the target child and her/his family.

REFERENCES

1. Kawasaki C, Nugent, JK, Miyshita H, Miyahara H, Brazelton TB. The cultural organization of infants' sleep. *Children's Environments Quarterly* 1994;11(2):135-141.
2. Blampied NM, France KG. A behavioural model of infant sleep disturbance. *Journal of Applied Behavior Analysis* 1993;26(4):477-492.
3. France KG, Blampied NM. Infant sleep disturbance: Description of a problem behaviour process. *Sleep Medicine Reviews* 1999;3(4):265-280.
4. France KG, Blampied NM, Henderson JMT. Infant sleep disturbance. *Current Paediatrics* 2003;13(3):241-246.
5. France KG, Henderson JMT, Hudson SM. Fact, act, tact: A three-stage approach to treating sleep problems of infants and young children. *Child & Adolescent Psychiatric Clinics of North America* 1996;5(3):581-599.
6. France KG, Hudson SM. Behavior management of infant sleep disturbance. *Journal of Applied Behavior Analysis* 1990;23(1):91-98.
7. Sadeh A. Assessment of intervention for infant night waking: Parental reports and activity-based home monitoring. *Journal of Consulting & Clinical Psychology* 1994;62(1):63-68.
8. Anders TF, Sostek AM. The use of time lapse video recording of sleep-wake behaviour in human infants. *Psychophysiology* 1976;13(2):155-158.
9. Erler T, Wischniewski E. Sleep medicine in infants – practicability and limitations. *Early Human Development* 2001;63(1):23-35.
10. Keren M., Feldman R., Tyano S. Diagnoses and interactive patterns of infants referred to a community-based infant mental health clinic. *Journal of the American Academy of Child & Adolescent Psychiatry* 2001;40(1):27-35.
11. Mindell JA, Moline ML, Zendell SM, Brown LW, Fry JM. Pediatricians and sleep disorders: Training and practice. *Pediatrics* 1994;94(2 Pt 1):194-200.
12. Anders TF, Eiben LA. Pediatric sleep disorders: A review of the past 10 years. *Journal of the American Academy of Child & Adolescent Psychiatry* 1997;36(1):9-20.
13. Thiedke CC. Sleep disorders and sleep problems in childhood. *American Family Physician* 2001;63(2):277-284.
14. Armstrong KL, Quinn RA, Dadds MR. The sleep patterns of normal children. *Medical Journal of Australia* 1994;161(3):202-206.
15. Scott G, Richards MPM. Night waking in 1-year old children in England. *Child: Care, Health & Development* 1990;16(5):283-302.
16. Mehlenbeck R, Spirito A, Owens J, Boegers J. The clinical presentation of childhood partial arousal parasomnias. *Sleep Medicine* 2000;1(4):307-312.
17. Anders TF. Neurophysiological studies of sleep in infants and children. *Journal of Child Psychology & Psychiatry & Allied Disciplines* 1982;23(1):75-83.
18. Gaultier C. Sleep apnoea in infants. *Current Paediatrics* 2003;13(1):64-68.
19. Willinger M, Hoffman HJ, Hartford RB. Infant sleep position and risk for sudden infant death syndrome: Report of meeting held January 13 and 14, 1994, National Institutes of health, Bethesda, MD. *Pediatrics* 1994;93(5):814- 819.

20. Gunn AJ, Gunn TR, Mitchell EA. Is changing the sleep environment enough? Current recommendations for SIDS. *Sleep Medicine Reviews* 2000;4(5):453-469.
21. Pollock JI. Predictors and long-term associations of reported sleeping difficulties in infancy. *Journal of Reproductive & Infant Psychology* 1992;10(3):151-168.
22. Pollock JI. Night-waking at five years of age: Predictors and prognosis. *Journal of Child Psychology & Psychiatry & Allied Disciplines* 1994;35(4):699-708.
23. Dahl RE. The regulation of sleep and arousal: Development and psychopathology. *Development & Psychopathology* 1996;8(1):3-27.
24. Stoleru S., Nottelman ED, Belmont B, Ronsaville D. Sleep problems in children of affectively ill mothers. *Journal of Child Psychology & Psychiatry & Allied Disciplines* 1997;38(7):831-841.
25. Kuhn BR, Elliott AJ. Treatment efficacy in behavioral pediatric sleep medicine. *Journal of Psychosomatic Research* 2003;54(6):587-597.
26. Kuhn BR, Weidinger D. Interventions for infant and toddler sleep disturbance: A review. *Child & Family Behavior Therapy* 2000;22(2):33-50.
27. Mindell JA. Empirically supported treatments in pediatric psychology: Bedtime refusal and night wakings in young children. *Journal of Pediatric Psychology* 1999;24(6):465-481.
28. Owens JL, France KG, Wiggs L. Behavioural and cognitive behavioural interventions for sleep disorders in infants and children: A review. *Sleep Medicine Reviews* 1999;3(4):281-302.
29. Ramchandani P, Wiggs L, Webb V, Stores G. A systematic review of treatments for settling problems and night waking in young children. *British Medical Journal* 2000;320(7229):209-213.
30. Chambless DL, Hollon SD. Defining empirically supported therapies. *Journal of Consulting & Clinical Psychology* 1998;66(1):7-18.
31. Adair R, Zuckerman B, Bauchner H, Philipp B, Levenson S. Reducing night waking in infancy: A primary care intervention. *Pediatrics* 1992;89(4):585-588.
32. Wolfson A, Lacks P, Futterman A. Effects of parent training on infant sleeping patterns, parents' stress and perceived parental competence. *Journal of Consulting & Clinical Psychology* 1992;60(1):41-48.
33. Pinilla T, Birch LL. Help me make it through the night: Behavioral entrainment of breast-fed infant's sleep patterns. *Pediatrics* 1993;91(2):436-444.
34. Kerr SM, Jowett SA, Smith, LN. Preventing sleep problems in infants: A randomized controlled trial. *Journal of Advanced Nursing* 1996;24(5):938-942.
35. Adams LA, Rickert VI. Reducing bedtime tantrums: Comparison between positive routines and graduated extinction. *Pediatrics* 1989;84(5):756-761.
36. Sanders MR, Bor B, Dadds MR. Modifying bedtime disruptions in children using stimulus control and contingency management techniques. *Behavioural Psychotherapy* 1984;12(2):130-141.
37. Ronnen T. Intervention package for treating sleep disorders in a four-year-old girl. *Journal of Behaviour Therapy & Experimental Psychiatry* 1991;22(2):141-148.
38. Piazza CC, Fisher W. A faded bedtime with response cost protocol for treatment of multiple sleep problems in children. *Journal of Applied Behavior Analysis* 1991;24(1):129-140.

39. France KG, Blampied NM, Wilkinson P. Treatment of infant sleep disturbance by trimeprazine in combination with extinction. *Journal of Developmental & Behavioral Pediatrics* 1991;12(5):308-314.
40. France KG, Blampied NM, Wilkinson P. A multiple-baseline, double-blind evaluation of the effects of trimeprazine tartrate on infant sleep disturbance. *Experimental & Clinical Psychopharmacology* 1999;7(4):502-513.
41. France KG. Handling parents' concerns regarding the behavioural treatment of infant sleep disturbance. *Behavior Change* 1994;11(2):101-109.
42. Durand VM, Mindell JA. Behavioral treatment of multiple childhood sleep disorders: Effects on child and family. *Behavior Modification* 1990;14(1):37-49.
43. Lawton C, France KG, Blampied NM. Treatment of infant sleep disturbance by graduated extinction. *Child & Family Behavior Therapy* 1991;13(1):39-56.
44. Minde K, Faucon A, Falkner S. Sleep problems in toddlers: Effects of treatment on their daytime behavior. *Journal of the American Academy of Child and Adolescent Psychiatry* 1994;33(8):1114-1121.
45. France KG. Behavior characteristics and security in sleep-disturbed infants treated with extinction. *Journal of Pediatric Psychology* 1992;17(4):467-475.
46. Ollendick TH, Hagopian LP, Huntzinger RM. Cognitive-behavior therapy with nighttime fearful children. *Journal of Behavior Therapy & Experimental Psychiatry* 1991;22(2):113-121.
47. Frank NC, Spirito A, Stark L, Owens-Stively J. The use of scheduled awakenings to eliminate childhood sleep walking. *Journal of Pediatric Psychology* 1997;22(3):345-353.
48. Mellon MW, McGrath ML. Empirically supported treatments in pediatric psychology. Nocturnal enuresis. *Journal of Pediatric Psychology* 2000;25(4):193-214.
49. Wiggs L, France KG. Behavioural treatments for sleep problems in children and adolescents with physical illness, psychological problems or intellectual disabilities. *Sleep Medicine Reviews* 2000;4(3):299-314.

To cite this document:

France KG, Blampied NM. Services and programs proven to be effective in managing pediatric sleep disturbances and disorders, and their impact on the social and emotional development of young children. In: Tremblay RE, Barr RG, Peters RDeV, eds. *Encyclopedia on Early Childhood Development* [online]. Montreal, Quebec: Centre of Excellence for Early Childhood Development; 2004:1-8. Available at: <http://www.child-encyclopedia.com/documents/France-BlampiedANGxp.pdf>. Accessed [insert date].

Copyright © 2004