



## **Prenatal Stress and Offspring Development in Nonhuman Primates**

**MARY L. SCHNEIDER, PhD, OTR**  
**COLLEEN F. MOORE, PhD**

*University of Wisconsin-Madison, USA*

*(Published online April 3, 2003)*

### **Topic**

*Stress (prenatal and perinatal)*

### **Introduction**

Psychosocial stress during pregnancy has been linked to adverse developmental outcomes in children, including low birthweight and shorter gestation duration, reduced neonatal attention and habituation to stimuli, and increased risk for Attention Deficit Hyperactivity Disorder, schizophrenia, speech impairments, and social abnormalities.<sup>1-6</sup> Primate studies conducted under controlled laboratory conditions have provided an inferential link between rodent studies and human epidemiological studies.

### **Subject**

Increased stress and violence in everyday life has resulted in a growing concern about the effects of maternal stress on child outcomes. This is particularly true in the economically underprivileged sector of our society, where individuals are more likely to be subjected to uncontrollable daily pressures and more serious stressors such as relocation and unemployment. Because women under stress are more likely to smoke cigarettes, consume alcohol, and engage in other stress-related behaviours, there is often a clustering of negative events, which can compound the adverse effects on developmental outcome.

### **Problems**

It is difficult to establish causal links between prenatal stress and developmental outcomes in human studies due to selection bias and possible confounding variables. Even with statistical corrections, it is impossible to ascertain whether major confounders have been eliminated or whether the method of adjustment has effectively removed confounders. In primate studies, one can use randomized experiments, inferring causal connections between prenatal stress on one hand, and offspring behaviour and development on the other. Nonhuman primate studies make an excellent model for studying prenatal stress because of the similarity in brain structure and biological processes of stress reactions to humans between nonhuman primates and humans. Other factors include the richness of nonhuman primate social organization and their complex cognitive capabilities, the opportunity to isolate prenatal stress from other factors and to employ a standard stress treatment, and the ability to investigate possible biological

underpinnings.<sup>7-9</sup> The disadvantages of nonhuman primate studies are that it is necessary to use small sample sizes, cost is high compared to rodent studies, and one must still generalize from animals to humans, albeit this inferential leap is smaller with nonhuman primates than it is with rodents.

### **Research Context**

Stressful events stimulate the release of stress hormones, which can cross the primate placenta (at least in small amounts), and can influence fetal development. Studies with rodents and with nonhuman primates have demonstrated that when dams were injected with stress a hormone (corticotrophin-releasing hormone) (CRH), or when pregnant rhesus monkeys were injected with adrenocorticotrophic hormone (ACTH), their offspring showed effects similar to those observed in prenatally stressed offspring.<sup>10,11</sup>

### **Key research questions**

Our research used daily exposure to 3 short bursts of noise (115 dB sound at 1 m, 1300 Hz) as a prenatal stressor to answer the following questions:

- a) Does unpredictable daily psychological stress during pregnancy have an adverse effect on birthweight, gestation duration, neonatal neurobehaviour and/or stress reactivity?
- b) Is there a sensitive period or time of enhanced vulnerability for prenatal stress effects?
- c) Are maternal stress hormones an important part of the mechanism for the prenatal stress effects on the offspring?
- d) Does prenatal stress interact with other potential negative events, such as fetal alcohol exposure?
- e) Is there continuity between effects observed during infancy and later cognitive functions?

### **Recent research results**

- a) Prenatal stress significantly reduced birthweight, but did not shorten gestation durations, although all birthweights were within the range of what is considered normal for rhesus monkeys.<sup>12</sup> Prenatal stress was also associated with a neurobehavioural profile that included reduced neonatal attention, motor maturity, and motor activity.<sup>13-15,10</sup> When exposed to stressful events, prenatally stressed monkeys showed more disturbance behaviour and higher levels of stress hormones, compared to controls.<sup>16-18,12</sup>
- b) The effects of prenatal stress on birthweight and early neurobehaviour appear to peak during early gestation and taper off during mid-to-late gestation.<sup>14</sup>
- c) Maternal endocrine activation was shown to be one of the underlying mechanisms for the effect, since administering ACTH to pregnant females resulted in effects similar to those observed in prenatally stressed monkeys.<sup>10</sup>
- d) When prenatal stress was combined with fetal alcohol exposure, the most adverse effects on birthweight and behaviour were observed, and dopamine innervation in the striatum was altered compared to controls.<sup>19-21</sup>
- e) There was a significant association between reduced neonatal attention and learning deficits during adolescence in nonmatch-to-sample, showing some continuity between early deficits and later outcomes.<sup>20</sup>

### **Conclusions**

Nonhuman primate studies, conducted under carefully controlled laboratory conditions, are an important link between rodent research and human studies. Primate studies indicate that prenatal stress induces lower birthweights, reduced early attention and motor maturity, slower learning and impaired emotion regulation in offspring. Early gestation appears to be a period of enhanced vulnerability, although effects are also observed from chronic stress during mid-to-late gestation. More research is needed to study the severity of stress, chronicity versus single episodes of stress, timing of stressful events, and the effects of clustering of negative events. In people, coping strategies are very important in ameliorating the impact of stress, and such strategies cannot be studied in animals. Research to date supports a tentative conclusion that there is a causal connection between prenatal stress and developmental outcomes in humans and nonhuman primates.

### **Implications**

Because development is shaped by a highly complex process involving the interplay of biological and environmental factors,<sup>22</sup> it is reasonable to add prenatal stress to the list of risk factors that may adversely affect development, especially when combined with other risk factors. Relevant public policy issues in this area of research include the identification and reduction of existing risk factors and the enhancement of available protective factors in pregnant women. The public needs to be educated about risk factors during pregnancy, including prenatal stress, and the clustering of negative events. Professional training for service providers should include information relevant to risk factors, including stress, and possible effects on offspring.

## REFERENCES

1. Huttunen MO, Niskanen P. Prenatal loss of father and psychiatric disorders. *Archives of General Psychiatry* 1978;35(4):429-431.
2. Lobel M. Conceptualizations, measurement, and effects of prenatal maternal stress on birth outcomes. *Journal of Behavioral Medicine* 1994;17(3):225-272.
3. McIntosh DE, Mulkins RS, Dean RS. Utilization of maternal perinatal risk indicators in the differential diagnosis of ADHD and UADD children. *International Journal of Neuroscience* 1995;81(1-2):35-46.
4. Meijer A. Child psychiatric sequelae of maternal war stress. *Acta Psychiatrica Scandinavica* 1985;72(6):505-511.
5. Oyemade UJ, Cole OJ, Johnson AA, Knight EM, Westney OE, Laryea H, Hill G, Cannon E, Formufod A, Westney LS, Jones S, Edwards CH. Prenatal predictors of performance on the Brazelton Neonatal Behavioral Assessment Scale. *Journal of Nutrition* 1994;124(suppl 6):1000S-1005S.
6. Wadhwa PD. Prenatal stress and life-span development. In: Friedman HS, ed. *Encyclopedia of Mental Health*. Vol. 3. San Diego, CA: Academic Press; 1998:265-280.
7. Goldman-Rakic PS, Brown RM. Postnatal development of monoamine content and synthesis in the cerebral cortex of rhesus monkeys. *Developmental Brain Research* 1982;4(3):339-349.
8. Newell-Morris L, Fahrenbruch CE. Practical and evolutionary considerations for use of the nonhuman primate model in prenatal research. In: Watts ES, ed. *Nonhuman primate models for human growth and development*. New York, NY: A.R. Liss; 1985:9-40.
9. Suomi SJ, Higley JD. Rationale and methodologies for developing nonhuman primate models of prenatal drug exposure. In: Kilby MM, Ashgar K, eds. *Methodological issues in controlled studies on effects of prenatal exposure to drug abuse*. Rockville, Md: National Institute of Drug Abuse; 1991:291-302. *NIDA Research Monograph*; vol. 114.
10. Schneider ML, Coe CL, Lubach GR. Endocrine activation mimics the adverse effects of prenatal stress on the neuromotor development of the infant primate. *Developmental Psychobiology* 1992;25(6):427-439.
11. Williams MT, Hennessy MB, Davis HN. CRF administered to pregnant rats alters offspring behavior and morphology. *Pharmacology, Biochemistry & Behavior* 1995;52(1):161-167.
12. Schneider ML, Moore CF. Effect of prenatal stress on development: A nonhuman primate model. In: Nelson CA, ed. *Minnesota Symposium on Child Psychology*. Vol. 31: Effects of early adversity on neurobehavioral development. Mahwah, NJ: Lawrence Erlbaum Associates Inc; 2000:201-244.
13. Schneider ML. The effect of mild stress during pregnancy on birthweight and neuromotor maturation in rhesus monkey infants (*Macaca mulatta*). *Infant Behavior and Development* 1992;15(4):389-403.
14. Schneider ML, Roughton EC, Koehler AJ, Lubach GR. Growth and development following prenatal stress exposure in primates: An examination of ontogenetic vulnerability. *Child Development* 1999;70(2):263-274.

STRESS (PRENATAL AND PERINATAL)

15. Schneider ML, Coe CL. Repeated social stress during pregnancy impairs neuromotor development of the primate infant. *Journal of Developmental and Behavioral Pediatrics* 1993;14(2):81-87.
16. Clarke AS, Schneider ML. Prenatal stress has long-term effects on behavioral responses to stress in juvenile rhesus monkeys. *Developmental Psychobiology* 1993;26(5):293-304.
17. Clarke AS, Schneider ML. Effects of prenatal stress on behavior in adolescent rhesus monkeys. *Annals of the New York Academy of Sciences* 1997;807:490-491.
18. Schneider ML. Prenatal stress exposure alters postnatal behavioral expression under conditions of novelty challenge in rhesus monkey infants. *Developmental Psychobiology* 1992;25(7):529-540.
19. Schneider ML, Roughton EC, Lubach GR. Moderate alcohol consumption and psychological stress during pregnancy induce attention and neuromotor impairments in primate infants. *Child Development* 1997;68(5):747-759.
20. Schneider ML, Moore CF, Kraemer GW. Moderate alcohol during pregnancy: Learning and behavior in adolescent rhesus monkeys. *Alcoholism: Clinical and Experimental Research* 2001;25(9):1383-1392.
21. Roberts AD, De Jesus OT, Schneider ML, Schueller MJ, Shelton SE, Nickles RJ. Dopamine system characterization of rhesus monkeys exposed to moderate dose alcohol in utero. Society of Nuclear Medicine 46th annual Meeting, Los Angeles, CA. *Journal of Nuclear Medicine* 1999;40(5):108.
22. Sameroff AJ. Developmental systems and psychopathology. *Development and Psychopathology* 2000;12(3):297-312.

To cite this document:

Schneider ML, Moore CF. Prenatal stress and offspring development in nonhuman primates. In: Tremblay RE, Barr RG, Peters RDeV, eds. *Encyclopedia on Early Childhood Development* [online]. Montreal, Quebec: Centre of Excellence for Early Childhood Development; 2003:1-5. Available at: <http://www.child-encyclopedia.com/documents/Schneider-MooreANGxp.pdf>. Accessed [insert date].

Copyright © 2003