



Autism and Its Impact on Child Development

TONY CHARMAN, PhD

Institute of Child Health, UNITED KINGDOM

(Published online September 16, 2005)

(Revised June 1, 2007)

Topic

Autism

Introduction

Autism and related pervasive developmental disorders are more common than previously recognized, affecting up to one in 100 individuals.¹ Although symptoms and accompanying functional impairments can improve with development and as a result of specific interventions, the condition is lifelong and results in considerable morbidity and cost to the individual, his or her family and society.

Subject

The term “autism spectrum disorders” is now commonly used to describe a range of neurodevelopmental conditions that differ in underlying etiology and vary in manifestation.² What they share in common with Kanner’s description of classically “autistic” children is a primary impairment in social relatedness and reciprocity.³ Once considered particular characteristics of rare individuals in the population, these impairments are now more commonly understood as a broad spectrum of individual differences that are widely distributed in the general population.⁴

Problems

The primary impairment in social reciprocity, peer relationships and emotional engagement is accompanied in different individuals by varying degrees of language and communication impairment, mental retardation and patterns of non-functional restricted, repetitive and stereotyped behaviours. Sensory abnormalities, including either hyposensitivity or hypersensitivity, and unusual interests in some sensations are common. A lack of imaginative play indicates an underlying difficulty with generation of ideas that is highly relevant in the development of understanding, and thinking about, other people and other situations. All of these characteristics can be seen in varying degrees of severity.

Both the DSM-IV⁵ and the ICD-10⁶ classification systems include diagnostic categories for individuals who show some but not the full complement of impairments necessary to meet criteria for autism: Asperger’s syndrome, atypical autism and pervasive developmental disorder (“unspecified” in ICD-10; “not otherwise specified” in DSM-IV).

This group is a mix of those with greater severity in one area than others, mild impairments in several areas or late onset (after three years of age).

Research Context

While it has been established that autism is an organic condition and that the condition is highly heritable, in only a minority of cases can a specific neurologic or genetic cause be identified.² As a result, considerable research activity is currently focused on identifying the genetic and neuro-pathological basis. However, as the autism phenotype may be arrived at by a number of pathogenic routes that overlap somewhere in brain development and function, as yet unidentified biological or genetic markers will not necessarily be present in all cases and case definition will continue to be reliant on the behavioural picture alone. Targets for behavioural research include establishing the neuropsychological processes that are impaired, earlier identification and establishment of accurate and reliable diagnosis in toddlers, and testing the efficacy and effectiveness of various intervention programs and approaches.

Key Research Questions and Recent Research Findings

Although etiology can be established in a few cases (e.g. children with fragile-X syndrome or tuberous sclerosis), there is evidence for complex polygenic inheritance. However, while attempts to identify susceptibility genes by association studies have produced several candidate genes on different chromosomes, no gene has yet been identified.⁷ The recurrence risk for subsequently born siblings is approximately 5-10%, although milder impairments of social communication skills or language are found in as many as 20% of relatives. Familial susceptibility to the “broader phenotype” of autism has implications for genetic counselling. Autism is more common in boys than girls (4:1), but no explanation for this discrepancy has been substantiated.⁸ Between 15% and 30% of children with autism experience a period of stasis or regression, most commonly in speech and social behaviour, usually between 12 and 20 months of age, although the causes of this regression are not well understood.⁹

There is agreement that social processing (of faces, emotions, mentalizing skills) is impaired and there is evidence that the brain systems that subserve such cognitive functions are structurally and functionally disrupted. However, the developmental cause of such impairments may be earlier disruptions in the development of brain circuits that underlie the social reward and social orienting systems.^{10,11}

There has been progress in the earlier identification and diagnosis of cases, in part via efforts to develop prospective screening instruments¹² and the prospective study of “high-risk” samples, such as younger siblings of already diagnosed children.¹³ However, this presents clinical challenges: establishing the reliability of early diagnosis, modification of treatment approaches for toddlers, use of assessment tools with younger children and the ability to indicate prognosis.¹⁴

There is some evidence for effectiveness of intensive applied behavioural analysis approaches to early intervention, but there are also limitations with respect to outcomes and the generalization of cue-dependent behaviours.¹⁵ There is also evidence for benefits

from social and communication-based approaches¹⁶ and approaches that provide visual cues and structure that many preschoolers with autism find difficult to generate themselves¹⁷. Important elements of intervention programs for preschool children with autism include a focus on the development of pragmatic and functional communication skills (whether verbal or non-verbal), joint engagement and joint social activities, promotion of emotional engagement and regulation, and helping parents to manage behavioural tantrums and maladaptive routines.¹⁵ For the most treatment-resistant children with the poorest prognosis (e.g. those with no verbal communication by mid-school years, accompanied by extreme social aloofness and mental retardation), there is a need to determine whether augmentative communication approaches might help adaptation.

Depending on family resources and access to support and services, the impact on the family can be considerable, especially at times of important transition (diagnosis, school entry, school transfer, entry to adulthood). Research into the effectiveness and acceptability of support services for families and for adults with autism is scant. One emerging trend is the identification of comorbid psychopathology (e.g. anxiety, OCD) in adolescents and adults, most notable in individuals with average IQs, that can lead to additional challenging behaviour.

Conclusions

Our understanding of autism, once considered a rare and almost always severe childhood disorder, has undergone a revolution in the past 20 years. It is not a rare disorder. Its manifestations can vary widely and it can present in high- as well as in low-IQ individuals. Together with increasing evidence of the positive benefits of early intervention, these changes in conceptualization and application of the diagnosis mean that our notions of outcome and likely progress are also undergoing a revolution. Alongside advances in genetic and neuro-scientific research, a reconceptualization of autism has led scientists to ask fundamental questions regarding social behaviour and communication that have relevance for populations of children in general and not just those relatively rare children who have impairments in these abilities sufficient to meet diagnostic criteria for autism.

Policy Implications

The costs of autism to individuals, families and society are considerable. Internationally, there is an impetus to improve early identification and treatment in order to minimize the impact and to ameliorate the negative secondary sequelae of late diagnosis and ineffective treatments. Community health practitioners and kindergarten staff require training in the identification and management of autism. Basic research into the etiology and underlying psychological impairments that characterize autism are required alongside more applied research into early identification, effective interventions and support for families. At a broader societal level, the recognition that aspects of autism are connected more generally to individual differences in social behaviour (for example, between males and females¹⁸) challenges the notion of autism as a distinct and necessarily “impaired” way of processing and understanding the social world. This calls for greater societal acceptance of differences in social engagement and social behaviour.

To learn more on this topic, consult the following sections of the Encyclopedia:

- [How important is it?](#)
- [What do we know?](#)
- [What can be done?](#)
- [According to experts](#)

REFERENCES

1. Baird G, Simonoff E, Pickles A, Chandler S, Loucas T, Meldrum D, Charman T. Prevalence of disorders of the autism spectrum in a population cohort of children in South Thames: the Special Needs and Autism Project (SNAP). *Lancet* 2006; 368:210-215.
2. Volkmar FR, Lord C, Bailey A, Schultz RT, Klin A. Autism and pervasive developmental disorders. *Journal of Child Psychology and Psychiatry* 2004;45(1):135-170.
3. Kanner L. Autistic disturbances of affective contact. *Nervous Child* 1943;2:217-250.
4. Constantino JN, Todd RD. Autistic traits in the general population: A twin study. *Archives of General Psychiatry* 2003;60(5):524-530.
5. American Psychiatric Association. *Diagnostic and statistical manual of mental disorders (DSM-IV)*. 4th ed. Text revision. Washington, DC: American Psychiatric Association; 2000.
6. World Health Organization. *The ICD-10 classification of mental and behavioural disorders: Diagnostic criteria for research*. Geneva, Switzerland: World Health Organization;1993. Available at: <http://www.who.int/classifications/icd/en/GRNBOOK.pdf>. Accessed May 20, 2005.
7. Folstein SE, Rosen-Sheidley B. Genetics of autism: complex aetiology for a heterogeneous disorder. *Nature Reviews Genetics* 2001;2(12):943-955.
8. Skuse DH. Imprinting, the X-chromosome, and the male brain: explaining sex differences in the liability to autism. *Pediatric Research* 2000;47(1):9-16.
9. Richler J, Luyster R, Risi S, Hsu WL, Dawson G, Bernier R, Dunn M, Hepburn S, Hyman S, McMahon W, Goudie-Nice J, Minshew N, Rogers S, Sigman M, Spence M, Golberg W, Tager-Flusberg H, Volkmar F, Lord C. Is there a 'Regressive Phenotype' of autism spectrum disorder associated with the measles-mumps-rubella vaccine? A CPEA Study. *Journal of Autism and Developmental Disorders* 2006;36(3): 299-316.
10. Dawson G, Webb S, Schellenberg GD, Dager S, Friedman S, Aylward E, Richards T. Defining the broader phenotype of autism: Genetic, brain, and behavioural perspectives. *Development and Psychopathology* 2002;14(3):581-611.

11. Mundy P. The neural basis of social impairments in autism: the role of the dorsal medial-frontal cortex and anterior cingulate system. *Journal of Child Psychology and Psychiatry* 2003;44(6):793-809.
12. Baird G, Charman T, Cox A, Baron-Cohen S, Swettenham J, Wheelwright S, Drew A. Current topic: Screening and surveillance for autism and pervasive developmental disorders. *Archives of Disease in Childhood* 2001;84(6):468-475.
13. Zwaigenbaum L, Thurm A, Stone W, Baranek G, Bryson S, Iverson J, Kau A, Klin A, Lord C, Landa R, Rogers S, Sigman M. Studying the emergence of autism spectrum disorders in high-risk infants: methodological and practical issues. *Journal of Autism and Developmental Disorders* 2006;37(3):466-80.
14. Charman T, Baird G. Practitioner review: Diagnosis of autism spectrum disorder in 2- and 3-year-old children. *Journal of Child Psychology and Psychiatry* 2002;43(3):289-305.
15. Committee on Educational Interventions for Children with Autism, Lord C, McGee JP, eds. *Educating Children with Autism*. Washington, DC: National Academy Press; 2001.
16. Aldred C, Green J, Adams C. A new social communication intervention for children with autism: pilot randomised controlled treatment study suggesting effectiveness. *Journal of Child Psychology and Psychiatry* 2004;45(8):1420-1430.
17. Howlin, P., Gordon K., Pasco G., Wade A, Charman T. The effectiveness of Picture Exchange Communication System (PECS) training for teachers of children with autism: a pragmatic, group randomized controlled trial. *Journal of Child Psychology and Psychiatry* 2007;48(5),473-481.
18. Baron-Cohen S, Knickmeyer RC, Belmonte MK. Sex differences in the brain: implications for explaining autism. *Science* 2005;310(5749):819-23.

To cite this document:

Charman T. Autism and its impact on child development. Rev ed. In: Tremblay RE, Barr RG, Peters RDeV, Boivin M, eds. *Encyclopedia on Early Childhood Development* [online]. Montreal, Quebec: Centre of Excellence for Early Childhood Development; 2007:1-5. Available at: http://www.child-encyclopedia.com/documents/CharmanANGxp_rev.pdf. Accessed [insert date].

Copyright © 2005-2007