

HYPERACTIVITY AND INATTENTION (ADHD)

ADHD and Treatment

¹Meghan Miller, PhD, ²Stephen P. Hinshaw, PhD

¹University of California, Davis, USA

²University of California, Berkeley and University of California, San Francisco, USA

September 2019, Rev. ed.

Introduction

Attention-deficit/hyperactivity disorder (ADHD) is a common and impairing neurodevelopmental disorder that originates in childhood and tends to persist across the lifespan. ADHD is strongly heritable, affects approximately 5-8% of young persons, and occurs more commonly in males than females. As described in the Diagnostic and Statistical Manual of Mental Disorders–Fifth Edition,¹ ADHD consists of symptoms that are developmentally extreme, highly impairing and cross-situationally displayed. Such symptoms fall into two categories: (a) inattention-disorganization and (b) hyperactivity-impulsivity. Individuals who display significant symptoms of inattention-disorganization (but not hyperactivity-impulsivity) are considered to have the Inattentive Presentation; those who display significant symptoms of hyperactivity-impulsivity (but not inattention) are categorized in Hyperactive-Impulsive Presentation. The most common clinical presentation of ADHD is the Combined Presentation, in which the individual displays significant symptoms from both symptom domains. Current evidence-based interventions for ADHD consist of psychotropic medications and behaviour therapy.^{2,3,4,5}

Subject

Determining the most effective intervention strategies for ADHD is highly relevant. It is critically important to determine which components of evidence-based treatments are most effective (including evaluation strategies for making this determination), to identify how treatment strategies can optimally be combined, to determine ways in which to individualize treatments in order to achieve optimal results, to establish the best means of promoting generalization and maintenance of treatment gains, and to determine the factors that contribute to ideal treatment outcomes.

Problems

Years of intervention research have identified and honed evidence-based treatments for ADHD, including

behavioural interventions and medication treatment. Yet such treatments, although evidence-based, are not curative and do not result in significant improvements for all who are treated. Additionally, effects of current evidence-based treatments are not generally long-term and tend not to generalize across settings.

Research Context

The voluminous research on risk factors, correlates, long-term outcomes and underlying processes related to ADHD has still not been fully translated into mechanism-specific interventions. Still, a number of well-controlled single-case reports and randomized, controlled clinical trials attest to the viability of behaviour therapy and medication interventions for ADHD.

Key Research Questions

A key research question involves evaluating the effectiveness of interventions for ADHD, including behavioural, medication and combined behavioural-medication interventions. Additional critical research issues focus on identifying factors that explain how and how well treatments work, and for whom.⁶ Such factors may include sex, ADHD subtype, developmental level, comorbidity, parental factors, medication dosage, cognitive changes, and family discipline styles.^{6,7,8,9,10}

Recent Research Results

Current evidence-based treatments for ADHD include medication and behavioural interventions. Medication treatments for ADHD typically consist of psychostimulants, although other types are often concurrently prescribed in order to address comorbid disorders. Psychostimulants used to address ADHD symptoms include methylphenidate, dextroamphetamine, and mixed amphetamine salts, which all enhance the transmission of dopamine. Atomoxetine, a norepinephrine reuptake inhibitor, has also been found to be effective. Both dopamine and norepinephrine are neurotransmitters (messengers in the brain) that are involved in many mental processes. Whereas such medications have been shown to reduce ADHD-related symptoms and functional impairments across settings,⁴ effects tend to last only as long as the medication is active within the body and brain.

As a result, and to promote active skill-building, non-medication treatments are often recommended as well. Behaviour therapy is the only consistently evidence-based intervention for childhood ADHD aside from medication. Behavioural treatments typically involve interventions with parents, teachers, and the child.^{2,11} Specific components of behavioural interventions for ADHD include direct contingency management and clinical behaviour therapy. Direct contingency management consists of teachers or counselors directly rewarding target skills and employing effective consequences when problems arise. More commonly-used clinical behaviour therapy procedures involve (1) parent training on topics including behaviour management (e.g., positive parental attention, rewards for appropriate behaviour, negative consequences for misbehaviour) and (2) teacher training on topics such as use of prompts and rewards in the classroom. Across most relevant investigations, the greatest likelihood of symptom reduction occurs when medication and behavioural treatments are combined, particularly with respect to functional outcomes.^{12,13,14}

Although evidence-based interventions for ADHD have been identified, few randomized controlled trials have

focused on identifying specific individual factors that influence treatment outcome. Key factors emerging from the Multimodal Treatment of children with ADHD (MTA) Study included the presence of a comorbid anxiety disorder, family public assistance, ethnicity/race, severity of ADHD, parental depressive symptomatology, child IQ, attendance, medication use in the community and negative/ineffective parental discipline.⁶

Finally, cognitive enhancements of contingency-based interventions (e.g., social skills training with parent training) as well as cognitive training to ameliorate neuropsychological deficits commonly associated with ADHD (i.e., executive function deficits, which include difficulties planning, staying organized, inhibiting inappropriate responses, setting and carrying out goals) may be viable but evidence is currently limited. A critical issue with the current evidence-based behavioural interventions for ADHD – including combination treatments of medication and behavioural interventions – is that treatment gains often are not maintained over periods of time, nor are gains generalized across settings.³

Research Gaps

A major issue with current evidence-based treatments for ADHD concerns generalization.¹⁶ Specifically, individuals with ADHD tend not to translate gains obtained in one setting to other key life settings. Thus, future treatment development efforts should focus on determining components of interventions that promote long-lasting maintenance of treatment gains across settings. An additional issue involves developing treatments that can address functional impairments that often accompany ADHD (e.g., social difficulties, academic struggles, organizational skills deficits). Successful interventions in these domains may have significant long-term effects, with the potential for contributing to the elusive goal of maintaining gains. Finally, a critical gap in the literature pertains to early, preventive interventions. Such interventions, if successful, could alter the trajectory of ADHD from the start, preventing later problems. Although this approach is being used in other neurodevelopmental disorders (e.g., autism spectrum disorder), such methodology has rarely been used in the study of ADHD.¹⁷ One key problem relates to false-positives: how many preschool-aged children at risk for ADHD will actually proceed to the full disorder?

Conclusions

ADHD is a common and impairing neurodevelopmental disorder that requires intensive intervention. Much research has focused on identifying evidence-based interventions for ADHD. Current evidence-based treatment options include behavioural interventions and medication treatment. Medication treatment provides the greatest reduction of core ADHD symptoms (inattention, hyperactivity-impulsivity) whereas evidence exists that the combination of medication and behavioural interventions results in the greatest improvement in associated impairments (e.g., parent-child relations, academic problems, anxiety). Although these treatment options are useful for decreasing core symptomatology, they do not appear to remediate core deficits related to ADHD, and they tend not to produce long-term, generalized gains. It will be essential to promote translational research linking biological and contextual risk factors to development of improved treatment strategies. Important areas for future research include identifying specific factors that influence treatment outcome, developing of interventions that produce effects that can be generalized and maintained over time, determining ways to address the functional impairments commonly present in individuals with ADHD, and ascertainment of the potential benefits of preventive measures.

Implications for Parents, Services and Policy

Professional organizations (e.g., American Association of Child and Adolescent Psychiatry and American Academy of Pediatrics)^{18,19} have published assessment and treatment guidelines for ADHD, but there is little or no enforcement of such professional guides, nor is it clear that reimbursement always or even usually covers such standards. Families need to be aware of the need for relevant health professionals to have demonstrated expertise in ADHD and its common comorbidities and impairments; policy-makers need to assure adequate assessment and treatment standards. Furthermore, large regional variation in rates of diagnosis and treatment of ADHD exists within the U.S.²⁰ and internationally.²¹ Policy-related factors such as high-stakes achievement testing, training of professionals, insurance coverage, and advertisements for treatment (particularly medication) may all be relevant regarding this wide variation. Overall, knowledge of ADHD, reduction of stigma regarding its identification and treatment, enlightened policies that provide for evidence-based services, and proper means of assessing treatment-related gains are essential goals if youth and their families are to receive optimal services.²²

References

1. American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 5th ed. Washington, DC: Author; 2013.
2. Pfiffner LJ, Haack LM. Nonpharmacologic treatments for childhood ADHD and their combination with medication. In: Nathan PE, Gorman JM, eds. A guide to treatments that work, 4th ed. New York: Oxford University Press; 2015:55-84.
3. Miller M, Hinshaw SP. Treatment for children and adolescents with ADHD. In: Kendall PC, ed. Child and Adolescent Therapy, 4th Edition: Cognitive-Behavioral Procedures. New York: Guilford Press; 2011:61-91.
4. Kass E, Posner J, Greenhill LL. Pharmacological treatments for attention-deficit/ hyperactivity disorder and disruptive behavior disorders. In: Nathan PE, Gorman JM, eds. A guide to treatments that work, 4th ed. New York: Oxford University Press; 2015:85-140.
5. Evans SW, Hoza B, eds. Treating attention deficit hyperactivity disorder: Assessment and intervention in developmental context. Kingston, NJ: Civic Research Institute; 2011.
6. Hinshaw SP. Moderators and mediators of treatment outcome for youth with ADHD: Understanding for whom and how interventions work. *Journal of Pediatric Psychology*. 2007;32(6):664-675.
7. Jensen PS, Hinshaw SP, Kraemer HC, Lenora N, Abikoff HB, Conners CK, et al. ADHD comorbidity findings from the MTA study: Comparing comorbid subgroups. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2001;40(2):147-158.
8. MTA Cooperative Group. Moderators and mediators of treatment response for children with ADHD: The MTA Study. *Archives of General Psychiatry*. 1999;56(12):1088-1096.
9. Owens EB, Hinshaw SP, Kraemer HC, Arnold LE, Abikoff HB, Cantwell DP, et al. Which treatment for whom for ADHD? Moderators of treatment response in the MTA. *Journal of Consulting and Clinical Psychology*. 2003;71(3):540-552.
10. Kazdin AE, Weisz JR. Identifying and developing empirically supported child and adolescent treatments. *Journal of Consulting and Clinical Psychology*. 1998;66(1):19-36.
11. Evans SW, Owens JS, Wymbs BT, Ray AR. Evidence-based psychosocial treatments for attention-deficit/hyperactivity disorder. *Journal of Clinical Child & Adolescent Psychology*. 2018;47(2):157-198.
12. Conners CK, Epstein JN, March JS, Angold A, Wells KC, Klaric J, et al. Multimodal treatment of ADHD in the MTA: An alternative outcome analysis. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2001;40(2):159-167.
13. Swanson JM, Kraemer HC, Hinshaw SP, Arnold LE, Conners CK, Abikoff HB, et al. Clinical relevance of the primary findings of the MTA: Success rates based on severity of ADHD and ODD symptoms at the end of treatment. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2001;40(2):168-179.
14. MTA Cooperative Group. A 14-month randomized clinical trial of treatment strategies for attention-deficit/hyperactivity disorder. *Archives of General Psychiatry*. 1999;56(12):1073-1086.
15. Hinshaw SP, Owens EB, Wells KC, Kraemer HC, Abikoff HB, Arnold LE, et al. Family processes and treatment outcome in the MTA: Negative/ineffective parenting practices in relation to multimodal treatment. *Journal of Abnormal Child Psychology*. 2000;28:555-568.
16. Abikoff H. ADHD psychosocial treatments: Generalization reconsidered. *Journal of Attention Disorders*. 2009;13(3):207-210.

17. Sonuga-Barke EJS, Halperin JM. Developmental phenotypes and causal pathways in attention deficit/hyperactivity disorder: Potential targets for early intervention? *Journal of Child Psychology and Psychiatry*. 2010;51(4):368-389.
18. American Association of Child and Adolescent Psychiatry. Practice parameter for the assessment and treatment of children and adolescents with attention-deficit/hyperactivity disorder. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2007;46(7):894-921.
19. American Academy of Pediatrics. ADHD: Clinical practice guidelines for the diagnosis, evaluation, and treatment of attention-deficit/hyperactivity disorder in children and adolescents. *Pediatrics*. 2011;128(5):1007-1022.
20. Fulton BD, Scheffler RM, Hinshaw SP, Levine P, Stone S, Brown TT, et al. National variation of ADHD diagnostic prevalence and medication use: Health care providers and educational policies. *Psychiatric Services*. 2009;60(8):1075-1083.
21. Hinshaw SP, Scheffler RM, Fulton BD, Aase H, Banaschewski T, Cheng W, et al. International variation in treatment procedures for ADHD: Social contexts and recent trends. *Psychiatric Services*. 2011;62(5):459-464.
22. Hinshaw SP. Attention deficit-hyperactivity disorder (ADHD): Controversy, developmental mechanisms, and multiple levels of analysis. *Annual Review of Clinical Psychology*. 2018;14:291-316.