



# **Child nutrition**

Last update: October 2025

**Topic Editor:** 

Myles S. Faith, PhD, University of North Carolina at Chapel Hill, USA

# **Table of content**

Synthesis	4
Feeding skill, Appetite and Feeding Behaviours of Infants and Young Children and Their Impact on Growth and Psychosocial Development MARIA RAMSAY, PHD, SEPTEMBER 2025	7
Feeding Behaviour of Infants and Young Children and Its Impact on Child Psychosocial and Emotional Development YI HUI LIU, MD MPH, MARTIN T. STEIN, MD, SEPTEMBER 2013	21
Helping Children Develop Healthy Eating Habits MAUREEN M. BLACK, PHD, KRISTEN M. HURLEY, PHD, SEPTEMBER 2013	28
Assessment and Treatment of Pediatric Feeding Disorders  ¹SUZANNE M. MILNES, PHD, ¹CATHLEEN C. PIAZZA, PHD, ²TAMMY CARROLL, PHD, SEPTEMBER 2013	38
Genetic Influences on Child Eating Behaviour CLARE LLEWELLYN, PHD, JANE WARDLE, PHD, SEPTEMBER 2013	 43
Screen time Use and Food Marketing Influences on Children's Eating Behaviors  ¹CHRISA ARCAN, PHD, MPH, RD, ²ANINDA SEN, MBBS, MPH, ³MEG BRUENING, PHD, MPH, RD, ⁴MARY STORY, PHD, RD, OCTOBER 2025	50
Eating Behaviour in Studies of Child Growth, Development and Health – Measure Precisely, Early and in Context: Commentary on Ramsay, Liu & Stein, Black & Hurley, Milnes, Piazza & Carrol, Llewellyn & Wardle, and Arcan, Bruening and Story  MYLES S. FAITH, PHD, SEPTEMBER 2013	67

### **Synthesis**

#### How important is it?

Oral motor skills such as sucking and chewing develop rapidly during the first year, allowing young children to discover an increasing number of foods and textures. Refinements in fine motor skills also allow infants to develop more autonomy in regards to feeding, and by the end of the second year, most children have acquired good feeding skills. At this age, children become increasingly influenced by external signals such as family, friends, and society, to dictate their hunger.

Feeding problems are common, and touch approximately 25% to 50% of young children. These problems are usually minor and temporary. They tend to be seen when children are exposed to new foods or events during mealtime, or when they are trying to master a new feeding skill. However, 1% to 2%of children experience chronic feeding problems, including overeating, malnutrition, problematic behaviours during meals and atypical eating choices. These problems are especially common in children born prematurely, and those with developmental disabilities. Eating problems can be highly stressful for parents and can result in a strained parent-child relationship.

#### What do we know?

Early feeding experiences can have long-lasting consequences on eating habits and food preferences later on in life. A healthy diet consisting of ample fruits and vegetables in the first two years may increase future preferences for healthy food. In contrast, unhealthy habits, such as favoring food that is high in fat and sugar tend to persist over time.

Feeding problems are associated with a number of poor developmental, psychological and health outcomes. For instance, children who overeat are more likely to develop medical conditions and potential psychological issues, whereas insufficient calorie intake can result in stunted growth.

Several factors play a role in the development of feeding problems. Feeding difficulties are likely to emerge when feeding has been paired with a painful or unpleasant experience that the child comes to associate with mealtime. Temperament traits such as the inability to self-regulate and

to communicate one's hunger level can make feeding routine challenging. Similarly, a genetic predisposition for big or small appetite can increase the likelihood of feeding problems.

Parents play a central role during feeding throughout the early years. Breastfeeding, for instance, not only protects against contagious diseases, but also allows infants to develop the ability to control their own calorie intake, an important healthy eating habit. Parents also contribute positively to their child's feeding by modeling healthy eating habits and making mealtimes enjoyable.

When feeding problems emerge, mealtimes can become loaded with anxiety and frustration for both children and parents. Misreading the child's signals can further aggravate problematic behaviours. Strategies to encourage eating such as punishment, distractions, and toys can work temporarily, but they tend to worsen problems over time. The most effective approach is responsive feeding, where reciprocal interactions during meals are based on the child's signals, and are age-appropriate.

At a larger scale, culture also contributes to feeding experiences by its influence on feeding choices, behaviours and exposure to different types of food. For instance, children's exposure to advertisements of unhealthy food is related to their preference for this type of food, and this is especially problematic in the African American and Hispanic American communities.

#### What can be done?

Given that feeding problems most often originate from multiple causes, multidisciplinary approaches that consider cultural and temperamental differences are strongly recommended. Although not all children respond equally well to interventions, several strategies have been successful at promoting a healthy diet, starting with those involving parents. Parents can help reduce overeating by cooking smaller quantity of food per meal, thereby limiting access to oversized portion, and by encouraging their children to slow down their eating speed by putting their utensils down after each bite. Parents can also encourage healthy habits by discussing feelings of satiety, modelling healthy eating habits, and limiting the amount of television watching to 1 to 2 hours a day of quality programming, and only in children over 2 years. Parents who discuss the content of advertisement messages and who regulate the type of food their young child eats are likely to be successful at discouraging unhealthy habits.

In cases of insufficient calorie intake, medications that increase appetite have been shown to positively affect weight gain, and can make psychosocial intervention more efficient. The acceptance of new foods can also be encouraged by pairing the new food with a preferred food until the new food becomes familiar. To make mealtimes pleasant, children should be given enough time to eat (between 10 to 30 minutes) and proper equipment (ex., highchairs, and small utensils). Feeding is an intricate skill that takes time to develop. Parents and professionals can be about the specific impact of problematic feeding behaviour and motivation. Information about healthy habits can be transmitted to parents during regular health check-ups.

At the policy level, several steps can be undertaken to reduce accessibility to unhealthy foods. These include the elimination of vending machines in schools, stricter advertising and menu regulations, quotas assigned to the number of fast food restaurants and changes in supermarket layouts.

# Feeding skill, Appetite and Feeding Behaviours of Infants and Young Children and Their Impact on Growth and Psychosocial Development

Maria Ramsay, PhD

McGill University, Canada

September 2025, 2e éd. rév.

#### Introduction

Feeding, like other sensorimotor skills, is a developmental skill that matures during the first two years of life. It is a highly complex sensorimotor process with developmental stages based on neurological maturation and experiential learning. However, feeding, unlike other sensori-motor skills, is heavily reliant on internal incentive or motivation to initiate ingestion, and is essential for survival of the newborn. Thus, the act of feeding is highly charged emotionally for the mother, whose primary responsibility, as viewed by the family, society and culture around her, is to ensure the early growth and well being of her child. Therefore, from the very beginning the mother-infant feeding relationship is influenced by physiologic as well as interactional forces at multiple levels. <sup>2</sup>

#### Subject

When feeding skills are intact and appetite is robust, feeding times, and later on, mealtimes are a source of pleasant socialisation resulting in adequate nutrient intake and good growth. Demanding food at regular intervals, sucking, eating and drinking with good rhythm, trying new food tastes or textures, and expressing satisfaction at the end of feeding are all considered good feeding behaviours by family and society. These pro-feeding behaviours invite praise and positive feeding interactions and thus reinforce the feeling of self-mastery in the young child and promote continued food acceptance and independent feeding behaviours.

However, when feeding skills are impaired (e.g.: poor oral-motor skills, taste and texture sensitivities) and or poor appetite (inadequate hunger), they manifest themselves in problematic feeding behaviours such as not signalling hunger, sucking or eating excessively slowly, gagging, and not bringing food to the mouth.<sup>3-7</sup> In addition, associative conditioning to painful

gastrointestinal cues is particularly powerful in young infants and this conditioning often manifests itself in problematic feeding behaviours.<sup>8,9</sup> Temperamental characteristics and regulatory capacities of the infant may further modulate feeding behaviours.<sup>10,11</sup> Maternal attempts at increasing her infant's nutrient intake by feeding more frequently or longer duration tend to result in stressful feeding experiences for both.<sup>12</sup> While these efforts may work well initially for maintaining good weight gain, they tend to become ineffective and maladjusted mealtime interactions and behavioural mismanagement prevail.<sup>2,13,15</sup> Maternal and family characteristics and societal expectations about the size of the young child and the type of food eaten further influence an already stressful feeding relationship.<sup>16,17</sup>

#### **Problems**

Feeding difficulties are one of the most common developmental disturbances in otherwise healthy infants and young children, often resulting in poor growth. Although an estimated 25%-50% of children experience transient feeding problems under two years of age, 18,19 most feeding issues resolve by the end of early childhood. However, an estimated 3-10% of children present with more severe forms of feeding problems which put children at risk for impaired growth, chronic illnesses and behavioural developmental problems.<sup>20</sup> As well, a large percentage of children with special needs, children with developmental disabilities and children born prematurely have severe and chronic feeding problems where families need support in resolving the feeding issues.<sup>21-23</sup> At the clinical level, the mother (and her paediatrician) is often not aware of the underlying reasons for problematic feeding behaviours. Thus, the mother's reactions to a poor feeder may be exposed to covert or overt family criticism, which often lead to internal doubt about her own ability to nurture.<sup>4</sup> At the policy level, there is a lack of education of professionals and young parents about feeding as a highly complex developmental skill, motivated by hunger and conditioned by parental reactions. Furthermore, professionals are still not trained to recognize that when either feeding skills or motivation or both are impaired, problematic feeding behaviours, stressed mealtime interactions and family conflicts are likely to result.

#### Research Context

Earlier cross-sectional clinical studies examined the relationship between feeding difficulties and attachment, maternal characteristics, family dynamics<sup>24</sup> and feeding practices.<sup>24-30</sup> These studies were conducted prospectively, that is, after the children were diagnosed with poor growth. Several observational studies focused on feeding interactions and problematic feeding

behaviours.<sup>31,32</sup> The development of feeding and patterns of food acceptance have been studied by numerous psychologists.<sup>33-36</sup> More recently, few researchers started to focus on possible pathophysiology (heart rate variability, hormonal balance) of poor growth and problematic feeding behaviours.<sup>37-39</sup> Other studies were conducted in the context of primarily behavioural interventions for problematic feeding behaviours in medically ill and very premature infants.<sup>40-43</sup>

#### **Key Research Questions**

The extensive research in the area of feeding problems and poor growth can be divided along the following three research questions:

- 1. How do maternal (family) characteristics (cognitive abilities, personality disorders, psychological status and early attachment history) influence feeding behaviours and growth?
- 2. How do infant characteristics (feeding skills, appetite, temperament and other physiological characteristics) influence feeding behaviours, mealtime interactions and growth?
- 3. How effective are behavioural and other forms of intervention for severe problematic feeding behaviours in medically ill infants?

#### **Recent Research Results**

Only questions 2 and 3 will be summarised here. With a focus on infant characteristics, studies have shown that feeding problems often co-occur with sleeping and behavioural disturbances (irritability, poor self-calming and intolerance to change), suggesting that these are symptoms of a common underlying constitutional "regulatory disorder" in infants and young children.<sup>44,45</sup> In a large whole-population survey of children's growth and development, a significant portion (36%) of the 47 children identified with failure to thrive at one year of age were found to have oral motor difficulties, suggesting that these children were biologically more vulnerable to poor eating from birth.<sup>46</sup> Another study showed that young infants with gastroesophageal reflux were significantly more likely to have delay in their feeding skills and readiness behaviour for solids than controls.<sup>47</sup> In a prospective study of a group of healthy term infants (n=330), infants with inefficient sucking, as measured by tracings on a polygraph, at one week and two months were significantly more likely to have mothers with greater effort at feeding than infants with efficient sucking.<sup>44</sup>

A number of studies have shown that children under 3 to 4 years of age eat primarily in response to appetite or hunger cues, whereas older children's eating are influenced by a variety of environmental (extra food available) and social factors. 48,49 As well, children with poor growth were observed to refuse offered food more often and fed themselves significantly less often than controls. 50 In terms of the third question, the literature reflects the reality that presently we are better at identifying factors contributing to feeding problems at any level of severity than treating them. 28,51,52 Treating feeding problems at the primary or secondary level, while desirable, is not always available for parents. 53 Treating feeding disorders associated with severe medical illnesses, developmental disabilities and gavage feedings requires the collaboration of multidisciplinary teams for successful outcome. 54,59 Lastly, studies have shown that appetite stimulating medications result in good weight gain, and thus making intervention more efficacious. 60,62

#### **Conclusions**

Understanding feeding behaviours requires the knowledge of feeding as a developmental skill that matures over time and is reliant on hunger (appetite) cues and experiential learning. Whereas feeding skills are well established by two years of age, hunger cues shift from primarily internal to external (family, school and societal) control only by 4-5 years. Thus, although initially problematic feeding behaviours tend to be reactions to internal cues, these behaviours can become conditioned to external (coaxing parents) and societal cues. Medical illnesses, prematurity and developmental disorders further interfere with the development of normal feeding behaviours.

In order to help identify feeding problem, a number of feeding scales have been devised, 63-65 but rarely used for assessment or treatment outcome. Yet, early behavioural intervention can play an important role in normalizing feeding behaviours and mealtime interactions, which in turn help promote independence and other self-help skills in the child. Most recently, an easy and short screening tool was developed for detecting problematic feeding behaviours in primary care offices, allowing early referral to appropriate feeding clinics. 66

An example of a Feeding Scale, which takes only 5 minutes to fill and may be filled by a pediatrician or nurse or other professionals, given that they possess the necessary information about infant/child feeding, or know where to refer the child with feeding problems.


#### **Implications for Policy and Services**

The major findings from this updated summary continue to be that the physiological make up of the infant, medical illnesses, developmental disabilities and parental responses all play an important role in the dynamic relationship in which problematic feeding behaviours develop. This finding has several implications for policy and services in hospitals where premature infants and infants with medical needs grow poorly, requiring enteral feedings. Thus, intervention with staff specialized in feeding disorders, usually occupational or speech therapists, help initiate oral feedings. As well, doctors or pediatricians following infants and young children need continued education in possible causes of poor growth in otherwise healthy infants.

- 1. At the primary care level, the use of easy to administer feeding scales for earlier detection and thus, treatment of feeding problems should continue to be advocated. (See an example of the Feeding Scale below).
- 2. The continuation of mandated multidisciplinary feeding clinics addressing feeding disorders in hospital settings. These feeding programs need to be easily accessible to parents, where effective behavioural intervention and preventive strategies may be implemented in the early stages of reported difficult feeding behaviours.
- 3. The training of experts in the field of feeding disorders, which should include training in the behavioural, developmental and interactional components of feeding, as well as understanding the physiological, as well as the medical components should be advocated.

#### The Pediatric Feeding Program' research at The Montreal Children's Hospital:

Feeding infants and young children can be challenging. Initially, the feeding scale was developed to have a common language with the mothers about feeding difficulties they experience with their children, and to assess the intervention. This scale (MCH-FS) has become known wordwide, and as of now, it has been translated and standardized on a healthy children population in Montreal and in several other countries. The author of this article has approved all translations by using back-translations before the translated version could be used in 8 languages. Depicted in the graphic below is only 7 as Canada used the results of the English and French languages combined. The scale was later translated and standardized in another four languages: Italian Mexican Turkish Turkish and Chinese.



See the MCH-FS above: Of note is that other infant/child feeding scales do not take into consideration the variability of infant/child level of appetite, which influences maternal approach to feeding and the child's acceptance of food.<sup>66-77</sup>

#### References

- Stevenson RD, Allaire JH. The development of normal feeding and swallowing.
   Development and behavior: The very young child. *Pediatric Clinics of North America*. 1991;38(6):1439-1453.
- 2. Ramsay M. Les problèmes alimentaires chez les bébés et les jeunes enfants : Une nouvelle perspective. *PRISME*. 1999;30:10-27.
- 3. Milla PJ. Feeding, tasting and sucking. In: Walker WA, Durie P, Hamilton J, Walker-Smith J, Watkins J, eds. *Pediatric gastrointestinal disease: pathophysiology, diagnosis, management*. Philadelphia, PA: DC Decker; 1991:217-223.
- 4. Ramsay M. Feeding disorder and failure to thrive. *Child and Adolescent Psychiatric Clinics of North America*. 1995;4(3):605-616.
- 5. Reau NR, Senturia YD, Lebailly SA, Christoffel KK. Infant and toddler feeding patterns and problems. Normative data and a new direction. *Journal of Developmental and Behavioral Pediatrics*. 1996;17(3):149-153.
- 6. Jacobi C, Agras WS, Bryson S. Hammer LD. Behavioral validation, precursors and concomitants of picky eating in childhood. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2003;42(1):76-84.
- 7. Ramsay M, Gisel EG, Boutry M. Nonorganic failure to thrive: A growth failure secondary to feeding skills disorder. *Developmental Medicine and Child Neurology*. 1993;35(4):285-297.
- 8. Davidson TL. Pavlovian occasion setting: a link between physiological change and appetitive behavior. *Appetite*. 2000;35(3):271-272.
- 9. Hamilton AB, Zeltzer LK. Visceral pain in infants. *Journal of Pediatrics*. 1994;125(6 Pt 2):S95-102.
- 10. Harris G, Blissett J, Johnson R. Food Refusal Associated with Illness. *Child Psychology and Psychiatry Review*. 2000;5(4):148-156.

- 11. DeGangi GA, Porges SW, Sickel RZ, Greenspan SI. Four-year follow-up of a sample of regulatory disordered infants. *Infant Mental Health Journal*. 1993;14(4):330-343.
- 12. Ferguson A, Blaymore Bier LA, Cucca J, Andereozzi L, Lester B. The quality of sucking in infants with colic. *Infant Mental Health Journal*. 1995;17(2):161-169.
- 13. Ramsay M, Gisel EG. Neonatal sucking and maternal feeding practices. *Developmental Medicine and Child Neurology*.1996;38(1):34-47.
- 14. Ramsay M, Gisel E, McCusker J, Bellevance F, Platt R. Infant sucking ability, nonorganic failure to thrive, maternal characteristics and feeding practices: A prospective cohort study. *Developmental Medicine and Child Neurology*. 2002;44(6):405-414.
- 15. Budd KS, McGraw TE, Farbisz R, Murphy TB, Hawkins D, Heilman N, Werle M, Hochstadt NJ. Psychosocial concomitants of children's feeding disorders. *Journal of Pediatric Psychology*. 1992;17(1):81-94.
- 16. Farrow C, Blissett J. Maternal cognitions, psychopathologic symptoms and infants temperament as predictors of early infant feeding problems: A longitudinal study. *International Journal of Eating Disorders*. 2006;39(2):128-134.
- 17. Knaapila A, Tuorila H, Silventoinen K, Keskitalo K, Kallela M, Wessman M, Peltonen L, Cherkas LF, Spector TD, Perola M. Food neophobia shows heritable variation in humans. *Physiology & Behavior*. 2007;91(5):573-578.
- 18. McDermott BM, Mamun AA, Najman JM, Williams GM, O'callaghan MJ, Bor W. Preschool children perceived by mothers as irregular eaters: Physical and psychosocial predictors of a birth cohort study. *Journal of Developmental and Behavioral Pediatrics*. 2008: 29(3):197-205.
- 19. Carruth BR, Zeigler PJ, Gordon A, Barr SI. Prevalence of picky eaters among infants and toddlers and their caregivers' decisions about offering a new food. *Journal of the American Dietetic Association*. 2004;104(1 Suppl 1):S57-S64.
- 20. Corbett SS, Drewett RF. To what extent is failure to thrive in infancy is associated with poorer cognitive development. A review and meta-analysis. *Journal of Child Psychology and Psychiatry, and Allied Disciplines*. 2004;45(3):641-654.
- 21. Lenscheid TR. Behavioral treatments for pediatric feeding disorders. *Behavior Modification* . 2006;30(1):6-23.

- 22. SJ, Harris G, Blissett J. Tube feeding in infancy: Implications for the development of normal eating and drinking skills. *Dysphagia*. 2005;20(1):46-61.
- 23. Cerro N, Zeunert S, Simmer KN, Daniels LA. Eating behaviour of children 1.5-3.5 years born preterm: Parents' perceptions. *Journal of Paediatrics and Child Health*. 2002;38(1):72-78.
- 24. Piwoz EG, Black RE, Lopez de Romana G, Creed de Kanashiro H, Brown KH. The relationship between infants' preceding appetite, illness, and growth performance and mothers' subsequent feeding practice decision. *Social Science & Medicine*. 1994;39(6): 851-860.
- 25. Benoit D. Failure to thrive and feeding disorders. In: Zeanah CH Jr, ed. *Handbook of infant mental health*. New York, NY: Guilford Press; 1993:317-331.
- Galler JR, Harrison RH, Biggs MA, Ramsey F, Forde V. Maternal moods predict breastfeeding in Barbados. *Journal of Developmental and Behavioral Pediatrics* .1999;20(2):80-87.
- 27. Polan HJ, Kaplan MD, Kessler DB, Schindledecker R, Mewmark M, Stern D, Ward MJ. Psychopathology in mothers of infants with failure to thrive. *Infant Mental Health Journal*. 1991;12(1):55-64.
- 28. Ward MJ, Kessler DB, Altman SC. Infant attachment in children with failure to thrive. *Infant Mental Health Journal*. 1993;14(3):208-220.
- 29. Lindberg L, Bohlin G, Hagekull B, Thurnström M. Early food refusal: Infant and family characteristics. *Infant Mental Health Journal*. 1994;15(3):262-277.
- 30. Parkinson NK, Drewett RF. Feeding behaviour in the weaning period. *Journal of Child Psychology and Psychiatry*. 2001;42(7):971-978.
- 31. Lindberg L, Bohlin G, Hagekull B. Interactions between mothers and infants showing food refusal. *Infant Mental Health Journal*. 1996;17(4):334-347.
- 32. Stein A, Woolley H, Cooper SD, Fairburn CG. An observational study of mothers with eating disorders and their infants. *Journal of Child Psychology and Psychiatry*. 1994;35(5):733-748.
- 33. Birch LL, Gunder L, Grimm-Thomas, Laong DG. Infants' consumption of a new food enhances acceptance of similar foods. *Appetite*. 1998;30(3):283-295.

- 34. Birch LL. Development of food acceptance patterns. *Developmental Psychology*. 1990;26(5):515-519.
- 35. Beauchamp GK, Mennella JA. Flavor perception in human infants: Developmental and functional significance. *Digestion*. 2001;63 Suppl 1:1-6.
- 36. Coulthard H, Harris G, Emmett P. Delayed introduction of lumpy foods to children during the complementary feeding period affects child's food acceptance and feeding at 7 years of age. *Maternal and Child Nutrition*. 2009;5(1):75-85.
- 37. Shaoul R, Kessel A, Toub, E, Lanir A, Glazer O, Jaffe M. Leptin and cytokines levels in children with failure to thrive. *Journal of Pediatric Gastroenterology and Nutrition*. 2003;37(4):487-491.
- 38. Steward DK, Moser DK, Ryan-Wenger NA. Biobehavioral characteristics of infants with failure to thrive. *Journal of Pediatric Nursing*. 2001;16(3)162-171.
- 39. Tannenbaum GS, Ramsay M, Martel C, Samia M, Zygmuntowicz C, Porporino M, Ghosh S. Elevated circulating acylated and total ghrelin concentrations along with reduced appetite scores in with failure to thrive. *Pediatric Research*. 2009;65(5):569-573.
- 40. Babbitt RA, Hoch RA, Coe DA, Cataldo MF, Kelly KJ, Stackhouse C, Perman JA. Behavioral assessment and treatment of pediatric feeding disorders. *Journal of Developmental and Behavioral Pediatrics*. 1994;15(4):278-291.
- 41. Burklow KA, Phelps AN, Schultz JR, McConnell, Rudolph C: Classifying complex pediatric feeding disorders. *Journal of Pediatric Gastroenterology and Nutrition*. 1998;27(2):143-147.
- 42. Werle MA, Murphy TB, Budd KS. Treating chronic food refusal in young children: Homebased parent training. *Journal of Applied Behavior Analysis*. 1993;26(3):421-433.
- 43. Burmucic K, Trabi T, Deutschmann A, Scheer PJ, Dunitz-Scheer M. Inpatient tube weaning in children with long-term feeding tube dependency: A retrospective analysis. *Infant Mental Health Journal*. 2010;31(6):664-681
- 44. St.James-Roberts I, Plewis I. Individual differences, daily fluctuations, and developmental changes in amounts of infant waking, fussing, crying, feeding, and sleeping. *Child Development*. 1996;67(6):2527-2540.
- 45. Wolke D, Gray P, Meyer R. Excessive infant crying: A controlled study of mothers helping mothers. *Pediatrics*. 1994;94(3):322-334.

- 46. Reilly SM, Skuse DH, Wolke D, Stevenson J. Oral-motor dysfunction in children who fail to thrive: organic or non-organic. *Developmental Medicine and Child Neurology*. 1999;41(2):115-122.
- 47. Mathisen B, Worrall L, Mase; J, Wall C, Shepherd RW. Feeding problems in infants with gastro-oesophageal reflux disease: a controlled study. *Journal of Paediatrics and Child Health*. 1999;35(2):163-169.
- 48. Fisher JO, Birch LL. Eating in the absence of hunger and overweight in girls from 5 to 7 year of age. *American Journal of Clinical Nutrition*. 2000;76(1):226-231.
- 49. Rolls BJ, Engell D, Birch LL. Serving portion size influences 5-year-old but not 3-year-old children's food intakes. *Journal of the American Dietetic Association*. 2000;100(2):232-234.
- 50. Drewett RF, Mambwe K-H, Wright C. Feeding behaviour in young children who fail to thrive. *Appetite*. 2003;40(1):55-60.
- 51. Wright C, Birks E. Risk factors for failure to thrive: A population-based survey. *Child: Care, Health and Development.* 2000;26(1):5-16.
- 52. Field D, Garland M, Wiliams K. Correlates of specific childhood feeding problems. *Journal of Paediatrics and Child Health*. 2002;39(4):299-304.
- 53. Hofacker NV, Papousek M. Disorders of excessive crying, feeding and sleeping: The Munich interdisciplinary research and intervention program. *Infant Mental Health Journal*. 1998;9(2):180-201.
- 54. Williams KE, Field DG, Seiverling L. Food refusal in children: review of the literature. *Research in Developmental Disabilities*. 2010;31(3):625-633.
- 55. Mason SJ, Harris G, Blissett J. Tube feeding in infancy: Implications for the development of normal eating and drinking skills. *Dysphagia*. 2005;20(1):46-61.
- 56. Byars KC, Burklow KA, Ferguson K, O'Flaherty T, Santoro K, Kaul A. A multicomponent behavioral program for oral aversion in children dependent on gastrostomy feedings. *Journal of Pediatric Gastroenterology and Nutrition*. 2003;37(4):473-480.
- 57. Kindermann A, Kneepkens CM, Stok A, van Dijk EM, Engels M, Douwes AC.

  Discontinuation of tube feeding in young children by hunger provocation. *Journal of Pediatric Gastroenterology and Nutrition*. 2008;47(1):87-91.

- Garro A, Thurman SK. Kerwin ML E, Ducette JP. Parent/caregiver stress during pediatric hospitalization for chronic feeding problems. *Journal of Pediatric Nursing*. 2005;20(4):268-275.
- 59. Greer AJ, Gulotta CS, Masler EA, Laud RB. Caregiver stress and outcomes of children with pediatric feeding disorders treated in an intensive interdisciplinary program. *Journal of Pediatric Psychology*. 2008;33(6):612-620.
- 60. Al-Yaarubi S, Ramsay M, Rodd C. Megesterol acetate promotes euglycemia and appetite in a child with persistent hyperinsulimenic hypoglycemia. *Acta Paediatrica*. 2004;93(4):422-423.
- 61. Blissett J, Harris G, Kirk J. Effect of growth hormone therapy on feeding problems and food intake in children with growth disorders. *Acta Paediatrica*. 2000;89(6):644-649.
- 62. Homnick DN, Homnick BD, Reeves R, Marks JH, Pimentel RS, Bonnema SK.

  Cyproheptadine is an effective appetite stimulant in cystic fibrosis. *Pediatric Pulmonology*. 2004;38(2):129-134.
- 63. Archer LA. Rosenblaum PL, Streiner DL. The children's eating behaviour inventory: Reliability and validity results. *Journal of Pediatric Psychology*. 1991;16(5):629-670.
- 64. Crist W, Napier-Phillips. Mealtime behaviours of young children: A comparison of normative and clinical data. *Journal of Developmental and Behavioral Pediatrics*. 2001;22(5):279-286.
- 65. Wardle J. Guthrie C, Sanderson S, Rapoport L. Development of the Children's Eating Behaviour Questionnaire. *Journal of Child Psychology and Psychiatry*. 2001;42(7):963-970.
- 66. Ramsay M, Martel C, Porporino M, Zygmuntowicz, C. The Montreal Children's Hospital Feeding Scale: A brief bilingual screening tool for identifying feeding problems. *Paediatrics and Child Health*. 2011;16(3):147-151.
- 67. Dijk Maijn van, Timmerman MF, Martel C, Ramsay M: Towards the development of a Duch screening instrument for the detection of feeding problems in young children. *Netherland Journal of Psychology*. 2011;66(4):112-119.
- 68. Benjasuwatep B, Rattanamongkolgul S, Ramsay M. The Thai version of the Montreal Children's Hospital Feeding Scale (MCH-FS): psychometric properties. *Journal of the Medical Association of Thailand*. 2015;(2)98:163-169.

- 69. Diniz B P, Fagondes S C, Ramsay M: Cross-Cultural adaptation and validation of the Montreal Children's Hospital Scale into Brazilian Portuguese. *Revista Paulista de Pediatria*. 2021 Feb 24:39:e2019377.
- 70. Garcin PRC, Ramsay M, Meyer SCL et al. Validation of the feeding scale "The Montreal Children's Hospital Feeding Scale" (mch-fs) to detect eating disorders in Mexican children. *Acta Pediátrica de México*. 2014;45(5):443-452.
- 71. Sforza E, Onesimo R, Triumbari EK, Leoni C, Giorgio V, Rigante D, Kuczynska EM Ramsay, M, Zampino G. Cross-cultural adaptation and validation of the Italian version of the Montreal Children's Hospital in a special needs' population. *Language and Communication Disorders*. 2023;(58):1223-1231.
- 72. Hauser, MP, Taborda, RA, Oiberman A, Ramsay M: Escala Argentina de Dificultades Alimentarias en Nino (EADAN): Propiedades Psicometrias. *Evaluar*. 2019;19(2):1-14.
- 73. Babik K, Dziechciarz P, Horwath A, Ostazsewski P. The Polish version of the Montreal Children's Hospital Feeding Scale (MCH-FS): Translation, cross-cultural adaptation, and validation. *Pediatria Polka*. 2019:94(5):299-305.
- 74. Lopez AC, Guimaraes C, Afonso C. Montreal Children's Hospital Feeding Scale: Translation and contribution for the validation into European Portugeses. *Revista Portuguesa De Terapia Da Fala*. 2015;(3):05-15.
- 75. Casaubon Garcin P, Ramsay M, Casaubon Lemmen Meyer S, Goldbard Rochman D, Oliva Meza Hernandez O, Baptista González H, Monroy Tenia Z, Barriguete Melendez J, Valdez M. Validación de la escala de alimentación "The Montreal Children's Hospital Feeding Scale" (mch-fs) para detectar trastornos de la alimentación en niños mexicanos. *Acta Pediátrica De México*. 2024;45(5):443-452.
- 76. Dogru I, Manav G, Uymaz P. Turkish validity and reliability study of the Montreal Children's Hospital Feeding Scale. *Current Pediatric Research*. 2022;26(3):1289-1296.
- 77. Zhongxia Ren, Hanglian Lan, Ignatius Man-Yau Szeto, AI Zhao. Feeding difficulty among chinese toddlers aged 1-3 years and its association with health and development. *Frontiers in Pediatrics*. 2021 Nov 23;9:758176.

# Feeding Behaviour of Infants and Young Children and Its Impact on Child Psychosocial and Emotional Development

Yi Hui Liu, MD MPH, Martin T. Stein, MD University of California San Diego, USA September 2013, 2e éd.

#### Introduction

Feeding is a primary event in the life of an infant and young child. It is the focus of attention for parents and other caregivers, and a source of social interaction through verbal and non-verbal communication. The eating experience provides not only sustenance but also an opportunity for learning. It affects not only children's physical growth and health but also their psychosocial and emotional development. The feeding relationship is affected by culture, health status and temperament.

#### Subject

The essential component of feeding behaviour in young children is the relationship between the child and the primary caregiver. The first three years of life are a particular challenge because a child's feeding abilities and needs change with motor, cognitive and social development. In the first stage (birth to three months) of self-regulation and organization, the child integrates experiences of hunger and satiety to develop regular feeding patterns. In the second stage (three to seven months), the infant and parent form an attachment that allows them to communicate with each other and the infant develops basic trust and self-soothing behaviours. In the third stage (six to 36 months), the child gradually "separates" emotionally from the parent and discovers a sense of independence or autonomy, making use of developing motor and language skills to control the environment and establish independent feeding.

With participation in family meals, the social component of feeding expands. The child begins to mimic eating choices, patterns and behaviours modelled by family members. The structure of family meals sets limits for the child as he or she achieves independent feeding skills. The accessibility of particular foods, modelling, media exposure and feeding interactions shape a

child's eating behaviour and food preferences.

The caregiver's behaviours and the child's temperament influence the feeding relationship. The parent who allows her infant to determine timing, amount and pacing of a meal helps her infant develop self-regulation and secure attachment. The parent who allows her toddler to explore the environment while providing structure and appropriate limits helps her child develop motor and social skills. The effective parent adjusts and responds appropriately to her child's temperament — the child's emotional reactivity, adaptability and response to change. Temperament can affect how a child approaches and responds to new foods and to a parent's feeding patterns.

Culture may significantly influence the feeding experience. It may determine not only the choice of infant feeding (breast milk or formula) but also associated behaviours (co-sleeping is linked to prolonged breastfeeding), the length of feeding method (later weaning in developing countries versus earlier weaning for working mothers in developed countries), and exposure to feeding environments outside the home (child care among families with mother who work outside the home).

#### **Problems**

Mild and transient feeding problems occur in 25% to 35% of young children while severe and chronic feeding problems occur in 1% to 2%.¹ Common conditions include overeating, poor eating, feeding behaviour problems and unusual or unhealthy food choices. Although medical disorders and inappropriate food selection can result in feeding problems, these conditions are often associated with early problems in parent-child feeding experiences. Problems with self-regulation, attachment, temperament and the development of autonomy can contribute. A poor attachment may result from substance abuse or mental illness in the caregiver, developmental delay or a medical condition in the child, and parent-child personality/temperament conflict.

While most feeding problems in infants and young child are temporary, emotional and social development may be impacted during late childhood, adolescence and adulthood. Obesity, cardiovascular disease, diabetes mellitus and behavioural problems are more frequent in those with early childhood feeding problems.

1. *Overeating*. The prevalence of overweight and obesity in the United States has increased to 10.4% in two- to five-year olds, 15.3% in six- to 11-year olds, and 15.5% in 12- to 19-year olds.<sup>2</sup> These children are not only at risk for medical problems (e.g. diabetes mellitus,

hypertension, orthopedic problems, obstructive sleep apnea), but also poor self-esteem, disturbed body image, social isolation, maladjustment, depression and eating disorders. Social stigmatization begins as early as preschool and continues into school-age as their peers may reject overweight children. Parental concerns about overeating and obesity may result in inappropriate restriction of their young child's diet.

- 2. Poor eating or not gaining sufficient weight. A parent may misperceive her child as having insufficient nutritional intake when the child is active and more interested in play and the environment than in meals. Some parents have inappropriate expectations about sufficient food portions and weight gain. Failure to thrive (FTT) occurs when a child's rate of weight gain has decreased to below the third to fifth percentile for gestation-corrected age and sex, or the child's weight has fallen and crossed two major percentiles in a standardized growth chart. Children with FTT may have impaired growth (e.g. height, head circumference) and developmental skills and are at risk for long-term developmental and behavioural problems.
- 3. Feeding behaviour problems. Parents may have difficulty making the transition from an infant who is cooperative during feeding to a toddler who seeks independence at mealtime. Limited food preferences may be normal and temporary during this period or may develop into a behavioural disorder. Food phobias or a post-traumatic feeding disorder may result from a painful episode (e.g. choking with a particular food) or a difficult experience associated with a food-induced allergic reaction.
- 4. *Unusual choices*. Pica, or the ingestion of non-food substances, is normal in children under two years of age who explore their environment through hand-to-mouth experiences. After two years of age, pica is a behavioural condition more frequent in children with insufficient stimulation, psychological disorders and mental retardation.
- 5. *Unhealthy food choices*. Food preferences are established through exposure and accessibility to foods, modelling and advertisements. Most "alternative" diets are not harmful, although specific nutrient deficiencies should be addressed with some (for example, iron and vitamin B12 in vegan diets).

#### **Research Context**

Early childhood feeding experiences affect both health and psychological well-being. Because many feeding problems have their roots in infancy and childhood, current research focuses on

determining the antecedents to these problems and the effectiveness of modifying various factors.

#### **Key Research Questions**

What are the most significant behavioural antecedents to childhood obesity that affect feeding? How can they be modified? How can behavioural changes be sustained? What are the most effective community-based interventions that have an impact on optimal nutritional choices and early feeding behaviours? What cultural determinants influence optimal feeding behaviours in early childhood? How can a better understanding of unique cultural values and habits influence medical and public-health programs to improve childhood nutrition?

#### **Recent Research Results**

Behavioural research in childhood feeding has focused on breastfeeding (choice, initiation and sustainability), teaching parents developmentally-appropriate feeding methods, and behavioural programs directed to specific feeding disorders, including obesity, failure to thrive and anorexia nervosa. In each case, principles of behaviour modification, health promotion and education have been applied effectively.

Many studies have examined the proposal that breastfeeding protects against the development of obesity later in life. While some have found an insignificant effect, 3.4.5.6.7 others have found a significant and even a dose-response relationship 9.10,11 between breastfeeding duration and lower risk of child obesity. Without a consensus, the benefits of breastfeeding (e.g. establishment of attachment, optimal nutrition and protection against certain infectious diseases), still support encouraging breastfeeding whenever possible. With breastfeeding, lower maternal control of food intake and greater maternal responsiveness to infant cues has a beneficial effect on infant-feeding style and food intake, acknowledges the infant's ability to self-regulate appropriate food intake, and may contribute to healthier eating patterns. 12

Child-feeding practices and behavioural interventions may modify patterns of intake. An overview of pediatric obesity treatment concluded that dietary changes accompanied by behaviour change methods, exercise and parental involvement are important in long-term success. Parental participation and modelling is instrumental in establishing and changing eating patterns in children. Modelling consumption of healthy foods, such as fruit and vegetables, has a positive effect on the consumption of those foods by children whereas

modelling dieting behaviours results in problems in regulating a child's intake.<sup>15</sup> Television has a powerful influence on the foods children request; limiting television viewing can lessen obesity.<sup>16</sup> Birch and Fisher have written an excellent review detailing the determinants of children's dietary intake and responses to their modification.<sup>17</sup>

#### **Conclusions**

Feeding infants and young children is a behavioural event influencing their growth and development. Early experiences with feeding set the stage for healthy feeding-associated behaviours in later childhood and adulthood. Understanding the development of normal feeding behaviour in infants and young children makes it easier to distinguish between self-limited concerns and those requiring further intervention. Parents and other caregivers need knowledge about both nutritional content and developmentally appropriate feeding behaviours. Since earlier onset of problems results in more significant consequences, prevention of feeding disorders and related behaviour problems should be targeted towards guiding the feeding behaviours of infants and young children and their feeding relationships with parents and caregivers. Obesity (especially in developed countries) and undernutrition (especially in developing countries) can be addressed only through a combination of making healthy food available, ensuring an understanding of age-appropriate feeding practices, and supporting the emotional health of families. Cultural differences and temperament variations should be incorporated into any recommendations.

#### **Implications for Policy and Service**

- 1. Establish national dietary guidelines that are specific for children and easily understood and applied by parents.
- 2. Promote and support breastfeeding. The goals of *Healthy People 2010* are to increase the proportion of mothers who breastfeed to 75% in the early postpartum period, 50% at six months and 25% at one year. Educate pregnant and new mothers on the advantages and maintenance of breastfeeding.
- 3. Advocate for nutrition in schools. Endorse and fund healthy school lunches and free school breakfasts (e.g. the United States federal government's *School Breakfast Program*<sup>19</sup>). Remove soda, sweetened beverages and unhealthy snacks from school campuses.<sup>20</sup> Support nutrition education in classrooms.

- 4. Require regular physical education in schools to promote a healthy lifestyle and to help decrease obesity.<sup>21</sup>
- 5. Restrict television advertisements endorsing unhealthy food choices. Use media to promote healthy eating and regular physical activity.
- 6. Increase the availability of affordable fresh foods, especially fruit and vegetables, in low socioeconomic communities.
- 7. Promote education about healthy eating habits through public-health messages and increase funding for public-health campaigns promoting breastfeeding, healthy foods and obesity prevention.
- 8. Fund research investigating the etiology, prevention and treatment of obesity; factors influencing choice of breastfeeding, food intake and physical activity; and child-feeding practices in differing socioeconomic and ethnic groups.
- 9. Form public and private partnerships to promote healthy eating. Coordinate efforts of policy-makers, health professionals, community leaders and parents.

#### References

- 1. Satter E. The feeding relationship: problems and interventions. Journal of Pediatrics 1990;117(2 Pt 2):S181-S189.
- 2. Ogden CL, Flegal KM, Carroll MD, Johnson CL. Prevalence and trends in overweight among US children and adolescents, 1999-2000. *JAMA Journal of the American Medical Association* 2002;288(14):1728-1732.
- 3. Baranowski T, Bryan GT, Rassin DK, Harrison JA, Henske JC. Ethnicity, infant-feeding practices, and childhood adiposity. *Journal of Developmental and Behavioral Pediatrics* 1990;11(5):234-239.
- 4. Elliott KG, Kjolhede CL, Gournis E, Rasmussen KM. Duration of breastfeeding associated with obesity during adolescence. *Obesity Research* 1997;5(6):538-541.
- 5. Hediger ML, Overpeck MD, Kuczmarski RJ, Ruan WJ. Association between infant breastfeeding and overweight in young children. *JAMA Journal of the American Medical Association* 2001;285(19):2453-2460.
- 6. Wolman PG. Feeding practices in infancy and the prevalence of obesity in preschool children. *Journal of the American Dietetic Association* 1984;84(4):436-438.
- 7. Zive MM, McKay H, Frank-Spohrer GC, Broyles SL, Nelson JA, Nader PR. Infant-feeding practices and adiposity in 4-y-old Anglo- and Mexican-Americans. *American Journal of Clinical Nutrition* 1992;55(6):1104-1108.
- 8. Kramer MS. Do breast-feeding and delayed introduction of solid foods protect against subsequent obesity? *Journal of Pediatrics* 1981;98(6):883-887.
- 9. Gillman MW, Rifas-Shiman SL, Camargo CA Jr., Berkey CS, Frazier AL, Rockett HR, Field AE, Colditz GA. Risk of overweight among adolescents who were breastfed as infants. *JAMA- Journal of the American Medical Association* 2001;285(19):2461-2467.
- 10. Liese AD, Hirsch T, von Mutius E, Keil U, Leupold W, Weiland SK. Inverse association of overweight and breast feeding in 9 to 10-y-old children in Germany. *International Journal of Obesity and Related Metabolic Disorders* 2001;25(11):1644-

1650.

- 11. von Kries R, Koletzko B, Sauerwald T, von Mutius E, Barnert D, Grunert V, von Voss H. Breast feeding and obesity: cross sectional study. *BMJ British Medical Journal* 1999;319(7203):147-150.
- 12. Fisher JO, Birch LL, Smiciklas-Wright H, Picciano MF. Breast-feeding through the first year predicts maternal control in feeding and subsequent toddler energy intakes. *Journal of the American Dietetic Association* 2000;100(6):641-646.
- 13. Epstein LH, Myers MD, Raynor HA, Saelens BE. Treatment of pediatric obesity. Pediatrics 1998;101(3 Pt 2):554-570.
- 14. Fisher JO, Mitchell DC, Smiciklas-Wright H, Birch LL. Parental influences on young girls' fruit and vegetable, micronutrient, and fat intakes. *Journal of the American Dietetic Association* 2002;102(1):58-64.
- 15. Johnson SL. Improving preschoolers' self-regulation of energy intake. Pediatrics 2000;106(6):1429-1435.
- 16. Robinson TN. Reducing children's television viewing to prevent obesity: a randomized controlled trial. *JAMA- Journal of the American Medical Association* 1999;282(16):1561-1567.
- 17. Birch LL, Fisher JO. Development of eating behaviors among children and adolescents. *Pediatrics* 1998;101(3 Pt 2):539-549
- 18. US Department of Health and Human Services. Maternal, infant and child health. In: *Healthy people 2010: Conference edition*. Vol 2. Washington, DC: US Government Printing Office; 2000:47-48. Available at: http://www.healthypeople.gov/Document/HTML/Volume2/16MICH.htm. Accessed September 20, 2005.
- 19. McBean LD, Miller GD. Enhancing the nutrition of America's youth. *Journal of the American College of Nutrition* 1999;18(6):563-571.
- 20. American Academy of Pediatrics, Committee on School Health. Soft drinks in schools. *Pediatrics* 2004;113(1 Pt 1):152-154.
- 21. American Academy of Pediatrics. Physical fitness and activity in schools. Pediatrics 2000;105(5):1156-1157.

## **Helping Children Develop Healthy Eating Habits**

Maureen M. Black, PhD, Kristen M. Hurley, PhD

University of Maryland School of Medicine, USA September 2013, 2e éd. rév.

#### Introduction

The first year of life is characterized by rapid developmental changes related to eating. As infants gain truncal control, they progress from sucking liquids in a supine or semi-reclined position to eating solid foods in a seated position. Oral motor skills progress from a basic suck-swallow mechanism with breast milk or formula to a chew-swallow mechanism with semi-solids, progressing to complex textures.<sup>1,2</sup> As infants gain fine motor control, they progress from being fed exclusively by others to at least partial self-feeding. Their diet extends from breast milk or formula, through purees and specially prepared foods, to the family diet. By the end of the first year of life, children can sit independently, can chew and swallow a range of textures, are learning to feed themselves and are making the transition to the family diet and meal patterns.

As children transition to the family diet, recommendations address not only food, but also the eating context. A variety of healthy foods promote diet quality, along with early and sustained food acceptance. Data gathered on infants and young children 6 to 23 months of age across 11 countries have demonstrated a positive association between dietary variety and nutritional status.<sup>3</sup> Exposure to fruits and vegetables in infancy and toddlerhood have been associated with acceptance of these foods at later ages.<sup>4-6</sup>

Children's eating patterns and food preferences are established early in life. When children refuse nutritious foods such as fruits or vegetables, mealtimes can become stressful or confrontational, and children may be denied both the nutrients they require and healthy, responsive interactions with caregivers. Caregivers who are inexperienced or stressed, and those who have poor eating habits themselves, may be most in need of assistance to facilitate healthy, nutritious mealtime behaviour with their children.

#### **Subject**

Problems associated with eating occur in 25% to 45% of all children, particularly when children are acquiring new skills and are challenged with new foods or mealtime expectations. For example, infancy and toddlerhood are characterized by bids for autonomy and independence as children strive to do things themselves. When these characteristics are applied to eating behaviours, children may be neophobic (hesitant to try new foods) and insist on a limited repertoire of foods, leading them to be described as picky eaters.

Most eating problems are temporary and easily resolved with little or no intervention. However, eating problems that persist can undermine children's growth, development, and relationships with their caregivers, leading to long-term health and developmental problems. Children with persistent eating problems whose caregivers do not seek professional advice until the problems become severe, may be at risk for growth or behaviour problems.

#### **Problems**

Eating patterns have developmental, family and environmental influences. As children become developmentally able to make the transition to family foods, their internal regulatory cues for hunger and satiety may be overridden by familial and cultural patterns. At the family level, children of caregivers who model healthy food intakes are likely to consume more fruits and vegetables than children of caregiver who do not, whereas children of caregivers who model less healthy, snack food intakes are likely to establish patterns of eating behaviours and food preferences that include excess amounts of fat and sugar. At the environmental level, children's frequent exposure to fast-food and other restaurants has led to increased consumption of high-fat foods, such as french fries, rather than more nutritious options, such as fruit and vegetables. In addition, caregivers may not realize that many commercial products marketed for children, such as sweetened drinks, may satisfy hunger or thirst, but provide minimal nutritional benefits.

National surveys have reported excessive caloric intakes during toddlerhood,<sup>13,14</sup> and many children continue to consume alarmingly low quantities of fruit and vegetables and essential micronutrients.<sup>15</sup> By elementary school, many children receive over half their beverage intake from sweetened drinks,<sup>16</sup> a pattern that undoubtedly begins during the toddler and preschool years. These poor nutritional patterns (high fat, sugar and refined carbohydrates; sweetened drinks; and limited fruit and vegetables) increase the likelihood of micronutrient deficiencies (e.g., Iron Deficiency Anemia) and excess weight gain in young children.<sup>17</sup>

#### **Research Context**

Eating is often studied through observational studies or caregiver reports of mealtime behaviour. Some investigators rely on clinical samples of children with growth or eating problems, while others recruit normative children.

#### **Key Research Questions**

Key questions include the progression of eating behaviours from infancy through toddlerhood, methods children use to signal hunger and satiety, and why some children (the so-called "picky" eaters) have selective food preferences. Key questions for caregivers and families are how to promote healthy eating behaviours in young children, how to encourage children to eat healthy food, and how to avoid problems in feeding and growth.

#### **Recent Research Results**

#### Attachment and eating

Healthy eating behaviour begins in infancy, as infants and their caregivers establish a partnership in which they recognize and interpret both verbal and non-verbal communication signals from one another. This reciprocal process forms a basis for the emotional bonding or attachment between infants and caregivers that is essential to healthy social functioning. <sup>18</sup> If there is a disruption in the communication between children and caregivers, characterized by inconsistent, non-responsive interactions, the attachment bond may not be secure, and eating may become an occasion for unproductive, upsetting battles over food.

Infants who do not provide clear signals to their caregivers or do not respond to their caregivers' efforts to help them establish predictable routines of eating, sleeping and playing are at risk for regulatory problems that may include eating. Infants who are premature or ill may be less responsive than healthy full-term infants and less able to communicate hunger or satiety. Caregivers who do not recognize their infants' satiety cues may overfeed them, causing infants to associate feelings of satiety with frustration and conflict.

#### The caregiver-child context of feeding

Variability in the caregiver-child feeding context is related to children's eating behaviour and growth.<sup>19</sup> The dimensions of parental structure and nurturance, which incorporate parents'

perceptions of their child's behaviour, have been applied to the feeding context (Figure 1). <sup>20,21,22</sup> Responsive feeding reflects a reciprocal pattern in which caregivers provide guidance and developmentally appropriate responses to their child's signals of hunger and satiety. Unresponsive feeding is marked by a lack of reciprocity between the caregiver and child, often characterized by the caregiver taking excessive control of the feeding context (forcing/pressuring or restricting food intake), the child controlling the feeding context (e.g., demanding a limited repertoire of food, indulgent feeding), or the caregiver ignoring the child's signals or failing to establish mealtime routines (uninvolved feeding). <sup>23,24</sup>

Figure 1. The Caregiver-Child Feeding Context: Patterns of Parenting and Feeding

	NURTURANCE	
	HIGH	LOW
	AUTHORITATIVE	AUTHORITARIAN
HIGH	• Involved	• Forceful
	• Nurturing	• Restrictive
	• Structured	• Structured
	Responsive	Controlling
	INDULGENT	UNINVOLVED
LOW	• Involved	<ul> <li>Unengaged</li> </ul>
	• Nurturing	<ul><li>Insensitive</li></ul>
	• Unstructured	• Unstructured
	Indulgent	Uninvolved

A responsive feeding style, high in nurturance and structure, a derivative of authoritative parenting, represents caregivers who form a relationship with their child that involves clear demands and mutual interpretation of signals and bids for mealtime interaction. Responsive feeding is characterized by interactions that are prompt, contingent on the child's behaviour and developmentally appropriate with an easy give-and-take.<sup>22,25,26</sup>

A controlling feeding style, high in structure and low in nurturance, represents caregivers who use forceful or restrictive strategies to control mealtimes. Controlling feeding is embedded in an overall authoritarian pattern of parenting and may include over-stimulating behaviours, such as speaking loudly, forcing foods or otherwise overpowering the child.<sup>27</sup> Controlling caregivers may override their child's internal regulatory cues for hunger and satiety.<sup>28</sup> The innate capacity that infants have to self-regulate their energy intake declines during early childhood in response to family and cultural patterns.<sup>29</sup>

An indulgent feeding style, high in nurturance and low in structure, is embedded in an overall indulgent style of parenting, and occurs when caregivers allow children to make decisions around meals, such as when and what they will eat.<sup>23</sup> Without parental guidelines, children are likely to be attracted to high salt/high sugar foods, rather than to a more balanced variety including vegetables.<sup>23</sup> Thus, an indulgent feeding style may be problematic, given infants' genetic predispositions to prefer sweet and salty tastes.<sup>30</sup> Children of caregivers who display an indulgent feeding style are often heavier than children of caregivers who use non-indulgent feeding styles.<sup>24</sup>

An uninvolved feeding style, low in both nurturance and structure, often represents caregivers who have limited knowledge and involvement in their child's mealtime behaviour.<sup>23</sup> Uninvolved child feeding styles may be characterized by little or no active physical help or verbalization during feeding, lack of reciprocity between the caregiver and child, a negative feeding environment and a lack of feeding structure or routine. Uninvolved feeders often ignore both child feeding recommendations and their toddler's cues of hunger and satiety and may be unaware of what or when their toddler is eating. Egeland and Sroufe<sup>31</sup> found that children of uninvolved or psychologically unavailable caregivers were more likely to be anxiously attached when compared with children of available caregivers. An uninvolved feeding style is embedded in an overall uninvolved style of parenting.<sup>23</sup>

Several recent systematic reviews report associations between parental feeding control and infant and early child weight gain and/or weight status.<sup>24,32,33</sup> Controlling feeding has been associated with increased weight gain (e.g., children of caregivers who use restrictive feeding practices tend to overeat)<sup>34</sup> and to decreased weight gain (e.g., children who are pressured to eat more, do not).<sup>35</sup> However, the cross-sectional design of most studies, along with a tendency to rely exclusively on caregiver behaviour, rather than consider the reciprocal nature of feeding interactions, has hindered the understanding of caregiver-child feeding interactions. A recent

randomized controlled trial among infants in Australia found that providing anticipatory guidance regarding infant feeding behaviour led to healthier weight gain and higher rates of self-reported responsive feeding behaviour.<sup>36</sup> Additional trials are needed to better understand strategies to promote healthy feeding interactions and healthy growth.

#### Food preferences

Children who are raised with caregivers who model healthy eating behaviours, such as a diet rich in fruit and vegetables, establish food preferences that include fruit and vegetables.<sup>4</sup>

Food preferences are also influenced by associated conditions. Children are likely to avoid food that has been associated with unpleasant physical symptoms, such as nausea or pain. They may also avoid food that has been associated with the anxiety or distress that often occurs during meals characterized by arguments and confrontations.

Children also accept or reject food based on qualities of the food, such as taste, texture, smell, temperature or appearance, as well as environmental factors, such as the setting, the presence of others and the anticipated consequences of eating or not eating. For example, consequences of eating may include relief from hunger, participation in a social function or attention from caregivers. Consequences of not eating may include additional time to play, becoming the focus of attention or getting snack food instead of the regular meal.

Increasing familiarity with the taste of a food increases the likelihood of acceptance.<sup>37,38</sup> Caregivers can facilitate the introduction of new foods by pairing the new food with preferred food and presenting the new food repeatedly until it is no longer "new."

#### Conclusions

Eating patterns are established early in life in response to internal regulatory cues, caregiver-child interactions, mealtimes routines, foods offered and modeling from family members. Exposing children to fruits and vegetables early in life establishes a pattern of fruit and vegetable preference and consumption throughout life. Research is needed to investigate the individual, interactive and environmental determinants of the caregiver-child feeding context, relationships between responsive/unresponsive feeding and children's eating behaviour and weight gain and population-specific validated tools to measure responsive/unresponsive feeding.

Early childhood eating behaviours are heavily influenced by caregivers and are learned through early experiences with food and eating. Education and support provided by health professionals (i.e., public health nurses, family physicians and pediatricians) and nutrition programs need to be strengthened to ensure that caregivers have the facilities needed to address issues of eating behaviours during childhood.

Caregivers should eat with children so modelling can occur and mealtimes are viewed as pleasant social occasions. Eating together lets children watch caregivers try new foods and helps children and caregivers communicate hunger and satiety, as well as enjoyment of specific foods.

Caregivers control both the food that is offered and the mealtime atmosphere. Their "job" is to ensure that children are offered healthy food on a predictable schedule in a pleasant setting. <sup>39</sup> By developing mealtime routines, caregivers help children learn to anticipate when they will eat. Children learn that feelings of hunger are soon relieved and there is no need to feel anxious or irritable. Children should not graze or eat throughout the day, so they develop an expectation and an appetite around mealtime. <sup>39</sup>

Mealtimes should be pleasant and family-oriented, with family members eating together and sharing the events of the day. When mealtimes are too brief (less than 10 minutes), children may not have enough time to eat, particularly when they are acquiring self-feeding skills and may eat slowly. Alternatively, sitting for more than 20 or 30 minutes is often difficult for a child and mealtimes may become aversive.

When meals are characterized by distractions from television, family arguments or competing activities, children may have difficulty focusing on eating. Caregivers should separate mealtime from playtime and avoid using toys, games, or television to distract the child during mealtime. Child-oriented equipment, such as highchairs, bibs and small utensils, may facilitate eating and enable children to acquire the skills of self-feeding.

#### **Implications**

Implications can be directed to environmental, family and individual levels. At the environmental level, encouraging fast-food and other restaurants to also provide healthy, palatable food options that are appealing to young children may reduce some of the feeding problems that occur when children are repeatedly exposed to high-fat foods, such as french fries, rather than to nutritious

options, such as fruit and vegetables. At the family level, guidelines for children's nutrition should include information on their nutritional needs and on strategies to promote healthy eating behaviour, including recognizing children's signals of hunger and satiety and use of appropriate feeding interactions, allocating time for meals, scheduling meals at relatively consistent times, promoting new foods through modelling and avoiding stress and conflict during meals. At the individual level, programs that help children develop healthy eating patterns by eating nutritious foods and eating to satisfy hunger, rather than to satisfy emotional needs, may prevent subsequent health and developmental problems.<sup>40</sup>

#### References

- 1. Bosma J. Development and impairments of feeding in infancy and childhood. In: Groher ME, ed. *Dysphagia: Diagnosis and management*. 3rd ed. Boston, MA: Butterworth-Heinemann; 1997:131-138.
- 2. Morris SE. Development of oral motor skills in the neurologically impaired child receiving non-oral feedings *Dysphagia* 1989;3:135-154.
- 3. Arimond M, Ruel MT. Dietary diversity is associated with child nutritional status: Evidence from 11 demographic and health surveys. *The Journal of Nutrition* 2004;134:2579-2585.
- 4. Skinner JD, Carruth BR, Bounds W, Ziegler P, Reidy K. Do food-related experiences in the first 2 years of life predict dietary variety in school-aged children? *Journal of Nutrition Education and Behavior* 2002;34(6):310-315.
- 5. Schwartz C, Scholtens PA, Lalanne A, Weenen H, Nicklaus S. Development of healthy eating habits early in life. Review of recent evidence and selected guidelines. *Appetite*. 2011;57(3):796-807.
- 6. Mennella JA, Nicklaus S, Jagolino AL, Yourshaw LM. Variety is the spice of life: strategies for promoting fruit and vegetable acceptance during infancy. *Physiol Behav.* 2008;22;94(1):29-38.
- 7. Linscheid TR, Budd KS, Rasnake LK. Pediatric feeding disorders. In: Roberts MC, ed. *Handbook of pediatric psychology*. New York, NY: Guilford Press; 2003:481-498.
- 8. Birch LL, McPhee L, Shoba BC, Pirok E, Steinberg L. What kind of exposure reduces children's food neophobia? Looking vs tasting. *Appetite* 1987;9(3):171-178.
- 9. Keren M, Feldman R, Tyano S. Diagnoses and interactive patterns of infants referred to a community-based infant mental health clinic. *Journal of the American Academy of Child & Adolescent Psychiatry* 2001;40(1):27-35.
- 10. Palfreyman Z, Haycraft E, Meyer C. Development of the Parental Modeling of Eating Behaviours Scale (PARM): links with food intake among children and their mothers. *Maternal and Child Nutrition*. 2012 [Epub ahead of print].
- 11. Zoumas-Morse C, Rock CL, Sobo EJ, Neuhouser ML. Children's patterns of macronutrient intake and associations with restaurant and home eating. *Journal of the American Dietetic Association* 2001;101(8):923-925.
- 12. Smith MM, Lifshitz F. Excess fruit juice consumption as a contributing factor in nonorganic failure to thrive. *Pediatrics* 1994;93(3):438-443.
- 13. Ponza M, Devaney B, Ziegler P, Reidy K, Squatrito C. Nutrient intakes and food choices of infants and toddlers participating in WIC. *Journal of the American Dietetic Association* 2004;104(1 Suppl 1):71-79.
- 14. Devaney B, Kalb L, Briefel R, Zavitsky-Novak T, Clusen N, Ziegler P. Feeding infants and toddlers study: overview of the study design. *Journal of the American Dietetic Association* 2004;104(1 Suppl 1):8-13.

- 15. Picciano MF, Smiciklas-Wright H, Birch LL, Mitchell DC, Murray-Kolb L, McConahy KL. Nutritional guidance is needed during dietary transition in early childhood. *Pediatrics* 2000;106(1):109-114.
- 16. Cullen KW, Ash DM, Warneke C, de Moor C. Intake of soft drinks, fruit-flavored beverages, and fruits and vegetables by children in grades 4 through 6. *American Journal of Public Health* 2002;92(9):1475-1477.
- 17. Brotanek JM, Gosz J, Weitzman M, Flores G. Secular trends in the prevalence of iron deficiency among US toddlers, 1976-2002. *Archives of Pediatrics & Adolescent Medicine* 2008;162:374-81.
- 18. Ainsworth MDS, Blehar MC, Waters E, Wall S. *Patterns of attachment: A psychological study of the strange situation*. New York: Psychology Press, 1978.
- 19. Rhee K. Childhood overweight and the relationship between parent behaviors, parenting style, and family functioning. *The Annals of the American Academy of Political and Social Science* 2008;615:11–37.
- 20. Baumrind D. Rearing competent children In: Damon W, ed. *Child development today and tomorrow*. San-Francisco, CA: Jossey-Bass Publishers; 1989:349-378.
- 21. Maccoby EE, Martin J. Socialization in the context of the family: parent-child interaction. In: Hetherington EM, ed. *Handbook of child psychology: Socialization, personality, and social development. Vol 4.* New York, NY: John Wiley; 1983:1-101.
- 22. Black MM & Aboud FE. Responsive feeding is embedded in a theoretical framework of responsive parenting. *Journal of Nutrition* 2011;141(3):490-4.
- 23. Hughes SO, Power TG, Fisher JO, Mueller S, Nicklas TA. Revisiting a neglected construct: Parenting styles in a child-feeding context. *Appetite* 2005;44(1):83-92.
- 24. Hurley KM, Cross MB, Hughes SO. A systematic review of responsive feeding and child obesity in high-income countries. *Journal of Nutrition* 2011;141:495-501.
- 25. Leyendecker B, Lamb ME, Scholmerich A, Fricke DM. Context as moderators of observed interactions: A study of Costa Rican mothers and infants from differing socioeconomic backgrounds. *International Journal of Behavioural Development* 1997;21(1):15-24.
- 26. Kivijarvi M, Voeten MJM, Niemela P, Raiha H, Lertola K, Piha J. Maternal sensitivity behaviour and infant behaviour in early interaction. *Infant Mental Health Journal* 2001;22(6):627-640.
- 27. Beebe B, Lachman F. *Infant research and adult treatment: Co-constructing interactions.* Hillsdale, NJ: The Analytic Press; 2002.
- 28. Birch LL, Fisher JO. Mothers' child-feeding practices influence daughters' eating and weight. *American Journal of Clinical Nutrition* 2000;71(5):1054-1061
- 29. Birch LL, Johnson SL, Andresen G, Peters JC, Schulte MC. The variability of young children's energy intake. *New England Journal of Medicine* 1991;324(4):232-235.
- 30. Birch LL. Development of food preferences. Annual Review of Nutrition 1999;19:41-62.
- 31. Egeland B, Sroufe LA. Attachment and early maltreatment. Child Development 1981;52(1):44-52.
- 32. DiSantis KI, Hodges EA, Johnson SL, Fisher JO. The role of responsive feeding in overweight during infancy and toddlerhood: a systematic review. *International Journal of Obesity* 2011;35:480-92.
- 33. Faith MS, Scanlon KS, Birch LL, Francis LA, Sherry B. Parent-child feeding strategies and their relationships to child eating and weight status. *Obesity Research* 2004;12(11):1711-1722.
- 34. Birch LL, Fisher JO, Davison KK. Learning to overeat: maternal use of restrictive feeding practices promotes girls' eating in the absence of hunger. *American Journal of Clinical Nutrition* 2003;78(2):215-220.

- 35. Fisher JO, Mitchell DC, Smiciklas-Wright H, Birch LL. Parental influences on young girls' fruit and vegetable, micronutrient, and fat intakes. *Journal of the American Dietetic Association* 2002;102(1):58-64.
- 36. Daniels LA, Mallan KM, Battistutta D, Nicholson JM, Perry R, Magarey A. Evaluation of an intervention to promote protective infant feeding practices to prevent childhood obesity: outcomes of the NOURISH RCT at 14 months of age and 6 months post the first of two intervention modules. *International Journal of Obesity (Lond)*. 2012 Oct;36(10):1292-8.
- 37. Birch LL. Children's preferences for high-fat foods. Nutrition Reviews 1992;50(9):249-255.
- 38. Birch LL, Marlin DW. I don't like it; I never tried it: effects of exposure on two-year old children's food preferences. *Appetite* 1982;3(4):353-360.
- 39. Satter E. Child of mine: Feeding with love and good sense. Palo Alto, CA: Bull Publishing; 2000.
- 40. Black MM, Cureton LA, Berenson-Howard J. Behaviour problems in feeding: Individual, family, and cultural influences. In: Kessler DB, Dawson P, eds. *Failure to thrive and pediatric undernutrition: A transdisciplinary approach*. Baltimore, Md: Paul H. Brookes Publishing Co.; 1999:151-169.

## Assessment and Treatment of Pediatric Feeding Disorders

<sup>1</sup>Suzanne M. Milnes, PhD, <sup>1</sup>Cathleen C. Piazza, PhD, <sup>2</sup>Tammy Carroll, PhD

<sup>1</sup>Munroe-Meyer Institute, University of Nebraska Medical Center, USA,

<sup>2</sup>University of Alabama, USA

September 2013, Éd. rév.

#### Introduction

A feeding disorder is identified when a child is unable or refuses to consume a sufficient quantity or variety of solids and liquids to maintain proper nutrition.<sup>1</sup> The complications from feeding problems range from mild (e.g., missed meals) to severe (e.g., severe malnourishment).<sup>2</sup> Mealtime difficulties occur in approximately 25% to 35% of typically developing children and up to 80% of individuals with developmental disabilities.<sup>3-6</sup> Feeding disorders may be manifested by total refusal to eat, dependence on supplemental feedings (e.g., gastrostomy tube), inappropriate mealtime behaviour, and selectivity by type and texture.

#### **Subject**

The causes of feeding disorders are equally varied. Feeding problems are often caused by a number of biological and environmental factors, which interact. For example, Rommel et al. evaluated 700 children referred to an interdisciplinary feeding team and found combined causes (e.g., medical, behavioural, oral-motor) of the feeding problem occurred in over 60% of patients.

Biological factors may include early experiences with medical procedures, chronic hospitalization, or medical problems, which cause eating to be painful. Even after the painful medical condition is treated, the child may continue to refuse food because if the child never or rarely eats, he or she never learns that eating is no longer painful. If the child refuses to eat, he or she does not have the opportunity to practice eating and does not develop the skills to become a capable eater. Refusal to eat may lead to growth failure, which contributes to poor feeding skills as undernourished children lack the energy to become capable eaters. Thus, a cycle develops in which the child refuses food, fails to learn that eating is no longer painful, misses opportunities to develop oral motor skills, and fails to gain weight.

#### **Problems**

Even when the cause of food refusal is a painful medical condition, caregiver responses to the child during meals may worsen the problem. Piazza and colleagues observed caregivers and children with feeding problems during meals, which showed that caregivers used a variety of strategies to encourage eating such as distracting, coaxing, and reprimanding; allowing the child to periodically take a break from or avoid eating; and providing preferred food or toys. All children displayed refusal behaviour and infrequently ate bites of food. When Piazza et al. analyzed the effects of caregiver behaviour during meals on child feeding behaviours, results indicated that the above listed strategies used by the caregivers to encourage eating actually worsened behaviour for 67% of the children.

These results are not surprising given the relation between the causes of feeding problems and caregiver behaviour. Parental strategies such as stopping the meal or coaxing may produce the immediate effect of temporarily stopping undesirable child behaviour. From the child's perspective, the study by Piazza et al. suggested that if refusal behaviour produces a "good outcome" for the child (e.g., the meal ends), refusal will continue.<sup>10</sup>

#### **Research Context**

The treatment strategies with the most scientific support are based on behaviour analysis. 11-16 Kerwin and Volkert and Piazza examined the research literature on treatment of pediatric feeding disorders to identify which treatments had enough scientific support to be labeled "effective." Kerwin and Volkert and Piazza found that behavioural interventions were the only treatments with enough scientific evidence to be labeled "effective." Similar analyses by Sharp and colleagues, Ledford and Gast, and Williams and colleagues supported those of Kerwin and Volkert and Piazza. 11,13-16

#### **Research Results and Conclusions**

Because children have feeding problems for a variety of reasons, treatment should focus on all of the components (i.e., biological, oral motor and psychological) that contribute to feeding problems and should be interdisciplinary.<sup>17,18</sup> One preliminary analysis of the outcomes for 50 children admitted to an interdisciplinary day-treatment feeding program indicated that over 87% of the goals for treatment were met by discharge from the program. When increases in calories consumed by mouth were the goal of treatment, 70% of patients reached their goal. Even when

children did not reach 100% of their oral intake goal, their levels of oral intake were increased substantially and within 20% of the goal One hundred percent of patients met their goals for increasing texture, decreasing bottle dependence, increasing self-feeding skills, and increasing variety of foods consumed.

All patients receiving their nutrition via tube experienced decreases in tube feedings and 70% of patients met their goals for decreases in tube feedings. Patients who entered the program with a *nasogastric tube* either left the program without the tube (75%), or the tube was removed shortly after discharge (100%). Ninety-seven percent of patients met their goals for decreasing inappropriate mealtime behaviour. Eighty-eight percent of caregivers were trained to implement the treatments with greater than 90% accuracy, and the treatment was transferred successfully to the home and community in 100% of cases.

Follow-up data indicated that the majority of patients continued to make progress toward agetypical feeding (e.g., volume increases *gastrostomy-tube* feeding decreases and initiation of self-feeding). Williams and colleagues, Greer and colleagues, and Laud and colleagues provided similar data in that interdisciplinary treatment with a behaviour analysis focus produces positive outcomes for children with severe feeding problems. 19-21

#### **Implications**

Interdisciplinary, intensive treatment of pediatric feeding disorders is successful in improving a wide variety of feeding problems, including dependence on supplemental feedings, selectivity by type and texture of food, inappropriate mealtime behaviour, failure to transition to age appropriate textures of food, and failure to self-feed to name a few. Successful treatment of these feeding problems has a number of important implications for children with feeding problems, their families, and society. Long-term, chronic feeding problems are associated with (a) health risks for the child, <sup>22</sup> (b) increased perceived stress for the child and family, <sup>23</sup> (c) mental health problems in families, <sup>24</sup> (d) increased risk of eating disorders such as anorexia, <sup>25</sup> and (e) increased health care costs for the child and family. <sup>19</sup> Therefore, treatment of pediatric feeding problems can result in (a) improved health of the child, (b) improved quality of life for the child and family, (c) decreased mental health problems in families, (d) reduced risk of long-term eating problems, and (e) reduced health care costs. Obviously, children who are dependent on technology such as gastrostomy-tubes (G-tube) for their nutritional needs have high health care costs. For example, the health care cost for a child with a G-tube is approximately \$41,811 for the first year. Over

two years, the health care cost for that child is estimated to be \$78,811 and after five years, the cost is \$189,811. These estimates are for uncomplicated care (e.g., no other significant medical problems related to the gastrostomy-tube) and do not include costs associated with family or individual therapy that may be necessary as a result of increased stress or psychopathology that has been documented in families of children with feeding problems. Moreover, the health care costs for these children may extend over many years if the child continues to need the gastrostomy-tube for nutrition or if the child develops eating problems such as anorexia later on. Williams and colleagues found that intensive, behavioural treatment was a cost-effective alternative to long-term supplemental feedings. Intensive, interdisciplinary treatment for feeding problems can eliminate the need for a gastrostomy-tube and result in age typical feeding, which can end the need for ongoing medical treatment in about 2 years. The estimated cost of intensive treatment for the feeding problem is approximately \$55,620 over 2 years. Thus, treatment of the feeding problems results in a savings of \$23,191 over a 2-year period and a cost savings of a minimum of \$134,191 over 5 years when compared to using a gastrostomy-tube for the problem.

Thus, not only are there obvious quality of life improvements for children with feeding problems and their families, but also there are significant cost savings when feeding problems are treated using interdisciplinary approaches with a behaviour-analytic focus.

#### References

- 1. Babbitt RL, Hoch TA, Coe DA, Cataldo MF, Kelly KJ, Stackhouse C, Perman JA. Behavioral assessment and treatment of pediatric feeding disorders. *Developmental and Behavioral Pediatrics* 1994;15(4):278-291.
- 2. Polan HJ, Kaplan MD, Kessler DB, Shindledecker R, Newmark M, Stern DN, Ward MJ. Psychopathology in mothers of children with failure to thrive. *Infant Mental Health Journal* 1991;12(1):55-64.
- 3. Field D, Garland M, Williams K. Correlates of specific childhood feeding problems. *Journal of Pediatric Child Health* 2003; 39: 299-304.
- 4. Gouge AL, Ekvall SW. Diets of handicapped children: Physical, psychological and socioeconomic correlations. *American Journal of Mental Deficiency* 1975;80(2):149-157.
- 5. Palmer S, Horn S. Feeding problems in children. In: Palmer S, Ekvall S, eds. *Pediatric nutrition in developmental disorders*. Springfield, Ill: Charles C. Thomas; 1978:107-129.
- 6. Perske R, Clifton A, McClean BM, Stein JI. *Mealtimes for severely and profoundly handicapped persons: New concepts and attitudes*. Baltimore, MD: University Park Press. 1977.
- 7. Burklow KA, Phelps AN, Schultz JR, McConnell K, Rudolph C. Classifying complex pediatric feeding disorders. *Journal of Pediatric Gastroenterology & Nutrition* 1998;27(2):143-147.
- 8. Rommel N, DeMeyer AM, Feenstra L, Veereman-Wauters G. The complexity of feeding problems in 700 infants and young children presenting to a tertiary care institution. Journal of Pediatric Gastroenterology and Nutrition 2003;37(1):75-84.

- 9. Troughton KE, Hill AE. Relation between objectively measured feeding competence and nutrition in children with cerebral palsy. *Developmental Medicine and Child Neurology* 2001;43(3):187-190.
- 10. Piazza CC, Fisher WW, Brown KA, Shore BA, Patel MR, Katz RM, Sevin BM, Gulotta CS, Blakely-Smith A. Functional analysis of inappropriate mealtime behaviors. *Journal of Applied Behavior Analysis* 2003;36(2):187-204.
- 11. Kerwin ME. Empirically supported treatments in pediatric psychology: severe feeding problems. *Journal of Pediatric Psychology* 1999;24(3):193-214.
- 12. Laud RB, Girolami PA, Boscoe J H, Gulotta C S. Treatment outcomes for severe feeding problems in children with autism spectrum disorders. *Behavior Modification* 2009; 33(5): 520-536.
- 13. Ledford JR, Gast DL. Feeding problems in children with autism spectrum disorders: A review. Focus on Autism and Other *Developmental Disabilities* 2006; 21: 153-166.
- 14. Sharp W G, Jaquess D L, Morton J F, Herzinger C V. Pediatric feeding disorders: A quantitative synthesis of treatment outcomes. *Clinical Child and Family Psychology Review* 2010.
- 15. Volkert VM, Piazza CC. Empirically supported treatments for pediatric feeding disorders. in: Sturmey P, Herson M, eds. *Handbook of Evidence Based Practice in Clinical Psychology*. Hoboken, NJ: Wiley, USA. in press
- 16. Williams KE, Field DG, Sieverling L. Food refusal in children: A review of the literature. *Research in Developmental Disabilities* 2010; 31: 625-633.
- 17. Cohen SA, Piazza CC, Navanthe A. Feeding and nutrition. In: Rubin IL, Crocker AC, eds. *Medical care for children and adults with developmental disabilities*. Baltimore, MD: Paul Brooks Publishing. 2006; 295-307.
- 18. Piazza CC. Feeding Disorders and behavior: What have we learned? *Developmental Disabilities Research Reviews* 2008; 14: 174-181.
- 19. Williams KE, Riegel K., Gibbons B, Field DG. Intensive behavioral treatment for severe feeding problems: A cost-effective alternative to tube feeding. *Journal of Developmental and Physical Disabilities* 2007; 19: 227-235.
- 20. Greer AJ, Gulotta CS, Masler EA, Laud RB. Caregiver stress and outcomes of children with pediatric feeding disorders treated in an intensive interdisciplinary program. *Journal of Pediatric Psychology* 2008; 33(6): 612-620.
- 21. Laud RB, Girolami PA, Boscoe JH, Gulotta CS. Treatment outcomes for severe feeding problems in children with autism spectrum disorder. *Behavior Modification* 2009; 33(5): 520-536.
- 22. Berezin S, Schwarz SM, Halata MS, Newman LJ. Gastroesophageal reflux secondary to gastrostomy tube placement. *American Journal of Diseases in Childhood* 1986;140(7):699-701.
- 23. Singer LT, Song L-Y, Hill BP, Jaffe AC. Stress and depression in mothers of failure-to-thrive children. *Journal of Pediatric Psychology* 1990;15(6):711-720.
- 24. Duniz M, Scheer PJ, Trojovsky A, Kaschnitz W, Kvas E, Macari S. Changes in psychopathology of parents of NOFT (non-organic failure to thrive) infants during treatment. *European Child and Adolescent Psychiatry* 1996;5(2):93-100.
- 25. Kotler LA, Cohen P, Davies M, Pine DS, Walsh TB. Longitudinal relationships between childhood, adolescent, and adult eating disorders. *Journal of the American Academy of Child & Adolescent Psychiatry* 2001;40(12):1434-1440.

#### **Genetic Influences on Child Eating Behaviour**

Clare Llewellyn, PhD, Jane Wardle, PhD

Health Behaviour Research Centre, University College London, United Kingdom September 2013

#### Introduction

Obesity is a global epidemic; its prevalence is increasing<sup>1</sup> and it is developing earlier in childhood.<sup>2</sup> It is therefore crucial to identify causal factors underlying early weight gain. Recent changes to the environment – such as increased opportunity to eat high calorie food and decreased opportunity to be physically active – have undoubtedly played a role. Nonetheless, not everyone is overweight. Body weight has a strong genetic basis,<sup>3,4</sup> leading to the hypothesis that genes may influence how vulnerable an individual is to gaining weight in the modern environment.<sup>5</sup> Genetically-determined susceptibility to the environment would help to explain how obesity can be both genetic and environmental.

#### **Subject**

Eating behaviour, or appetite, has been proposed as one mechanism through which genes influence obesity susceptibility.<sup>5,6</sup> In particular, individuals who inherit a more avid appetite – high responsiveness to external food cues and low sensitivity to internal satiety (fullness) processes – may be more likely to take advantage of the multiple eating opportunities presented by the modern environment, and consequently gain more weight. That is, genes may influence how big or small an individual's appetite is, and ultimately these genes therefore impact on their weight, so-called "inherited behavioural susceptibility to obesity."

#### **Problems and Research Context**

Eating behaviours of interest are those with evidence for a causal role in early life weight gain.<sup>7-11</sup> Broadly these can be thought of as "food approach" behaviours which indicate a bigger appetite and a greater interest in food (e.g., enjoyment of food<sup>11</sup> and responsiveness to external food cues <sup>8,11</sup>) and predispose to weight gain, and 'food avoidance' behaviours which suggest better appetite regulation and a lower interest in food (e.g., sensitivity to satiety<sup>9,11</sup> and slower eating speed<sup>7,10,11</sup>) and protect against weight gain. Observational measurement of eating behaviour

provides detailed information, but is time-consuming and expensive so the number of observations is limited, which is problematic for genetic research which requires large samples (see below). It is also possible that participants alter their eating behaviour in response to being observed, especially self-conscious overweight adults. Using children instead of adults helps get around this issue, and the development of a parent-report measure of child eating behaviour (the Child Eating Behavior Questionnaire [CEBQ]<sup>12,13</sup> or the Baby Eating Behavior Questionnaire [BEBQ]<sup>14</sup>) has enabled information to be collected reliably on very large samples, for a range of eating behaviours.

Two different approaches allow researchers to explore genetic influences on eating behaviour. So-called "quantitative genetic" studies estimate broadly the extent to which behaviour is influenced by genes or environment. 15 Studies compare family members who differ in their genetic relatedness - if relatives who are more genetically similar are also more similar for eating behaviour, genetic influence is inferred. Twins are commonly used because identical twins (monozygotic pairs, MZs) are 100% genetically identical, while non-identical or "fraternal" twins (dizygotic pairs, DZs) share on average only about 50% of their genes, like regular siblings. At the same time, MZs and DZs can be assumed to share environmental factors to a very similar extent (e.g., they are born at the same time into the same family), so they can be compared. Greater similarity between MZs compared to DZs suggests that genes influence eating behaviour. 15 Heritability is the statistic that is derived from twin studies, and it indicates the extent to which individual differences in the sample are explained by genetic variation. The statistic ranges from 0% (genetic variation does not contribute to individual differences) to 100% (individual differences can be explained entirely by genetic variation).[1] Twin studies are limited in that they cannot tell us anything about the actual genes involved, they simply indicate the relative importance of genes versus environment.

Molecular genetic studies try to identify the specific genes. Early research studied individuals and families with extreme manifestations of the trait – e.g., severe early onset obesity – to find genes. These studies identified genes responsible for rare and serious genetic disorders, but not common genes influencing variation across the general population (e.g., body weight). Recent technological advances combined with the completion of the Human Genome Project in 2000 have made it possible to explore the impact of millions of genetic variants on traits across the wider population, using a method called genome-wide association (GWAS). Large samples are needed for quantitative genetic studies and even larger samples for molecular genetic studies.

#### **Key Research Questions**

- 1. Is eating behaviour heritable?
- 2. Which genes influence eating behaviour?
- 3. Do weight-related genes influence weight via eating behaviour?

#### **Recent Research Results**

The high heritability of body weight in children and adults (~70%) is a long-standing finding established from a wealth of twin and family studies.<sup>3,4</sup> Similar heritability estimates have been found for infant and child eating behaviour. The CEBQ was used in 5,435 pairs of 10 year old British twins to demonstrate that the majority of individual differences in responsiveness to food cues and satiety responsiveness are determined by genes (75% and 63% respectively).<sup>5</sup> A similar estimate was obtained for an observational measure of eating speed (63%) in a subsample of 254 of these twins at age 11.<sup>18</sup> An infant-version of the CEBQ (the BEBQ<sup>19</sup>) showed comparably high heritability estimates for enjoyment of food (53%), food responsiveness (59%), satiety responsiveness (72%) and slowness in eating (84%), in 2,402 infant twin pairs.<sup>20</sup> The same sample of infant twins was also used to show that about a third of the genes that influence eating behaviours influence weight as well, supporting the idea that genes influence weight via their effects on eating behaviour.<sup>21</sup>

Early studies of obese individuals and families identified major mutations in one of a few genes resulting in severe early-onset obesity, as well as extreme appetite voraciousness and no apparent satiety.<sup>22</sup> These genes are fundamental regulators of the *leptin-melanocortin pathway* that controls hunger and satiety centrally, and is located in an ancient part of the brain called the hypothalamus. While the discovery of these genetic mutations provided important insights into the biology of body weight and eating behaviour, they are extremely rare and therefore do not explain weight variation at the population level.

Genome-wide association studies (GWAS) have identified more than 30 common genetic variants associated with body weight in adults and children.<sup>23</sup> The first to be discovered was the "fat mass and obesity-associated gene," FTO.<sup>24</sup> Nearly half of us carry at least one of the weight-related FTO variants; and those of us carrying two are on average 3 kilograms heavier than those carrying none. Not only is FTO expressed primarily in the hypothalamus, but it is associated with observational measures of food responsiveness measured at test meals,<sup>25</sup> and with satiety

responsiveness as measured by the CEBQ.<sup>26</sup> Many of the other variants discovered through GWAS are also expressed in the hypothalamus, suggesting that common genetic variants, like the rare mutations, influence body weight through eating behaviour. However, as yet relationships between these other common variants with eating behaviour have not been explored.

#### Research Gaps

Although research has made headway in establishing that eating behaviour, like weight, has a strong genetic basis, we still know very little about the specific genes involved. A good starting point would be to explore whether the weight-related common genes identified so far are also associated with eating behaviour. The biology linking genes to behaviour also needs to be characterized. In addition, it is crucial to test the feasibility of modifying eating behaviour in individuals who are genetically susceptible to obesity. There has been little research into modifying eating behaviour so far, but some studies are promising. Epstein and colleagues succeeded in slowing the eating speed of 7 year old children over a 6-month period by encouraging them to put their knife and fork down after each bite.<sup>27</sup> Another study demonstrated that it is possible to train 4-5 year old children to become better at recognizing and attending to their internal satiety sensations.<sup>28</sup>

#### **Conclusions**

Twins studies have established that differences between individuals in eating behaviour is explained partly by genetic variation, in typical western societies at this time; although the actual genes involved remain to be identified. Twin studies have also shown that the same genes which influence eating behaviour also influence weight, suggesting that the size of an individual's appetite ultimately influences their vulnerability to obesity in the modern permissive environment. In support of this, the most important common weight-related gene (FTO) appears to influence body weight through its effects on satiety sensitivity in children. This so-called inherited behavioural susceptibility to obesity helps to overcome the seeming paradox of both genetic and environmental determination of weight.

# Figure 1. Interactions between individual genetics susceptibility to obesity and the food environment




Figure 1 is a hypothetical demonstration of the percentage of children who are obese under three different environmental conditions, according to whether they have a low, average or high genetic susceptibility to obesity. Under conditions of famine there would be no obese children, regardless of genetic susceptibility to obesity; when the food supply is limited slightly more of the children at high genetic susceptibility would be obese than those at average susceptibility, but none of the children at low genetic susceptibility would be obese; under conditions of abundance (like the modern food environment in the U.S.) the majority of children at high genetic susceptibility would be obese, a considerable number at average genetic susceptibility, but very few children at low genetic susceptibility would be obese.

#### **Implications for Parents, Services and Policy**

This research suggests that some individuals are more vulnerable to overeating in response to the multiple opportunities presented by the modern food environment, by virtue of their genes, and more likely to gain weight. This calls into question the notion of personal responsibility for obesity. It is a commonly held belief that a child's weight reflects the rearing strategies of the parent. However, this research suggests that some children (more often those who have overweight parents and are therefore likely to have inherited higher risk genes.<sup>31</sup> may find it harder than others to regulate their food intake appropriately because they have a more avid

appetite, which is partly due to genes inherited from their parents. This can make it more difficult to resist over-consuming in response to the many eating opportunities presented by the modern environment.

[An effective strategy might be tighter state regulation of the wider food environment to reduce opportunities and incentives for overconsumption, such as removal of vending machines from schools, tighter control of food marketing to children, limitation of the number of fast food venues, and regulation of supermarket layouts.<sup>32</sup> Parents can also modify the home environment – e.g., by cooking smaller amounts of dinner to remove the temptation for "seconds," keeping problem foods out of sight (or better, out of the home), and teaching their children how to recognize feelings of fullness.

#### References

- 1. Finucane MM, Stevens GA, Cowan MJ, Danaei G, Lin JK, Paciorek CJ, Singh GM, Gutierrez HR, Lu Y, Bahalim AN, Farzadfar F, Riley LM, Ezzati M. National, regional, and global trends in body-mass index since 1980: systematic analysis of health examination surveys and epidemiological studies with 960 country-years and 9.1 million participants. *Lancet* 2011;377(9765):557-567.
- 2. Lee JM, Pilli S, Gebremariam A, Keirns CC, Davis MM, Vijan S, Freed GL, Herman WH, Gurney JG. Getting heavier, younger: trajectories of obesity over the life course. *Int J Obes (Lond)* 2010;34(4):614-623.
- 3. Elks CE, den Hoed M, Zhao JH, Sharp SJ, Wareham NJ, Loos RJ, Ong KK. Variability in the heritability of body mass index: a systematic review and meta-regression. *Front Endocrinol (Lausanne)* 2012;3:29.
- 4. Maes HH, Neale MC, Eaves LJ. Genetic and environmental factors in relative body weight and human adiposity. *Behav Genet* 1997;27(4):325-351.
- 5. Carnell S, Haworth CM, Plomin R, Wardle J. Genetic influence on appetite in children. *Int J Obes (Lond)* 2008;32(10):1468-1473.
- 6. Carnell S, Wardle J. Appetite and adiposity in children: evidence for a behavioral susceptibility theory of obesity. *Am J Clin Nutr* 2008;88(1):22-29.
- 7. Agras WS, Kraemer HC, Berkowitz RI, Korner AF, Hammer LD. Does a vigorous feeding style influence early development of adiposity? *J Pediatr* 1987;110(5):799-804.
- 8. Rodin J, Slochower J. Externality in the nonobese: effects of environmental responsiveness on weight. *J Pers Soc Psychol* 1976;33(3):338-344.
- 9. Parkinson KN, Drewett RF, Le Couteur AS, Adamson AJ. Do maternal ratings of appetite in infants predict later Child Eating Behavior Questionnaire scores and body mass index? *Appetite* 2010;54(1):186-190.
- 10. Stunkard AJ, Berkowitz RI, Schoeller D, Maislin G, Stallings VA. Predictors of body size in the first 2 y of life: a high-risk study of human obesity. *Int J Obes Relat Metab Disord* 2004;28(4):503-513.
- 11. van Jaarsveld CH, Llewellyn CH, Johnson L, Wardle J. Prospective associations between appetitive traits and weight gain in infancy. *Am J Clin Nutr* 2011;94(6):1562-1567.
- 12. Carnell S, Wardle J. Measuring behavioral susceptibility to obesity: validation of the child eating behavior questionnaire. *Appetite* 2007;48(1):104-113.

- 13. Wardle J, Guthrie CA, Sanderson S, Rapoport L. Development of the Children's Eating Behavior Questionnaire. *J Child Psychol Psychiatry* 2001;42(7):963-970.
- 14. Llewellyn CH, van Jaarsveld CH, Johnson L, Carnell S, Wardle J. Development and factor structure of the Baby Eating Behavior Questionnaire in the Gemini birth cohort. *Appetite* 2011;57(2):388-396.
- 15. Plomin R, DeFries JC, McClearn GE, McGuffin P. Behavioral Genetics. 5 ed. New York, US: Worth Publishers; 2008.
- 16. Faroogi IS. Genetic aspects of severe childhood obesity. Pediatr Endocrinol Rev 2006;3 Suppl 4:528-536.
- 17. Manolio TA. Genomewide association studies and assessment of the risk of disease. N Engl J Med 2010;363(2):166-176.
- 18. Llewellyn CH, van Jaarsveld CH, Boniface D, Carnell S, Wardle J. Eating rate is a heritable phenotype related to weight in children. *Am J Clin Nutr* 2008;88(6):1560-1566.
- 19. Llewellyn CH, van Jaarsveld CH, Johnson L, Carnell S, Wardle J. Development and factor structure of the Baby Eating Behavior Questionnaire in the Gemini birth cohort. *Appetite* 2011;57(2):388-396.
- 20. Llewellyn CH, van Jaarsveld CH, Johnson L, Carnell S, Wardle J. Nature and nurture in infant appetite: analysis of the Gemini twin birth cohort. *Am J Clin Nutr* 2010;91(5):1172-1179.
- 21. Llewellyn CH, van Jaarsveld CH, Plomin R, Fisher A, Wardle J. Inherited behavioral susceptibility to adiposity in infancy: a multivariate genetic analysis of appetite and weight in the Gemini birth cohort. *Am J Clin Nutr* 2012;95(3):633-639.
- 22. Barsh GS, Farooqi IS, O'Rahilly S. Genetics of body-weight regulation. Nature 2000;404(6778):644-651.
- 23. Speliotes EK, Willer CJ, Berndt SI et al. Association analyses of 249,796 individuals reveal 18 new loci associated with body mass index. *Nat Genet* 2010;42(11):937-948.
- 24. Frayling TM, Timpson NJ, Weedon MN et al. A common variant in the FTO gene is associated with body mass index and predisposes to childhood and adult obesity. *Science* 2007;316(5826):889-894.
- 25. Wardle J, Llewellyn C, Sanderson S, Plomin R. The FTO gene and measured food intake in children. *International Journal of Obesity* 2009;33(1):42-45.
- 26. Wardle J, Carnell S, Haworth CMA, Farooqi IS, O'Rahilly S, Plomin R. Obesity associated genetic variation in FTO is associated with diminished satiety. *J Clin Endocrinol Metab* 2008;93(9):3640-3643.
- 27. Epstein LH, Parker L, Mccoy JF, Mcgee G. Descriptive analysis of eating regulation in obese and nonobese children. *J Appl Behav Anal* 1976;9(4):407-415.
- 28. Johnson SL. Improving Preschoolers' self-regulation of energy intake. Pediatrics 2000;106(6):1429-1435.
- 29. Rokholm B, Silventoinen K, Tynelius P, Gamborg M, Sorensen TI, Rasmussen F. Increasing genetic variance of body mass index during the Swedish obesity epidemic. *PloS one* 2011;6(11):e27135.
- 30. Demerath EW. The genetics of obesity in transition. Coll Antropol 2012;36(4):1161-1168.
- 31. Whitaker KL, Jarvis MJ, Beeken RJ, Boniface D, Wardle J. Comparing maternal and paternal intergenerational transmission of obesity risk in a large population-based sample. *Am J Clin Nutr* 2010;91(6):1560-1567.
- 32. Gostin LO. Law as a tool to facilitate healthier lifestyles and prevent obesity. JAMA 2007;297(1):87-90.

# Screen time Use and Food Marketing Influences on Children's Eating Behaviors

<sup>1</sup>Chrisa Arcan, PhD, MPH, RD, <sup>2</sup>Aninda Sen, MBBS, MPH, <sup>3</sup>Meg Bruening, PhD, MPH, RD, <sup>4</sup>Mary Story, PhD, RD

<sup>1</sup>Renaissance School of Medicine, Stony Brook University, USA, <sup>2</sup>University of North Carolina at Chapel Hill, USA, <sup>3</sup>PennState College of Health and Human Development, USA, <sup>4</sup>Duke University, USA

October 2025

#### Introduction

Among children and adolescents, a key determinant of food preferences is taste which is shaped by early environmental exposure to food during infancy and childhood. Taste preferences are acquired through learning processes, including repeated exposure and positive messaging about various foods. Additionally, two of the strongest dietary predictors appear to be familiarity and sweetness of food. Repeated exposure and high frequency to food marketing, often of unhealthy foods, influences food preference, purchase requests, consumption, and importantly long term brand preferences, with negative health outcomes for youth. 5-7

Screentime, including time spent in front of a television (TV) and other digital platforms results in increased exposure to food advertising increasing the risk of overweight or obesity among children and adolescents. In recent years, the expansion of digital media platforms has caused a shift from TV viewership to digital media providing food companies with additional platforms to expand their food marketing strategies to children. This shift is more prevalent among adolescents whereby TV viewership has decreased while digital media use has increased leading the food and beverage companies to increase their investment in food advertising in digital media spaces. Trends also suggest that adolescents are more likely to have unhealthy diets compared to younger children, partly due to aggressive marketing practices. Despite the decrease in TV viewing among children in recent years, it still remains an important source of exposure to food advertising among these age groups. Additionally, TV advertisements for certain food products viewed by children exert a persuasive and motivating effect through visual, emotional, and sensorineural stimulation, exerting a large influence on purchase requests, purchases, and ultimate consumption of energy dense and unhealthy foods, compared to adults

viewing such advertisements, thus highlighting the importance of children's screentime towards shaping their dietary practices. 13,14

There are associations between increased screen time among children and adolescents and lower intakes of fruits and vegetables higher intakes of energy-dense snacks, beverages, and fast foods, and a higher percentage of energy intake from dietary fats and added sugars leading to subsequent obesity. <sup>5,15-17</sup> From 1999 to 2023, the prevalence of obesity among 2-19 year olds in the US has increased from 13.9% to 21.1%, while the prevalence of severe obesity has nearly doubled from 3.6% to 7%. <sup>18</sup> Studies have shown that due to psychological and neurological mechanisms affecting children's judgement, they tend to make more immediate food-related decisions than older adults following exposure to food advertisements. This leads to decisions favoring taste over health <sup>19</sup> leading to over-consumption of foods high in calories, refined sugars and salt, all of which are associated with outcomes such as obesity and other negative health outcomes. <sup>20</sup>

#### Prevalence and types of screentime use among children and adolescents

In the US, there has been a decline in traditional TV viewership by 57% and 60% among 2-11 years old children and 12-17 year old adolescents, respectively.<sup>21</sup> During this same period, the amount of entertainment-related screen time has increased among all age groups, with 8-12-year-old children and 13-18-year-old adolescents reporting 5.5 hours per day and 8 hours per day of screen time, respectively. This is largely attributed to digital media platforms, especially those accessed via mobile devices.<sup>21</sup>

With the growth of digital media and the use of mobile devises, the pervasiveness of advertising to children has increased as well. In the US, the most popular social media sites among children aged 5 years and older and adolescents aged 11-17 years are TikTok, Snapchat, and Instagram, while younger viewers aged 3-5 years primarily watch YouTube and YouTube kids. <sup>21,22</sup> This highlights that social media, video streaming, branded websites, and gaming sites take up the majority of younger viewers' screen time and have replaced traditional cable TV as the preferred alternative platform for entertainment access. In turn, the increase in screen time choices has presented food companies with a plethora of digital venues to engage in food marketing to children. Due to its diverse nature and unique characteristics, digital food marketing compared to traditional advertisement (TV, print media) presents a challenge in assessing children's exposures and impact on their health outcomes and in developing policies and solution to

address potential harms. The diversity and constantly evolving media platforms (e.g. mobile phones, gaming, social media, websites, etc.) and message sources allow for targeted and personalized messaging (e.g., sociodemographic) using creative approaches (e.g., influencers, branding, artificial intelligence, etc.) are a few of the characteristics of food digital marketing.<sup>21</sup>

#### Food and beverage marketing strategies and budget

Advertisements on TV, and more recently, on digital media platforms, have been identified as the two most popular venues where unhealthy, energy-dense, nutrient poor foods<sup>23</sup> are advertised to children and adolescents by employing sponsored content, influencers, data collection technology, and personalized behavioral marketing using machine learning.<sup>7,24</sup> To appeal to younger audiences, companies use strategies, such as celebrity endorsements, promotions, gifts / incentives (e.g., toys), games, branding, and other salient claims such as sporting prowess to promote products.<sup>7</sup> Studies indicate that in the US, food, beverage, and restaurant companies spend around \$14 billion each year for advertisements, with over 80% of that budget dedicated towards food which are generally considered unhealthy for children and adolescents, such as fast food, sugary drinks, candy, and snacks.<sup>25</sup> The food and beverage advertising budget dedicated to TV reaches \$11 billion annually.<sup>26</sup> In Canada, in 2019, advertisers spent an estimated \$628.6 million dollars on food advertising and 79.5% of this went to ads on television (67.7%) and digital media (11.8%). According to a study in Canada,<sup>27</sup> there were over 54 million food display ads (e.g., pop-up ads and banner ads) in one year on the most popular Canadian children's websites.

#### **Research Context**

Influence of TV viewing and digital use and advertising on eating behaviours of children

As one in five of U.S. children and adolescents have obesity and diets high in added sugars, sodium, and saturated fats are associated with a host of diet-related chronic diseases in adulthood, it is critical to examine the extent to which TV viewing and digital media use and food marketing negatively influence current and future eating behaviours among children and adolescents. Thus, it is important to discuss potential regulations that can protect children from TV and digital deceptive marketing practices. Finally, given the ubiquitous nature of food marketing using traditional and digital platforms, implications for policy makers and parents will be discussed.

#### **Key Research Questions**

- Is there an association between food marketing through traditional and digital platforms and eating behaviors in children and adolescents?
- Does advertising in TV and digital media platforms impact the quality of diet among children and adolescents? What are the effects of digital food marketing on children?
- Are there racial and socioeconomic differences in children's exposure to food marketing?
- Are TV viewing and digital media use and food advertising associated with diet-related health issues, such as obesity in children?
- What are the policy implications of existing marketing practices employed by food companies?

### The role of TV and digital platforms in shaping the diet quality of children and adolescents

The overall diet quality of children and especially adolescents remains low with over half of all youth ages 5-18 years in the US having poor dietary patterns with an average Healthy-Eating Index score of about 50 out of a possible 100.28 It has been documented that among young children, TV viewing is significantly associated with increased consumption of unhealthy foods, including fast food,29 increased requests of foods seen on TV, and more positive attitudes toward unhealthy foods.12,13,30-32 With the addition of digital food marketing, the increase in the overall screentime exposure has exposed children to more sources of high salt, sugar, and high energy-dense foods, including snacks, sugar-sweetened beverages, and fast food.433,34

While children younger than 7 years old display limited ability to recognize the persuasive nature of food advertisements, children from 7-14 tend to show greater awareness of the intent behind such advertisements. Despite this, adolescents are still undergoing cognitive development, and companies target this inability to comprehend disguised, persuasive marketing strategies, by influencing attitudes, food preferences, and poor impulse control. Adolescents are particularly vulnerable to celebrity endorsements, music sponsorships, social media and influencer-based marketing of potentially unhealthy food products leading to high levels of engagement and trust towards many brands across both traditional and digital marketing platforms. While adolescents might be more aware against such marketing tactics, their social desire to fit in and peer influence can overcome their skepticism towards media advertisements, thus creating a positive emotional associations to advertised products and making them feel more open to trying

out new foods and beverages. 32,36

Overall, both traditional TV advertising and digital food marketing have the same negative impact on children's food-related behaviors and more broadly on social norms and family relations; additionally, efforts to advertise healthy foods through digital media do not increase consumption of these foods neither do they decrease consumption of unhealthy foods. <sup>21</sup> Based on World Health Organization (WHO) reviews, unhealthy food marketing exposure increases immediate food intake and request for purchases. The frequent and amplified effects of food exposure through all multiple media platforms also has an effect on intermediate outcomes including brand awareness and loyalty, favorable ads, taste preferences, and purchasing intent for unhealthy foods. <sup>37</sup>

#### Racial and socioeconomic differences in children's exposure to food marketing

Targeted advertisements associated with TV and digital media usage can lead to disparities in diet-related health outcomes. Food and beverage advertising companies employ marketing practices targeting younger Hispanic and Black consumers in a disproportionate manner, which can have detrimental effects on the health of children and teenagers of ethnically diverse backgrounds. Reports indicate that fast food companies increased their budget for advertisements on Spanish speaking TV channels by 33% from 2012 to 2019, compared to a 9% increase across all channels combined. As a result of such discriminatory practices, viewing of fast-food advertisement among Hispanic children and adolescents increased in 2019. During the same time, Black children and adolescents also reported viewing 75% more fast food advertisements compared to their white counterparts, due to increased advertisements during TV programming popular among Black youth. Overall, despite the decrease in overall screen time associated with TV programming among all children and adolescents, the number of TV advertisements viewed by Black youth was higher compared to the number of advertisements viewed by youth in other racial groups.

Recognizing the rising presence of Black and Hispanic users in digital medial platforms, companies have also employed food marketing geared towards younger audiences from these population groups in recent years. 42 Recent studies found that social media accounts of food and beverage companies on platforms such as Facebook, Instagram, and TikTok have amassed significantly higher numbers of Hispanic and Black youth followers compared to social media accounts of other commercial products. Hispanic and Black youth have even reported feeling

more valued by food and beverage companies who employ racially targeted advertisements, which can lead to higher levels of online brand engagement compared to consumers from other racial backgrounds, affecting purchasing intent and actual purchases of unhealthy food and beverages. <sup>42</sup> These findings parallel the demographic differences in the prevalence of obesity among youth across different racial and ethnic groups, with Hispanic and Black children and adolescents being more likely to have obesity compared to their white counterparts. <sup>43,44</sup>

Children and adolescents in socioeconomically disadvantaged families are at a relatively higher risk of exposure to food and beverage advertisements from TV viewing compared to other groups. While the types of foods included both healthy and unhealthy items, the differences in exposure to unhealthy food advertisements is higher among younger viewers from lower socioeconomic backgrounds compared to the more affluent families.<sup>38</sup> This may be due to children from lower socioeconomic backgrounds spending more time in front of screens compared to children from higher socioeconomic backgrounds resulting in a subsequent doseresponse relationship, with higher screen-time and advertisement exposure leading to increased purchases and higher consumption of unhealthy food and beverages.<sup>38,40</sup>

Parental educational attainment has been shown to play a role on a child's exposure to food and beverage advertisements, especially on mobile devices, with children of lesser educated parents reporting higher advertisement exposure.<sup>38</sup> The reason for this discrepancy seems to be more related to the type of content viewed rather than duration of screen time, as families reporting lower socioeconomic status are found to be more likely to avoid higher subscription fees associated with subscription only streaming services. This results in more frequent commercial breaks in-between online entertainment, thus, greater exposure to food marketing practices.<sup>45</sup>

#### Policies to limit TV and digital media food advertising

The World Health Organization at the 63<sup>rd</sup> World Health Assembly in 2010, called for Member States to introduce policies that reduce the impact of advertising to children of foods high in sodium, sugars and saturated fat.<sup>46</sup> In 2023 WHO updated their statement calling for global action to reduce the impact of marketing of these foods to children. Specifically, they called for comprehensive and mandatory polices that reduce exposure of unhealthy food and beverage advertising among both children and adolescents based on nutrition criteria developed by governments.<sup>47</sup> Currently, 20 countries have mandatory policies, that vary in scope and enforcement.48 For example, in 2023, Norway's parliament agreed to new regulations to restrict

all advertisements of unhealthy foods to children under the age of 18 and will ban the marketing of unhealthy foods and beverages that are high in fats, sugars and salt in all media that children are likely to see, such as television, radio, the internet, and social media.<sup>47</sup> Chile and Portugal have the most comprehensive policies restricting all marketing and persuasive techniques in broadcast and non-broadcast platforms in children and adolescents.<sup>49</sup> Chile requires manufacturers to include front-of-package warning labels on prepackaged foods high in saturated fats, added sugars, and sodium or beverages, and bans their marketing to children under the age of 14 in media intended for children. These marketing restrictions apply to schools, product packaging, point-of-sale, print media, radio, television, and websites.<sup>47,48</sup>

In the US, in 2006, in partnership with the Council of Better Business Bureau's Children's Food and Beverage Advertising Initiative (CFBAI), a coalition of 21 of the largest food companies pledged to improve the nutritional quality of foods advertised to children under the age of 12 years. <sup>50</sup> Companies that participate in the CFBAI pledge that will only advertise foods and beverages that meet only "strict" nutrition criteria in "advertising primarily directed to children under age 13 in the U.S." Together, these stakeholders pledged to advertising only products which meet CFBAI criteria on children-directed media. <sup>51</sup> Prior evaluation of these policies has found that they excluded certain marketing venues, such as product packaging, in-store marketing, company own digital marketing, brand marketing etc. It is important to note that adolescents are not offered the same protections under the CFBAI increases their vulnerability for exposure to unhealthy food and the potential for altered food preferences, over-consumption, and subsequent health issues. <sup>51</sup>

In response to the rise in digital media usage among children, in 2018 the CFBAI updated their core principles to expand the digital media section of the agreement that was implemented in 2020. 52 Additionally, Children's Advertising Review Unit (CARU) is the primary self-regulatory mechanism for the advertising industry. 53 Disney and Google have established additional policies addressing food marketing to children on their own digital platforms.

Even after the updated CFBAI principles, analysis shows that the total number of products which are regulated by the CFBAI through self-regulatory practices across traditional TV and digital media platforms represent a very small portion of the overall calories consumed by children, which negates the possibility of such principles having any significant public health impact in curtailing childhood obesity rates. <sup>52</sup> Food and beverage companies continue to advertise products with low nutritional value to young audiences, encouraging unhealthy dietary habits that lead to

obesity and related negative health outcomes. 51,54 Recent studies also indicate that while many companies avoid marketing their brands on made-for-kids channels on platforms like YouTube and YouTube Kids, they can by-pass existing CFBAI guidelines through strategic product placements and endorsements of candy, sugar-sweetened beverages, salty snack and junk food on child-influencer videos. 55-57 In addition, the lack of protection for adolescents through the CFBAI policies, and the various vulnerabilities of this age group to targeted advertisement campaigns have so far not been addressed by any regulatory authorities as public health priority. Given the failure of the food industry's self-regulation initiatives, stronger policies are needed to protect children and adolescents from the harmful impact of food marketing. There is a need for government agencies in the US to implement and enforce mandatory regulatory policies on unhealthy marketing to children and youth.

In Quebec, the Consumer Protection Act prohibits commercial advertising, including all foods directed at children under the age of 13. In Canada, in 2016 the Minister of Health launched the Healthy Eating Strategy to help improve the food environment. Restricting food advertising to children is an important component of this Strategy. Health Canada considers the monitoring of food advertising to children as a priority and the Department of Health is monitoring the strategies that are most frequently used in advertising directed at children.<sup>59</sup>

#### **Research Findings and Gaps**

Some progress has been made in assessing the degree of exposure and content of TV and digital advertisement to children and adolescents, including expansion on newer popular media forms such as social media (e.g., Instagram), video streaming (e.g., Netflix, YouTube, TikTok), and livestream gaming (Twitch). Based on a panel report published by Healthy Eating Research (HER),<sup>21</sup> recent studies have expanded on the growing trend of influencer videos and other creative strategies in these platforms indicating the increased prevalence of branded posts and their impact on children. As adolescents engage more frequently with social media, recent studies have focused on adolescents compared to older studies that mostly included younger children. They have also observed that exposure to food advertising through influencers have immediate impact on young adolescents' consumption of unhealthy foods, despite providing ad disclosures.

Additional research gaps need to be addressed in future research.<sup>21</sup> In addition to traditional TV advertisements, the plethora of digital platforms and the constantly evolving creative marketing

strategies, require continuous research to understand the long-term impact of the cumulative and synergistic exposure on children's eating behaviors, health risks, and socio-cultural outcomes. While some research has been conducted to record unhealthy food marketing to racially and socioeconomically diverse children, more extensive research is needed to understand and address the impact of digital food marketing on the health inequities among these groups. It's important to develop programs addressing the impact of the expanding food marketing practices on children's eating behaviors as well as to evaluate the effectiveness of policies.

#### **Conclusions**

The types and amount of screen time that children and adolescents are exposed to has dramatically increased mainly due to the increase in the availability of digital entertainment platforms. While TV viewership has decreased, the increase in digital media has provided a plethora of food and beverage advertisements that amplified the persuasive massaging seen in traditional forms of media. Unfortunately, the vast majority of foods and beverages that are advertised to younger audiences across all digital platforms are unhealthy in nature<sup>7</sup> supporting the link between increased screentime, food marketing, and childhood and early life obesity.<sup>60</sup> The ubiquitous digital entertainment offerings in digital platforms and social media, (TikTok, Snapchat, Instagram) gaming platforms and video streaming services and the creative marketing strategies blur the lines between food promotion and entertainment with immediate and long term effect on food tastes and preferences of older children and adolescents.<sup>22,36</sup>

Youth from certain racial minority and socioeconomic backgrounds have been disproportionately affected as companies that cater to these audiences use targeted food marketing strategies in specific platforms and time-slots. Parental educational attainment has also been shown to contribute towards disproportionate exposure to unhealthy food advertisements among children and adolescents.<sup>38</sup> The exposure and impact of these diverse and targeted food marketing practices across different age, racial, and socioeconomic groups need to be further evaluated in order to develop policies aimed at protecting youth especially from vulnerable backgrounds.

There are no uniform policies and restrictions on food advertising to children internationally. <sup>61</sup> In North America, only Quebec and Mexico have mandatory restrictions on food advertising to children. <sup>59</sup>

#### **Implications for Parents, Providers, and Policy**

Parents' and caregivers' efforts to promote healthy eating at home are undermined by the high volume and influence of food advertisements that children see and hear where they learn, live, and play. Parents play an important role in the lives of young children, however the evolving landscape of food advertising to children undermines the parents' and caregivers' efforts to promote healthy eating to their children. When possible, parents can opt for child friendly plans when it comes to TV and digital media viewing, which will use high-quality media content that are considered less harmful in influencing unhealthy eating behavior. Clear and open communication between parents and children on how to assess marketing tactics on TV, social media, gaming websites and other screen-based platforms can equip young audiences with better skills for navigating unhealthy foods such as fatty foods, sugary-drinks, salted snacks, candy and others which can lead to both short and long-term negative health outcomes. Steps for building awareness within families of children who are disproportionately targeted by aggressive marketing will be key in shaping future dietary outcomes among minority and socioeconomically disadvantaged communities. School meetings, community events, and even places of worship can be used for disseminating information regarding digital media literacy, so that parents, guardians, and children of all ages can become better equipped in building their awareness of marketing strategies. Parents should also understand the importance of refraining from watching TV or using mobile devices during meals, removing TV from children's bedrooms, and generally limiting their children's exposure to TV and other devices. According to American Academy of Pediatrics, children below the age of 2 years should not watch TV and anyone older than 2 years should only watch 1 to 2 hours of quality programming per day. 62 Health care providers should be abreast of the latest research and policies regarding TV viewing and digital media use and children's dietary behaviours and obesity. At well-child visits, health care practitioner should discuss with families their media habits and inform them about the negative impact of food advertising on children's dietary behaviours.

Policy makers should monitor the media and strategies employed by food advertisers and collaborate with the food industry to develop evidence-based strategies and enforce regulatory guidelines to limit the TV and digital marketing of unhealthy products to children in all age groups. The regulatory principles can be extended in order to incorporate children 14 years old and older in order to offer added protection to adolescents during this crucial developmental stage in their lives, when lifelong dietary habits are being shaped and consolidated. Finally,

digital media platforms can be pressured in order to restrict their usage of personal data from mobile phones, tablets and other streaming and gaming devices which make it easier for advertisers to target certain groups for advertising food and beverages under the guise of social desirability and increased celebrity / influencer engagement.

#### References

- 1. Institute of Medicine (US) Committee on Food Marketing and the Diets of Children and Youth. Food Marketing to Children and Youth: Threat or Opportunity? Washington, DC: National Academies Press; 2006. doi:10.17226/11514
- Beckerman JP, Alike Q, Lovin E, Tamez M, Mattei J. The development and public health implications of food preferences in children. *Frontiers in Nutrition*. 2017;4:66. doi:10.3389/fnut.2017.00066
- 3. De Cosmi V, Scaglioni S, Agostoni C. Early taste experiences and later food choices. Nutrients. 2017;9(2):107. doi:10.3390/nu9020107
- 4. Scaglioni S, De Cosmi V, Ciappolino V, Parazzini F, Brambilla P, Agostoni C. Factors influencing children's eating behaviours. *Nutrients*. 2018;10(6):706. doi:10.3390/nu10060706
- 5. Boyland E, Backholer K, Potvin Kent M, et al. Unhealthy food and beverage marketing to children in the digital age: global research and policy challenges and priorities. *Annual Review of Nutrition*. 2024;44(1):471-497. doi:10.1146/annurev-nutr-062322-014102
- 6. Boyland EJ, Nolan S, Kelly B, et al. Advertising as a cue to consume: a systematic review and meta-analysis of the effects of acute exposure to unhealthy food and nonalcoholic beverage advertising on intake in children and adults. *American Journal of Clinical Nutrition*. 2016;103(2):519-533. doi:10.3945/ajcn.115.120022
- 7. World Health Organization. Food Marketing Exposure and Power and Their Associations
  With Food-Related Attitudes, Beliefs and Behaviours: A Narrative Review. Geneva: World
  Health Organization; 2022. Published February 7, 2022. Accessed June 17, 2025.
  https://www.who.int/publications/i/item/9789240041783
- 8. Fang K, Mu M, Liu K, He Y. Screen time and childhood overweight/obesity: a systematic review and meta-analysis. *Child: Care, Health and Development*. 2019;45(5):744-753. doi:10.1111/cch.12701

- 9. Elliott C, Truman E, Black JE. Tracking teen food marketing: participatory research to examine persuasive power and platforms of exposure. *Appetite*. 2023;186:106550. doi:10.1016/j.appet.2023.106550
- 10. Potvin Kent M, Soares Guimaraes J, Pritchard M, et al. Differences in child and adolescent exposure to unhealthy food and beverage advertising on television in a self-regulatory environment. *BMC Public Health*. 2023;23(1):555. doi:10.1186/s12889-023-15027-w
- 11. Liu J, Rehm CD, Onopa J, Mozaffarian D. Trends in diet quality among youth in the United States, 1999-2016. *JAMA*. 2020;323(12):1161-1174. doi:10.1001/jama.2020.0878
- 12. Ilieva RT, Gottlieb N, Christian H, Freudenberg N. Exposure to and impact of unhealthy food marketing on adolescents and young adults: a narrative review and research agenda. *Obesity Reviews*. Published online June 10, 2025:e13957. doi:10.1111/obr.13957
- 13. Boyland E. Would reducing children's exposure to food advertising prevent unhealthy weight gain? *Current Obesity Reports*. 2025;14(1):1-10. doi:10.1007/s13679-025-00648-6
- 14. Harris JL, Khanal B, Fleming-Milici F, Andreyeva T. Children's cereal purchases by U.S. households: associations with child versus adult TV ad exposure. *American Journal of Preventive Medicine*. 2025;68(3):598-606. doi:10.1016/j.amepre.2024.11.022
- Lapierre MA, Fleming-Milici F, Rozendaal E, McAlister AR, Castonguay J. The effect of advertising on children and adolescents. *Pediatrics*. 2017;140(Suppl 2):S152-S156. doi:10.1542/peds.2016-1758V
- 16. Robinson TN, Banda JA, Hale L, et al. Screen media exposure and obesity in children and adolescents. *Pediatrics*. 2017;140(suppl 2):S97-S101. doi:10.1542/peds.2016-1758K
- 17. Staiano AE, Button AM, Goldfield GS, et al. Screen media, obesity, and nutrition. In: Christakis DA, Hale L, eds. *Handbook of Children and Screens: Digital Media*, Development, and Well-Being from Birth Through Adolescence. Cham, Switzerland: Springer Nature; 2025:73-80. doi:10.1007/978-3-031-69362-5\_11
- QuickStats: Prevalence of obesity and severe obesity among persons aged 2-19 years United States, 1999-2000 through 2021-2023. MMWR. Morbidity and Mortality Weekly Report. 2024;73:941. doi:10.15585/mmwr.mm7341a5
- Bruce AS, Pruitt SW, Ha OR, et al. The influence of televised food commercials on children's food choices: Evidence from ventromedial prefrontal cortex activations. *Journal* of *Pediatrics*. 2016;177:27-32.e1. doi:10.1016/j.jpeds.2016.06.067

- Zhao L, Ogden CL, Yang Q, et al. Association of usual sodium intake with obesity among US children and adolescents, NHANES 2009-2016. *Obesity (Silver Spring)*. 2021;29(3):587-594. doi:10.1002/oby.23102
- 21. Healthy Eating Research. Evidence-based recommendations to mitigate harms from digital food marketing to children ages 2-17. Published 2024. Accessed June 9, 2025. https://healthyeatingresearch.org/research/digital-marketing-recommendations/
- 22. Anderson M, Perrin A. Children's engagement with digital devices, screen time. Pew Research Center. Published July 28, 2020. Accessed July 7, 2025. https://www.pewresearch.org/internet/2020/07/28/childrens-engagement-with-digital-devices-screen-time/
- 23. World Health Organization. Policies to Protect Children from the Harmful Impact of Food Marketing: WHO Guideline. Geneva, Switzerland: WHO; 2023. Accessed June 24, 2025. https://www.who.int/publications/i/item/9789240075412
- 24. Radesky J, Chassiakos YRL, Ameenuddin N, Navsaria D; Council on Communication and Media. Digital advertising to children. *Pediatrics*. 2020;146(1):e20201681. doi:10.1542/peds.2020-1681
- 25. UConn Rudd Center for Food Policy and Health. Food marketing. Published April 20, 2020. Accessed June 17, 2025. https://uconnruddcenter.org/research/food-marketing/
- 26. UConn Rudd Center for Food Policy and Health. TV & digital media food marketing. Published April 20, 2020. Accessed June 16, 2025. https://uconnruddcenter.org/research/food-marketing/tv-digitalmedia/
- 27. Potvin Kent M, Pauzé E, Bagnato M, et al. Food and beverage advertising expenditures in Canada in 2016 and 2019 across media. *BMC Public Health*. 2022;22(1):1458. doi:10.1186/s12889-022-13823-4
- 28. U.S. Department of Agriculture; U.S. Department of Health and Human Services. *Dietary Guidelines for Americans*, 2020-2025. 9th ed. Published 2020. Accessed July 24, 2025. https://www.dietaryguidelines.gov/resources/2020-2025-dietary-guidelines-online-materials
- 29. Chang HH, Nayga RM. Television viewing, fast-food consumption, and children's obesity. Contemporary Economic Policy. 2009;27(3):293-307. doi:10.1111/j.1465-7287.2008.00119.x

- 30. Chamberlain LJ, Wang Y, Robinson TN. Does children's screen time predict requests for advertised products? Cross-sectional and prospective analyses. Archives of Pediatrics & Adolescent Medicine. 2006;160(4):363-368. doi:10.1001/archpedi.160.4.363
- 31. Dixon HG, Scully ML, Wakefield MA, White VM, Crawford DA. The effects of television advertisements for junk food versus nutritious food on children's food attitudes and preferences. *Social Science & Medicine*. 2007;65(7):1311-1323. doi:10.1016/j.socscimed.2007.05.011
- 32. Harris JL, Sacco SJ, Fleming-Milici F. TV exposure, attitudes about targeted food ads and brands, and unhealthy consumption by adolescents: Modeling a hierarchical relationship. *Appetite*. 2022;169:105804. doi:10.1016/j.appet.2021.105804
- 33. Maksi SJ, Keller KL, Dardis F, et al. The food and beverage cues in digital marketing model: Special considerations of social media, gaming, and livestreaming environments for food marketing and eating behavior research. *Frontiers in Nutrition*. 2024;10:1325265. doi:10.3389/fnut.2023.1325265
- 34. Norman J, Kelly B, Boyland E, McMahon AT. The impact of marketing and advertising on food behaviours: Evaluating the evidence for a causal relationship. *Current Nutrition Reports*. 2016;5(3):139-149. doi:10.1007/s13668-016-0166-6
- 35. Fleming-Milici F, Harris JL. Adolescents' engagement with unhealthy food and beverage brands on social media. *Appetite*. 2020;146:104501. doi:10.1016/j.appet.2019.104501
- 36. Harris JL, Yokum S, Fleming-Milici F. Hooked on junk: Emerging evidence on how food marketing affects adolescents' diets and long-term health. *Current Addiction Reports*. 2021;8(1):19-27. doi:10.1007/s40429-020-00346-4
- 37. Kelly B, King L, Chapman K, Boyland E, Bauman AE, Baur LA. A hierarchy of unhealthy food promotion effects: Identifying methodological approaches and knowledge gaps.

  American Journal of Public Health. 2015;105(4):e86-e95. doi:10.2105/AJPH.2014.302476
- 38. Backholer K, Gupta A, Zorbas C, et al. Differential exposure to, and potential impact of, unhealthy advertising to children by socio-economic and ethnic groups: A systematic review of the evidence. *Obesity Reviews*. 2021;22(3):e13144. doi:10.1111/obr.13144
- 39. Cervi MM, Agurs-Collins T, Dwyer LA, Thai CL, Moser RP, Nebeling LC. Susceptibility to food advertisements and sugar-sweetened beverage intake in non-Hispanic Black and non-Hispanic White adolescents. *Journal of Community Health*. 2017;42(4):748-756.

#### doi:10.1007/s10900-016-0313-4

- 40. Coleman PC, Hanson P, van Rens T, Oyebode O. A rapid review of the evidence for children's TV and online advertisement restrictions to fight obesity. *Preventive Medicine Reports*. 2022;26:101717. doi:10.1016/j.pmedr.2022.101717
- 41. Harris JL, Fleming-Milici F, Phaneuf L, et al. Fast food advertising: Billions in spending, continued high exposure by youth. Rudd Center for Food Policy and Health. Published June 2021. Accessed July 2025. https://uconnruddcenter.org/research/fast-food-advertising/
- 42. Rummo PE, Arshonsky JH, Sharkey AL, Cassidy OL, Bragg MA. Social media accounts of food and beverage brands have disproportionately more Black and Hispanic followers than White followers. *Health Equity*. 2021;5(1):414-423. doi:10.1089/heq.2020.0068
- 43. Ogden CL, Fryar CD, Hales CM, Carroll MD, Aoki Y, Freedman DS. Differences in obesity prevalence by demographics and urbanization in US children and adolescents, 2013-2016. *Journal of the American Medical Association*. 2018;319(23):2410-2418. doi:10.1001/jama.2018.5158
- 44. Stierman B, Afful J, Carroll MD, et al. National Health and Nutrition Examination Survey 2017–March 2020 prepandemic data files—Development of files and prevalence estimates for selected health outcomes. *National Health Statistics Reports*. 2021;158:1-20. doi:10.15620/cdc:106273
- 45. Kenney EL, Mozaffarian RS, Norris J, Fleming-Milici F, Bleich SN. Estimating young children's exposure to food and beverage marketing on mobile devices. *Current Developments in Nutrition*. 2024;8(12):104505. doi:10.1016/j.cdnut.2024.104505
- 46. World Health Organization. Set of Recommendations on the Marketing of Foods and Non-Alcoholic Beverages to Children. WHO; 2010. Accessed July 24, 2025. https://www.who.int/publications/i/item/9789241500210
- 47. Childhood Healthy Living Foundation. What Are Other Countries Doing? Stop Marketing to Kids Coalition. 2024. Accessed July 24, 2025. https://stopmarketingtokids.ca/what-are-other-countries-doing/
- 48. World Health Organization. *Policies to Protect Children from the Harmful Impact of Food Marketing: WHO Guideline*. WHO; 2023.
- 49. Global Food Research Program. Countries With National Policies Regulating Food Marketing to Children. 2022. Accessed July 25, 2025.

#### https://www.globalfoodresearchprogram.org/

- 50. Kraak VI, Story M, Wartella EA, Ginter J. Industry progress to market a healthful diet to American children and adolescents. *American Journal of Preventive Medicine*. 2011;41(3):322-333. doi:10.1016/j.amepre.2011.05.029
- 51. BBB National Programs. *CFBAI Core Principles & Criteria* | *BBB National Programs*. 2024. Accessed June 23, 2025. https://bbbprograms.org/programs/children/cfbai/principles
- 52. Healthy Eating Research. Assessing the Public Health Impacts of the Children's Food and Beverage Advertising Initiative. August 2021. Accessed June 28, 2025. https://healthyeatingresearch.org/research/assessing-the-public-health-impacts-of-the-childrens-food-and-beverage-advertising-initiative/
- 53. BBB National Programs. *Children's Advertising Review Unit (CARU)* | *BBB National Programs*. March 24, 2025. Accessed July 8, 2025.

  https://bbbprograms.org/programs/children/caru
- 54. Jensen ML, Fleming-Milici F, Harris JL. Are U.S. food and beverage companies now advertising healthy products to children on television? An evaluation of improvements in industry self-regulation, 2017–2021. *International Journal of Behavioral Nutrition and Physical Activity.* 2023;20(1):118. doi:10.1186/s12966-023-01517-y
- 55. Fleming-Milici F, Phaneuf L, Harris J. Prevalence of food and beverage brands in "made-for-kids" child-influencer YouTube videos: 2019-2020. *Pediatric Obesity*. 2023;18(4):e13008. doi:10.1111/ijpo.13008
- 56. Fleming-Milici F, Gershman H, Agresta HO, McCann M, Harris J. Young children's (aged 3 to 8 years) food and beverage brand exposure on YouTube and YouTube Kids: An observational study and content analysis. *Journal of the Academy of Nutrition and Dietetics*. Published online June 25, 2025:S2212-2672(25)00188-1. doi:10.1016/j.jand.2025.05.010
- 57. Meyerding SGH, Marpert JD. Modern pied pipers: Child social media influencers and junk food on YouTube: A study from Germany. *Appetite*. 2023;181:106382. doi:10.1016/j.appet.2022.106382
- 58. Harris JL, Heard A, Schwartz MB. Older but still vulnerable: All children need protection from unhealthy food marketing. Published online 2014.
- 59. Health Canada. *Policy Update on Restricting Food Advertising Primarily Directed at Children: Policy Background*. April 25, 2023. Accessed July 24, 2025.

https://www.canada.ca/en/health-canada/services/food-nutrition/healthy-eating-strategy/policy-update-restricting-food-advertising-primarily-directed-children/policy-background.html

- 60. Jebeile H, Kelly AS, O'Malley G, Baur LA. Obesity in children and adolescents: epidemiology, causes, assessment, and management. *Lancet Diabetes & Endocrinology*. 2022;10(5):351-365. doi:10.1016/S2213-8587(22)00047-X
- 61. World Cancer Research Fund. *WCRF Policy Database*. Accessed July 24, 2025. https://policydatabase.wcrf.org/
- 62. American Academy of Pediatrics, Hill D, Ameenuddin N, et al. Media and young minds. *Pediatrics*. 2016;138(5):e20162591. doi:10.1542/peds.2016-2591

# Eating Behaviour in Studies of Child Growth, Development and Health - Measure Precisely, Early and in Context: Commentary on Ramsay, Liu & Stein, Black & Hurley, Milnes, Piazza & Carrol, Llewellyn & Wardle, and Arcan, Bruening and Story

Myles S. Faith, PhD

Gillings School of Global Public Health, University of North Carolina at Chapel Hill, USA September 2013

#### Introduction

Eating is essential for healthy growth and development. Along with physical activity, eating is the major behavioural conduit through which energy balance is regulated, through which food choices are made, and around which numerous interactions with family members and peers occur. Children's food choices and diet composition influence health status during development and, potentially, later life. Perhaps there is no better example of this than that of pediatric obesity, which tracks through adolescence and into adulthood and predicts a number of health disorders. For all these reasons, it is imperative that clinical guidelines and policy recommendations intended to foster healthy eating by children be guided by rigorous scientific studies and methodology. Quality measurement is fundamental to this endeavour and, as noted below, is a focal point of the papers<sup>1-6</sup> in this chapter.

The papers<sup>1-6</sup> reviewed in this section address a variety of topics related to the development of feeding and eating patterns throughout development, starting in infancy. The papers address biological and environmental drivers of refined eating phenotypes within "normal" populations as well as more specialized patient samples (e.g., feeding disorders).<sup>1-6</sup> They present some of the strongest scientific innovations in the pediatric ingestive behaviour fields, while highlighting real-world clinical and policy implications where real lives are touched. This is where the "rubber meets the road" of real-world eating behaviour.

A cross-cutting, if not unifying theme, across these papers<sup>1-6</sup> is the critical role of measurement in driving scientific advances. Armed with quality measurement, new insights have emerged which ultimately may better inform policy. Three conclusions can be drawn upon review of these

papers,<sup>1-6</sup> with respect to measurement: (1) Measure eating behaviour precisely; (2) Measure eating behaviour early; and (3) Measure eating behaviour in context. These are discussed next.

#### **Research and Conclusions**

Measure eating behaviour precisely

The contributions in this chapter<sup>1-6</sup> reveal insights that can be gleaned by using quality assessment tools that capture precise, refined dimensions of child eating behaviour. For example, as reviewed by Llewellyn and Wardle, development of the Child Eating Behavior Questionnaire over the past decade has yielded critical insights into the development of children's "food responsiveness" and "satiety awareness" - both of which are linked to childhood obesity.<sup>8,9</sup> The tool has been used internationally and allowed for replication of findings across diverse populations. This suggests novel targets for intervention (e.g., training children to recognize hunger/fullness cues) that, with rare exception, 10 have not been explored to date. Milnes et al.'s4 contribution illustrates how researchers can quantify specific parenting strategies in the context of pediatric feeding disorders (e.g., coaxing, reprimanding) and how these may impact on specific child eating patterns (e.g., bites of food, food refusal). Ramsay's¹ contribution illustrates how inefficient sucking, assessed by polygraph among young infants with gastroesophageal reflux, predicted poorer feeding skills and readiness for solid foods. Interestingly, there is other evidence that voracious sucking by infants is a risk factor for childhood obesity. 11,12 These examples underscore a point that is implicit throughout these contributions - but which needs explicit highlighting - that precise measurement has been essential for these scientific advances and will be critical to the next generation of discovery.

#### Measure eating behaviour early

As illustrated in this chapter,<sup>1-6</sup> eating behaviours can be measured reliably at earlier ages including infancy. This is illustrated by the Baby Eating Behaviour Questionnaire,<sup>13-15</sup> discussed by Llewellyn and Wardle,<sup>5</sup> the subscales of which may reflect early markers for obesity risk. As Black et al.<sup>3</sup> discuss, excess caloric intake during toddlerhood has been documented in population-based research, along with insufficient intake of fruits and vegetables and essential micronutrients. Ramsay's<sup>1</sup> contribution raises the topic of temperament and to what extent it might impact on feeding behaviour. Interestingly, there is emerging evidence that a "difficult" infant temperament in the first years of life may be risk factor for childhood obesity.<sup>16</sup> With the

development of new tools to assess eating behaviour in early life, researchers will be able to explore new causal models for (un)healthy growth and development.

#### Measure eating behaviour in context

The contributions to this chapter<sup>1-6</sup> reveal that eating behaviour occurs in a multi-level context. This context includes individual-level factors (e.g., genetics; temperament), as well as family-level factors (e.g., parental feeding practices; family relations) and societal/cultural factors (e.g., television commercials; laws regulating advertisements). This framework is consistent with the "socio-ecological model"<sup>17,18</sup> and has been endorsed by an American Heart Association task force on the promotion of healthy eating.<sup>19</sup> Understanding multi-level systems is scientifically challenging yet a pressing research and policy need.<sup>20</sup> This includes research linking genetics to individual child eating behaviour (as illustrated by Llewellyn and Wardle<sup>5</sup>) as well as individual child food preferences and intake to advertising policy (as illustrated by Arcan et al.<sup>6</sup>).

#### What's missing?

The contributions to this chapter<sup>1-6</sup> reveal the importance of measurement and its potential to provide refined characterizations of infant/toddler/child eating dimensions. This is profoundly important. Armed with these assessment tools, one of the greatest research needs can be better addressed: Better understanding the children for whom, and conditions under which, specific eating behaviours will promote or protect against disease onset. Answering these questions will require birth cohorts that are tracked across the development and – ideally – into adulthood. For example, are refined traits such as sucking intensity,<sup>12</sup> food neophobia,<sup>21</sup> food responsiveness,<sup>8</sup> and negative affect when eating<sup>22</sup> causally related to childhood obesity onset? If so, for whom and under what environmental conditions are these associations intensified or attenuated?

#### **Development and Policy Implications**

The authors provide thoughtful discussions regarding the policy implications for their respective topics. The findings, collectively, suggest an important implication for policy-level changes striving to modify child eating behaviour: It is unclear that all children necessarily will respond the same way to a given intervention. Whatever the policy may be (e.g., providing fruits and vegetables to schools; limiting the portion sizes of sugary beverages), one potentially should anticipate individual differences that drive non-uniform responses to the same policy. These individual differences might, conceivably, be related to factors such food responsiveness or

satiety awareness (which may have a sizable genetic loading), temperament, family interactions, neighborhood characteristics, or other unknown factors that influence how children eat. Being positioned to assess individual differences may help to reveal the conditions under which, and children for whom, certain policy changes will be more/less effective in fostering a healthier diet.

Finally, given the many factors that potentially impact on child eating behaviour, experimental studies that allow for strong causal inference are needed to guide policy development for healthy eating and obesity prevention. Indeed, laboratory studies and quasi-experimental designs can be enormously informative in this regard. This is illustrated by a recent review by Epstein et al.<sup>23</sup> which examined the impact of taxes and subsidies on food purchases. The review focused on studies using experimental designs, including laboratory-based investigations.

#### References

- Ramsay M. Feeding skill, appetite and feeding behaviours of infants and young children and their impact on growth and
  psychosocial development. Faith MS, topic ed. In: Tremblay RE, Boivin M, Peters RDeV, eds. Encyclopedia on Early
  Childhood Development [online]. Montreal, Quebec: Centre of Excellence for Early Childhood Development and Strategic
  Knowledge Cluster on Early Child Development; 2013:1-8. Available at: http://www.childencyclopedia.com/documents/RamsayANGxp2.pdf. Accessed May 15, 2013.
- Liu YH, Stein MT. Feeding behaviour of infants and young children and its impact on child psychosocial and emotional development. In: Tremblay RE, Barr RG, Peters RDeV, eds. Encyclopedia on Early Childhood Development [online].
   Montreal, Quebec: Centre of Excellence for Early Childhood Development; 2005:1-7. Available at: http://www.child-encyclopedia.com/documents/LiuSteinANGxp.pdf. Accessed May 15, 2013.
- 3. Black MM, Hurley KM. Helping children develop healthy eating habits. Faith MS, topic ed. In: Tremblay RE, Boivin M, Peters RDeV, eds. *Encyclopedia on Early Childhood Development* [online]. Montreal, Quebec: Centre of Excellence for Early Childhood Development and Strategic Knowledge Cluster on Early Child Development; 2013:1-10. Available at: <a href="http://www.child-encyclopedia.com/documents/Black-HurleyANGxp3-Eating.pdf">http://www.child-encyclopedia.com/documents/Black-HurleyANGxp3-Eating.pdf</a>. Accessed May 15, 2013.
- 4. Milnes SM, Piazza CC, Carroll TA. Assessment and treatment of pediatric feeding disorders. Faith MS, topic ed. In: Tremblay RE, Boivin M, Peters RDeV, eds. Encyclopedia on Early Childhood Development [online]. Montreal, Quebec: Centre of Excellence for Early Childhood Development and Strategic Knowledge Cluster on Early Child Development; 2013:1-5. Available at: http://www.child-encyclopedia.com/documents/Milnes-Piazza-CarrollANGxp2.pdf. Accessed May 15, 2013.
- 5. Llewellyn C, Wardle J. Genetic influences on child eating behaviour. Faith MS, topic ed. In: Tremblay RE, Boivin M, Peters RDeV, eds. *Encyclopedia on Early Childhood Development* [online]. Montreal, Quebec: Centre of Excellence for Early Childhood Development and Strategic Knowledge Cluster on Early Child Development; 2013:1-7. Available at: <a href="http://www.child-encyclopedia.com/documents/Llewellyn-WardleANGxp1.pdf">http://www.child-encyclopedia.com/documents/Llewellyn-WardleANGxp1.pdf</a>. Accessed May 15, 2013.
- 6. Arcan C, Bruening M, Story M. Television (TV) and TV advertisement influences on children's eating behaviour. Faith MS, topic ed. In: Tremblay RE, Boivin M, Peters RDeV, eds. Encyclopedia on Early Childhood Development [online]. Montreal, Quebec: Centre of Excellence for Early Childhood Development and Strategic Knowledge Cluster on Early Child Development; 2013:1-9. Available at: <a href="http://www.child-encyclopedia.com/documents/Arcan-Bruening-StoryANGxp1.pdf">http://www.child-encyclopedia.com/documents/Arcan-Bruening-StoryANGxp1.pdf</a>. Accessed May 15, 2013. Liu and Stein
- 7. Wardle J, Guthrie CA, Sanderson S, Rapoport L. Development of the Children's Eating Behaviour Questionnaire. *Journal of child psychology and psychiatry, and allied disciplines* 2001;42:963-70.

- 8. Carnell S, Wardle J. Appetite and adiposity in children: evidence for a behavioral susceptibility theory of obesity. *American journal of clinical nutrition* 2008;88:22-9.
- 9. Carnell S, Wardle J. Measuring behavioural susceptibility to obesity: validation of the child eating behaviour questionnaire. *Appetite* 2007;48:104-13.
- 10. Boutelle KN, Zucker NL, Peterson CB, Rydell SA, Cafri G, Harnack L. Two novel treatments to reduce overeating in overweight children: a randomized controlled trial. *Journal of consulting and clinical psychology* 2011;79:759-71.
- 11. Berkowitz RI, Moore RH, Faith MS, Stallings VA, Kral TV, Stunkard AJ. Identification of an obese eating style in 4-year-old children born at high and low risk for obesity. *Obesity* (Silver Spring) 2010;18:505-12.
- 12. Agras WS, Kraemer HC, Berkowitz RI, Korner AF, Hammer LD. Does a vigorous feeding style influence early development of adiposity? *Journal of pediatrics* 1987;110:799-804.
- 13. Johnson L, Llewellyn CH, van Jaarsveld CH, Cole TJ, Wardle J. Genetic and environmental influences on infant growth: prospective analysis of the Gemini twin birth cohort. *PloS one* 2011;6:e19918.
- 14. Llewellyn CH, van Jaarsveld CH, Johnson L, Carnell S, Wardle J. Nature and nurture in infant appetite: analysis of the Gemini twin birth cohort. *American journal of clinical nutrition* 2010;91:1172-9.
- 15. Llewellyn CH, van Jaarsveld CH, Johnson L, Carnell S, Wardle J. Development and factor structure of the Baby Eating Behaviour Questionnaire in the Gemini birth cohort. *Appetite* 2011;57:388-96.
- 16. Anzman-Frasca S, Stifter CA, Birch LL. Temperament and childhood obesity risk: a review of the literature. *Journal of developmental and behavioral pediatrics*: JDBP 2012;33:732-45.
- 17. Robinson T. Applying the socio-ecological model to improving fruit and vegetable intake among low-income African Americans. *Journal of community health* 2008;33:395-406.
- 18. Townsend N, Foster C. Developing and applying a socio-ecological model to the promotion of healthy eating in the school. *Public health nutrition* 2011:1-8.
- 19. Gidding SS, Lichtenstein AH, Faith MS, Karpyn A. Implementing American Heart Association pediatric and adult nutrition guidelines. *Circulation*. 2009;119: 1161-75.
- 20. Huang TT, Glass TA. Transforming research strategies for understanding and preventing obesity. *JAMA : Journal of the American Medical Association* 2008;300:1811-3.
- 21. Faith MS, Heo M, Keller KL, Pietrobelli A. Child food neophobiais heritable, associated with less compliant eating, and moderates familial resemblance for BMI. *Obesity* (Silver Spring) 2013.
- 22. Faith MS, Hittner JB. Infant temperament and eating style predict change in standardized weight status and obesity risk at 6 years of age. *Int J Obes* (Lond) 2010;34:1515-23.
- 23. Epstein LH, Jankowiak N, Nederkoorn C, Raynor HA, French SA, Finkelstein E. Experimental research on the relation between food price changes and food-purchasing patterns: a targeted review. *American journal of clinical nutrition* 2012;95:789-809.