

PHYSICAL ACTIVITY

Physical Activity Recommendations for Early Childhood

Rachel A. Jones, PhD, Anthony D. Okely, EdD

Early Start, Faculty of Social Sciences, University of Wollongong, Australia

February 2020, Éd. rév.

Introduction

Early childhood (0-5 years) is a critical time for the development of healthy behaviours, such as physical activity.¹ Participation in regular physical activity from birth prevents short- and long-term health complications, such as overweight and obesity, cardiovascular disease and musculoskeletal health.²⁻⁴ Furthermore, the promotion of physical activity should start as early as possible as physical activity levels track from early childhood to childhood and adolescence.⁵

Subject and Research Context

Given the international trend of less than optimal physical activity levels among young children, several countries have recently developed physical activity recommendations for the Early Years.⁶⁻¹⁰ Additionally, the World Health Organization has developed physical activity recommendations for children aged birth to 5 years.¹¹ These recommendations are evidence-based and provide a guide for how much and what type of physical activity is appropriate for infants (birth-1 year), toddlers (1-3 years) and preschool aged children (3-5 years). The majority of the guidelines also

recognize the importance of adequate sleep (in conjunction with physical activity and sedentary behaviours (specifically screen time) and are operationalized as 24-hour Movement Guidelines.^{6-8,10} All physical activity guidelines support the notion that physical activity is a natural and life-long activity that should be encouraged from birth. Parents and/or caregivers are encouraged to be positive role models and provide daily physical activity opportunities incorporating developmentally appropriate activities. Both structured and unstructured physical activity opportunities should be spread throughout the day and be provided in safe indoor and outdoor environments and the emphasis should be on “fun” and “participation” rather than competition.

Key Research Questions

The aim of this chapter is to summarize the empirical research supporting recently developed physical activity recommendations, in multiple countries, for children birth to 5 years of age.

The key research questions addressed in this chapter are:

1. What is the current evidence supporting the association between physical activity and health outcomes in early childhood?
2. Based on the current evidence, how much time should young children spend in physical activity and what type of physical activity should young children participate in.

Current evidence

Canada was one of the first countries to update their physical activity guidelines for children aged 0-5 years.⁷ Researchers involved in the development of these guidelines published a comprehensive systematic review which investigated the associations between physical activity and health-related outcomes.¹² This systematic review provides an excellent summary of the latest evidence.¹² Studies included in the review varied significantly in design and sample size and included randomized controlled trials, cross-sectional studies and longitudinal studies.¹² This chapter provides an overview of data presented in the review, as well as providing updated evidence since publication of the review. The association between physical activity and several health outcomes, namely, adiposity, motor development, psychosocial health, cognitive development, fitness, bone and skeletal health and cardiometabolic health are reviewed.

Fifty-seven studies investigated the relationship between physical activity and adiposity. Mixed results were reported with some studies reporting significant relationships between physical

activity and adiposity and others reporting no relationships. Mixed findings were reported irrespective of the design of the study. For example, of the 40 cross sectional studies, physical activity was positively associated with adiposity in 12 studies¹³⁻²⁴ and of the seven longitudinal studies physical activity was positively associated with adiposity in three studies.²⁵⁻²⁷ For all studies the relationship was stronger if a more direct measure of adiposity was used, for example, percent body fat as opposed to body mass index. The methodological quality of the studies, irrespective of design, was deemed as low or very low. Twenty-three studies investigated the association between physical activity and motor development. Seventy five percent of randomized controlled trials (n=4) reported positive associations²⁸⁻³⁰ as did 50% of clustered randomized controlled trials (n=2),³¹ 70% of cross-sectional studies (n=10)^{15,32-37} and 83% of the non-randomized interventions (n=6).³⁸⁻⁴²

In the Canadian systematic review,¹² less than 15 studies reported on the associations between physical activity and psychosocial health and cognitive development and less than 10 studies reported on the associations between physical activity and fitness, bone and skeletal health and cardiometabolic health. Two randomized controlled trials reported greater increases for psychosocial health outcomes in the intervention group compared to the control group^{28,43} and among the two longitudinal studies, physical activity, assessed as sport participation, was positively associated with psychosocial health in one study.⁴⁴ The association between physical activity and cognitive development were mixed. For the clustered randomized controlled trials (n=4), significant positive associations between physical activity and cognitive development were reported.⁴⁵⁻⁴⁸ These four studies had high methodological quality. Physical activity was positively associated with fitness in all studies (n=3).^{14,25,49} The majority (83%, 5/6) of cross-sectional studies reporting the association between physical activity and bone and skeletal health identified significant associations.⁵⁰⁻⁵⁴ Nine studies assessed the association between physical activity and cardiometabolic health. Mixed results were reported for studies investigating the relationship between blood pressure, cholesterol or triglycerides in both longitudinal studies and cross-sectional studies.^{26,55}

In summary, there is now more convincing evidence to suggest that physical activity of at least moderate- to vigorous-intensity is consistently associated with better health indicators.^{13,14,56,57} Furthermore, the evidence pertaining to younger children (0-2 years) has increased thus highlighting the importance of participating in physical activity from birth. Based on these findings the following guidelines were recommended (Table 1). The new guidelines specify the intensity of physical activity (i.e., whether the activity is light, moderate or vigorous) for preschool aged

children. Sixty minutes of moderate- to vigorous-intensity, also termed energetic play that results in ‘huff and puff’ is now recommended as part of the total 180 minutes per day.⁶⁻¹¹ Additionally, the most recent guidelines recommend a time for tummy time for infants (i.e., 30 minutes per day, accumulated throughout the day).

Table 1: Country-specific physical activity recommendations for children birth to five years

Country	Infants	Toddlers	Preschoolers
Australia* (released 2017) Infants: 0-1 year Toddlers: 1-3 years Preschoolers: 3-5 years ⁶	Be physically active several times in a variety of ways, particularly through interactive floor-based play; more is better. For those not yet mobile, this includes at least 30 minutes of tummy time spread throughout the day while awake.	At least 180 minutes spent in a variety of physical activities including energetic play, spread throughout the day; more is better.	At least 180 minutes spent in a variety of physical activities, of which at least 60 minutes is energetic play, spread throughout the day; more is better.
Canada* (released 2017) Infants: 0-1 year Toddlers: 1-2 years Preschoolers: 3-4 years ⁷	Be physically active several times in a variety of ways, particularly through interactive floor-based play—more is better. For those not yet mobile, this includes at least 30 minutes of tummy time spread throughout the day while awake.	At least 180 minutes spent in a variety of physical activities at any intensity, including energetic play, spread throughout the day—more is better.	At least 180 minutes spent in a variety of physical activities spread throughout the day, of which at least 60 minutes is energetic play—more is better.

New Zealand* Provide fun activities that support physical, social, emotional and spiritual growth (at least three hours every day for toddlers and preschoolers, Recommendations for spread throughout the day). Include plenty of opportunities for active play: that develop movement competence and confidence; that provide sufficient challenges to build resilience and encourage creativity through exploration; where children are by themselves as well as interacting with others, such as parents, siblings, friends, whanau/family and other caregivers that include a variety of indoor and outdoor activities, especially activities involving nature.

(released 2017)
Recommendations for specific ages groups not provided⁸

<p>United Kingdom (released 2019) Infants: 0-1 year Toddlers: 1-2 years Preschoolers: 3-4 years⁹</p>	<p>Babies should be encouraged to be active throughout the day, every day in a variety of ways, including crawling. If they're not yet crawling, encourage them to be physically active by reaching and grasping, pulling and pushing, moving their head, body and limbs during daily routines, and during supervised floor play. Try to include at least 30 minutes of tummy time spread throughout the day when they're awake. Once babies can move around, encourage them to be as active as possible in safe and supervised play environments.</p>	<p>Toddlers should be physically active every day for at least 180 minutes (3 hours). The more the better. This should be spread throughout the day, including playing outdoors. The 180 minutes can include light activity such as standing up, moving around, rolling and playing, as well as more energetic activity like skipping, hopping, running and jumping. Active play, such as using a climbing frame, riding a bike, playing in water, chasing games and ball games, is the best way for this age group to get moving.</p>	<p>Pre-schoolers should spend at least 180 minutes (3 hours) a day doing a variety of physical activities spread throughout the day, including active and outdoor play. The more the better. The 180 minutes should include at least 60 minutes of moderate-to-vigorous intensity physical activity. Children under 5 should not be inactive for long periods, except when they're asleep. Watching TV, travelling by car, bus or train, or being strapped into a buggy for long periods are not good for a child's health and development.</p>
--	--	--	---

<p>South Africa* (released 2018) Infants: 0-1 year Toddlers: 1-3 years Preschoolers: 3-5 years¹⁰</p>	<p>Be physically active several times a day in a variety of ways through interactive floor-based play, including crawling. For babies not yet mobile, this included at least 30 minutes of tummy time spread throughout the day while awake, and other movements such as reaching and grasping.</p>	<p>At least 180 minutes spent in a variety of physical activities including energetic play, spread through the day, more is better.</p>	<p>At least 180 minutes spent in a variety of physical activities of which at least 60 minutes is energetic play that raises heart rate and makes children “huff and puff” (i.e., running, jumping, dancing), spread throughout the day, more is better</p>
--	---	---	---

<p>World Health Organization* (released 2019) Infants: 0-1 year Toddlers: 1-2 years Preschoolers: 3-4 years¹¹</p>	<p>Be physically active several times a day in a variety of ways, particularly through interactive floor-based play; more is better. For those not yet mobile, this includes at least 30 minutes in prone position (tummy time) spread throughout the day while awake.</p>	<p>Spend at least 180 minutes in a variety of types of physical activities at any intensity, including moderate-to-vigorous-intensity physical activity, spread throughout the day; more is better.</p>	<p>Spend at least 180 minutes in a variety of types of physical activities at any intensity, of which at least 60 minutes is moderate- to vigorous intensity physical activity, spread throughout the day; more is better.</p>
---	--	---	--

*Included as part of 24-hour Movement behavior Guidelines

Research Gaps

Since the release of the revised country-specific physical activity recommendations, substantial international collaborative work has continued to address the research gaps. For example, surveillance studies which assess the proportion of children who meet the recommendations are

currently underway. The SUNRISE study is currently investigating this in 31 countries; two thirds of which are low- or middle-income countries. Data collection for the pilot study will be completed in 2020 (<https://sunrise-study.com>). Simple well-designed interventions have been reported, for example a recent study showed that by increasing the number of scheduled outdoor times in early childhood education and care settings, children spent significant more time in moderate- to vigorous-intensity physical activity.⁵⁸ Further innovative studies are called for to ensure optimal levels of physical activity are achieved.

Additional studies monitoring the awareness and uptake of the recommendations by stakeholders such as health professionals, childcare workers and parents are needed, however this type of investigation is often hindered by difficulty in securing funding. There have been small gains in this area, with some countries successfully increasing awareness of the recommendations among key stakeholders, however progress is generally slow.

Many studies which provided the evidence for the revised country-specific guidelines were low methodological quality. Based on the GRADE framework, in the Carson review,¹² only a few studies were deemed to have moderate or high methodological quality.^{28,45} It is important that all studies are methodologically sound to ensure the most robust evidence is provided which will in turn better inform policy and practice.

Conclusions

Life-long physical activity habits need to be established in the first five years of life as participation in regular physical activity has many health benefits. The release of a number of country-specific physical activity recommendations in the last five years provides clear recommendations on the appropriate type, intensity and amounts of physical activity for young children. Adhering to such recommendations will enhance health outcomes for children and provide the best possible start. Establishing healthy physical activity habits from a young age, through adherence to evidence-based physical activity recommendations, will be beneficial.

Implications for Parents, Services and Policy

The development of physical activity recommendations for children from birth to 5 years will have several notable implications for parents, services and policy makers. Current prescriptive physical activity recommendations, which are based on solid empirical evidence, will:

1. Assist key stakeholders to understand the importance of physical activity for health benefits among young children;
2. Inform government policy in relation to health-promoting physical activity for children birth to 5 years of age;
3. Assist consumers, childcare workers and other health professionals to understand the importance of physical activity for health in children; and
4. Underpin and support health promotion activities and intervention by workers across a range of sectors and all levels of government.

References

1. Ward DS, Vaughn A, McWilliams C, Hales D. Interventions for increasing physical activity at childcare. *Medicine and Science in Sports and Exercise*. 2010;42:526-534.
2. Timmons BW, LeBlanc AG, Carson V, Connor Gorber S, Dillman C, Janssen I, Kho ME, Spence JC, Stearns J, Tremblay MS. Systematic review of physical activity and health in the early years (aged 0-4 years). *Applied Physiology Nutrition and Metabolism*. 2012;37:773-792.
3. LeBlanc AG, Spence JC, Carson V, Connor Gorber S, Dillman C, Janssen I, Kho ME, Stearns J, Timmons BW, Tremblay MS. Systematic review of sedentary behavior and health indicators in the early years (aged 0-4 years). *Applied Physiology Nutrition and Metabolism*. 2012;37:753-772.
4. Janz K, Burns T, Levy S. Tracking of activity and sedentary behaviors in childhood. The Iowa Bone Development Study. *American Journal of Preventive Medicine*. 2005;29:171-178.
5. Jones RA, Hinkley T, Okely AD, Salmon J. Tracking physical activity and sedentary behavior in childhood: a systematic review. *American Journal of Preventive Medicine*. 2013;44:651-658. doi:10.1016/j.amepre.2013.03.001.
6. Australian Government. Department of Health. *Australian 24-Hour movement guidelines for children and young people (5-17 years): an integration of physical activity, sedentary behaviour and sleep*. <https://www1.health.gov.au/internet/main/publishing.nsf/Content/health-24-hours-phys-act-guidelines> Updated April 12, 2019. Accessed November 2019.
7. Canadian Society for Exercise Physiology. *Canadian 24-hour movement guidelines: an integration of physical activity, sedentary behavior and sleep*. <https://csepguidelines.ca>. Accessed November 2019.
8. New Zealand Government. Ministry of Health. *Sit less, move more, sleep well: active play guidelines for under-fives*. <https://www.health.govt.nz/publication/sit-less-move-more-sleep-well-active-play-guidelines-under-fives> Published May 31, 2017. Accessed November 2019.
9. UK National Centre for Sport and Exercise Medicine. UK Physical activity guidelines for early years. <https://www.laureus.co.za/moving-playing-sleeping-starting-early-with-healthy-habits/>. Accessed November 2019.
10. Laureus. *Moving, playing, sleeping: starting early with health habits*. <https://www.laureus.co.za>. Accessed November 2019.
11. World Health Organization. Guidelines on physical activity, sedentary behavior and sleep for children under 5 years of age. <https://www.who.int/publications-detail/guidelines-on-physical-activity-sedentary-behaviour-and-sleep-for-children-under-5->

years-of-age. Published April 2, 2019. Accessed November 2019.

12. Carson V, Lee EY, Hewitt L, Jennings C, Hunter S, Kuzik N, et al. Systematic review of the relationships between physical activity and health indicators in the early years (0-4 years). *BMC Public Health* 2017;17:854. doi:10.1186/s12889-017-4860-0
13. Eijkemans M, Mommers M, de Vries SI, van Buuren S, Stafleu A, Bakker I, Thijs C. Asthmatic symptoms, physical activity, and overweight in young children: a cohort study. *Pediatrics*. 2008;121(3):e666-e672.
14. Leppänen M, Nyström CD, Henriksson P, Pomeroy J, Ruiz J, Ortega F, Pomeroy J, Ruiz JR, Löf M. Physical activity intensity, sedentary behavior, body composition and physical fitness in 4-year-old children: results from the MINISTOP trial. *International Journal of Obesity*. 2016;40:1126-1133.
15. Lin LY, Cherng RJ, Chen YJ. Relationship between time use in physical activity and gross motor performance of preschool children. *Australian Occupational Therapy Journal*. 2016;64:49-57. doi:10.1111/1440-1630.12318
16. Pallan MJ, Adab P, Sitch AJ, Aveyard P. Are school physical activity characteristics associated with weight status in primary school children? A multilevel cross-sectional analysis of routine surveillance data. *Archives of Disease in Childhood*. 2014;99(2):135-141.
17. Ansari A, Pettit K, Gershoff E. Combating obesity in head start: outdoor play and change in children's body mass index. *Journal of Developmental and Behavioral Pediatric*. 2015;36(8):605-612.
18. Lioret S, Maire B, Volatier J, Charles M. Child overweight in France and its relationship with physical activity, sedentary behaviour and socioeconomic status. *European Journal of Clinical Nutrition*. 2007;61(4):509-516.
19. Trost SG, Sirard JR, Dowda M, Pfeiffer KA, Pate RR. Physical activity in overweight and nonoverweight preschool children. *International Journal of Obesity*. 2003;27(7):834-839.
20. Kagamimori S, Yamagami T, Sokejima S, Numata N, Handa K, Nanri S, Saito T, Tokui N, Yoshimura T, Yoshida K. The relationship between lifestyle, social characteristics and obesity in 3-year-old Japanese children. *Child Care Health and Development*. 1999;25(3):235-247.
21. Nelson JA, Carpenter K, Chiasson MA. Diet, activity, and overweight among preschool-age children enrolled in the special supplemental nutrition program for women, infants, and children (WIC). *Preventive Chronic Disease*. 2006;3(2):1-12.
22. Chen LP, Ziegenfuss JY, Jenkins SM, Beebe TJ, Ytterberg KL. Pediatric obesity and self-reported health behavior information. *Clinical Pediatrics*. 2011;50(9):872-875.
23. Shapiro LR, Crawford PB, Clark MJ, Pearson DL, Raz J, Huenemann RL. Obesity prognosis: a longitudinal study of children from the age of 6 months to 9 years. *American Journal of Public Health*. 1984;74(9):968-972.
24. Jones RA, Okely AD, Gregory P, Cliff DP. Relationships between weight status and child, parent and community characteristics in preschool children. *International Journal Pediatric Obesity*. 2009;4(1):54-60.
25. DuRant RH, Baranowski T, Rhodes T, Gutin B, Thompson WO, Carroll R, Greaves KA. Association among serum lipid and lipoprotein concentrations and physical activity, physical fitness, and body composition in young children. *Journal of Pediatrics*. 1993;123(2):185-192.
26. Klesges RC, Klesges LM, Eck LH, Shelton ML. A longitudinal analysis of accelerated weight gain in preschool children. *Pediatrics*. 1995;95(1):126-130.
27. Sijtsma A, Sauer PJ, Stolk RP, Corpeleijn E. Infant movement opportunities are related to early growth—GECKO Drenthe cohort. *Early Human Development*. 2013;89(7):457-461.
28. Porter LS. The impact of physical-physiological activity on infants' growth and development. *Nursing Research*. 1972;21(3):210-219.
29. Teixeira Costa HJ, Abelairas-Gomez C, Arufe-Giráldez V, Pazos-Couto JM, Barcala-Furelos R. Influence of a physical education plan on psychomotor development profiles of preschool children. *Journal of Human Sport Exercise*. 2015;10(1):126-140.

30. Mostafavi R, Ziaee V, Akbari H, Haji-Hosseini S. The effects of spark physical education program on fundamental motor skills in 4-6 year-old children. *Iran Journal of Pediatrics*. 2014;23(2):216-219.
31. Jones RA, Riethmuller A, Hesketh K, Trezise J, Batterham M, Okely AD. Promoting fundamental movement skill development and physical activity in early childhood settings: a cluster randomized controlled trial. *Pediatric Exercise Science*. 2011;23(4):600-615.
32. Williams HG, Pfeiffer KA, O'Neill JR, Dowda M, McIver KL, Brown WH, Pate RR. Motor skill performance and physical activity in preschool children. *Obesity*. 2008;16(6):1421-1426.
33. Pfeiffer KA, Dowda M, McIver KL, Pate RR. Factors related to objectively measured physical activity in preschool children. *Pediatric Exercise Science*. 2009; 21(2):196-208.
34. Kuo Y-L, Liao H-F, Chen P-C, Hsieh W-S, Hwang A-W. The influence of wakeful prone positioning on motor development during the early life. *Journal of Development and Behavioral Pediatrics*. 2008;29(5):367-376.
35. de Kegel A, Peersman W, Onderbeke K, Baetens T, Dhooge I, Van Waelvelde H. New reference values must be established for the Alberta infant motor scales for accurate identification of infants at risk for motor developmental delay in Flanders. *Child Care Health and Development*. 2013;39(2):260-267.
36. Dudek-Shriber L, Zelazny S. The effects of prone positioning on the quality and acquisition of developmental milestones in four-month-old infants. *Pediatric Physical Therapy*. 2007;19(1):48-55.
37. Fisher A, Reilly JJ, Kelly LA, Montgomery C, Williamson A, Paton JY, Grant S. Fundamental movement skills and habitual physical activity in young children. *Medicine Science and Sports Exercise*. 2005;37(4):684-688.
38. Krombholz H. The impact of a 20-month physical activity intervention in child care centers on motor performance and weight in overweight and healthy-weight preschool children. *Perceptual and Motor Skills*. 2012;115(3):919-932.
39. Draper CE, Achmat M, Forbes J, Lambert EV. Impact of a community-based programme for motor development on gross motor skills and cognitive function in preschool children from disadvantaged settings. *Early Child Development and Care*. 2012;182(1):137-152.
40. Livonen S, Sääkslahti A, Nissinen K. The development of fundamental motor skills of four- to five-year-old preschool children and the effects of a preschool physical education curriculum. *Early Child Development and Care*. 2011;181(3):335-343.
41. Venetsanou F, Kambas A. How can a traditional Greek dances programme affect the motor proficiency of pre-school children? *Research in Dance Education*. 2004;5(2):127-138.
42. Sigmundsson H, Hopkins B. Baby swimming: exploring the effects of early intervention on subsequent motor abilities. *Child Care Health and Development*. 2010;36(3):428-430.
43. Lobo YB, Winsler A. The effects of a creative dance and movement program on the social competence of head start preschoolers. *Social Development*. 2006;15(3):501-519.
44. Vella SA, Cliff DP, Magee CA, Okely AD. Associations between sports participation and psychological difficulties during childhood: a two-year follow up. *Journal of Science and Medicine in Sport*. 2015;18(3):304-309.
45. Mavilidi M-F, Okely AD, Chandler P, Cliff DP, Paas F. Effects of integrated physical exercises and gestures on preschool children's foreign language vocabulary learning. *Educational Psychology Review*. 2015;27(3):413-426.
46. Mavilidi M-F, Okely AD, Chandler P, Pass F. Infused physical activity into the classroom: effects on preschool children's geography learning. *Mind Brain and Education*. 2016;10(4):256-263.
47. Mavilidi M-F, Okely AD, Chandler P, Pass F. Effects of Integrating Physical Activities into a Science Lesson on Preschool Children's Learning and Enjoyment. *Applied Cognitive Psychology*. 2017;31(3):281-290.
48. Mavilidi M-F, Okely AD, Chandler P, Louise Somazet S, Pass F. Immediate and delayed effects of integrating physical activity into preschool children's learning of numeracy skill. *Journal of Experimental Child Psychology*. 2018;166:502-519.

49. Kolpakov V, Bespalova T, Tomilova E, Larkina NY, Mamchits E, Chernogrivova M, Kopytov AA. Functional reserves and adaptive capacity of subjects with different levels of habitual physical activity. *Human Physiology*. 2011;37(1):93-104.
50. Xu H, Zhao Z, Wang H, Ding M, Zhou A, Wang X, Zhang P, Duggan C, Hu FB. Bone mineral density of the spine in 11,898 Chinese infants and young children: a cross-sectional study. *PLoS One*. 2013;8(12):e82098.
51. Jazar AS, Takturi HR, Khuri-Bulos NA. Vitamin D status in a sample of preschool children aged from 1 to 6 years visiting the pediatrics clinic at Jordan University hospital. *Jordan Medical Journal*. 2012;45(4):308-316.
52. Kensarah OA, Jazar AS, Azzeh FS. Hypovitaminosis D in healthy toddlers and preschool children from western Saudi Arabia. *International Journal of Vitamin and Nutrition Research*. 2015;85:50-60.
53. Harvey N, Cole Z, Crozier S, Kim M, Ntani G, Goodfellow L, Robinson SM, Inskip HM, Godfrey KM, Dennison EM, Wareham N, Ekelund U, Cooper C; SWS Study Group. Physical activity, calcium intake and childhood bone mineral: a population-based cross-sectional study. *Osteoporosis International*. 2012;23(1):121-130.
54. Herrmann D, Buck C, Sioen I, Kouride Y, Marild S, Molnár D, et al. Impact of physical activity, sedentary behaviour and muscle strength on bone stiffness in 2-10-year-old children—cross-sectional results from the IDEFICS study. *International Journal of Behavioral Nutrition and Physical Activity*. 2015;12:112.
55. Wilson DK, Klesges LM, Klesges RC, Eck LH, Hackett-Renner CA, Alpert BS, Dalton ET. A prospective study of familial aggregation of blood pressure in young children. *Journal of Clinical Epidemiology*. 1992;45(9):959-969.
56. Butte NF, Puyau MR, Wilson TA, Liu Y, Wong WW, Adolph AL, Zakeri IF. Role of physical activity and sleep duration in growth and body composition of preschool-aged children. *Obesity*. 2016;24(6):1328-1335.
57. Collings PJ, Brage S, Ridgway CL, Harvey NC, Godfrey KM, Inskip HM, Cooper C, Wareham NJ, Ekelund U. Physical activity intensity, sedentary time, and body composition in preschoolers. *American Journal of Clinical Nutrition*. 2013;97(5):1020-1028.
58. Razak LA, Yoon SL, Wiggers J, Morgan PJ, Jones J, Finch M, Sutherland R, Lecathelnais C, Gillham K, Clinton-McHarg T, Wolfenden L. Impact of scheduling multiple outdoor free-play periods in childcare on child moderate-to-vigorous physical activity: a cluster randomised trial. *International Journal of Behavioral Nutrition and Physical Activity*. 2018;15:34.