

## PROSOCIAL BEHAVIOUR

---

# Empathy, Prosocial Behaviour and Adjustment: Clinical Aspects of Surfeits and Deficits in Concern for Others

Carolyn Zahn-Waxler, PhD, Andrew Schoen, BS

University of Wisconsin, USA

February 2016

### Introduction

Prosocial behaviours provide benefit to others. They include sharing, help, comfort, protection and defense of others, and related traits of kindness and generosity. These adaptive behaviours reflect social-emotional competence. Prosocial actions evoked by others' distress are often motivated by feelings of empathy/sympathy and a desire to alleviate their suffering.<sup>1</sup> Under some circumstances these feelings and actions can be maladaptive.<sup>2</sup> Some psychiatric disorders and psychological problems are characterized, in part, by extremes of empathy, both surfeits and deficits that undermine the capacity to care for others in a healthy manner.<sup>3,4,2</sup> The study of extremes can provide insights into processes associated with different forms of psychopathology.<sup>3,4,5,2</sup>

### Subject

Expressions of concern for others begin during the first and second year of life.<sup>6,7</sup> They are

manifest in facial and vocal expressions of empathy/sympathy, the forms of prosocial actions noted above, and cognitive awareness of the other's experience. From early on, however, three types of extremes are present<sup>8,9,10</sup> that may be precursors of later psychological and psychiatric problems.

## 1. Surfeits

High levels of empathy and prosocial behaviours are sometimes associated with anxiety and depression.<sup>11,2</sup> From early on in development extreme concern may be a sign of these internalizing problems. Extreme concern can also be seen in a genetic disorder, William's Disease; it includes mild to moderate mental disability and high sociability, where extreme prosociality can create danger.<sup>12</sup>

## 2. Deficits

- a. Callousness and hostility. This appears in the second and third years of life, after concern for others emerges and is expressed toward both adults and peers.<sup>13,7</sup> It is seen in laughter as children enjoy the person's distress, anger/aggression, and blaming the victim. It is relatively rare and may signal later conduct problems and psychopathic traits.
- b. Lack of response. This can be seen in children on the autism spectrum.<sup>3</sup> It is also seen in children high in inhibition,<sup>14</sup> which predicts later anxiety and depression in adolescence, especially in girls.<sup>15</sup>

The two types of empathy deficits are referred to as active versus passive deficits, respectively.<sup>16</sup> DSM-V psychiatric nomenclature describes disorders defined, to a significant degree, by these deficits.

## **Problem**

Surfeits and deficits in caring emotions and behaviours in children and adolescents can undermine quality of social relationships and long-term adjustment.

Others' lives also are affected (e.g., parents, siblings, peers). Research on surfeits of concern for others has been hampered by reluctance to pathologize behaviours that seem so mature and considerate. This began to change with the advent of a developmental psychopathology approach and assessment of risk factors implicating high levels of concern for others in anxiety and depression.<sup>2</sup> There is still, however, more research on active and passive deficits in concern

for others.

## **Research Context**

Because others' distress may be infrequent and unpredictable, naturalistic observations are difficult to obtain. Reports from children, adolescents, parents, teachers, and clinicians are used to assess both concern for others<sup>17,18,19</sup> and lack of regard for others.<sup>20,21</sup> Prosocial and antisocial themes also are evoked in symbolic play.<sup>22,9</sup> In early naturalistic studies<sup>23</sup> we trained mothers to make detailed, reliable, observations of children's responses to others' distress.

Structured probes (e.g., when an examiner or parent simulates pain or sorrow) are used extensively in both laboratory and home contexts. These probes first were used in studies of early normative development of concern for others<sup>24,7</sup> and then under conditions likely to evoke extreme concern, e.g. having a depressed parent.<sup>8</sup> Distress simulations also are used to study concern in older children and youth as well as those likely to have deficits, i.e. antisocial patterns and conduct problems<sup>25,26</sup> or autism spectrum children.<sup>27,3</sup> Longitudinal designs can assess whether early extremes predict later problems.<sup>25</sup>

## **Key Research Questions**

1. What environmental conditions elicit (a) surfeits of concern for others, (b) active deficits, and (c) passive deficits?
2. What biological/hormonal/genetic conditions elicit (a) surfeits of concern for others, (b) active deficits, and (c) passive deficits?
3. How do biological/gene and environmental processes interact to produce extremes? Research provides some starting points.<sup>28,29,30</sup>
4. How are surfeits and deficit in concern for others implicated in different psychological and psychiatric problems?
5. How do gender differences in extremes inform us about etiology of different forms of psychopathology?

## **Recent Research Results**

### *Surfeits of concern for others*

High-risk environments, e.g. exposure to parental depression and marital conflict<sup>31,8,9,10</sup> can evoke

higher than normative levels of concern and prosocial behaviour toward parents. Between 2 and 4 years of age, some children attempt to comfort parents in distress and mediate their conflicts. This may indicate parentification/role reversal and dissolution of boundaries, as parents' needs supersede those of their children. Children's initial empathy-based concern can fuse with anxiety and pathological guilt as children feel responsible, i.e. a cause of parental distress. Global attributions of being blameworthy or at fault are central to attributional theories of how depression develops. This may be exacerbated by depressed parent's use of guilt-induction<sup>32</sup> and other negative practices.<sup>33</sup> Subclinical and clinical anxiety and depression are present by 3 years,<sup>34,35</sup> hence early extreme concern may signal developing internalizing problems, diminished self-development, and problems with peers.<sup>36,37</sup>

More recent research confirms these findings and extends them to other populations, e.g. other forms of parental psychopathology and personality problems, alcoholism/substance abuse, early parenthood, poverty. A common theme across many studies<sup>2</sup> is that girls more often than boys are likely to show extreme concern. Other recent studies explore the multi-faceted nature of high concern in adolescents identifying (a) both costs and benefits in their friendships, peer relationships and involvement in parental conflict<sup>38,39,40</sup> and (b) high caring as a "risky strength."<sup>41</sup> Here too, girls are more affected. Possible brain and behavioural sex differences in empathy can help to explain females' susceptibilities and strengths in this domain.<sup>42</sup> Hormonal differences may be at play, as lower fetal testosterone has been linked to higher levels of empathy (though not always extremes) both in boys and girls.<sup>43</sup> In general, there has been little research on biological processes associated with surfeits of concern for others, because most researchers in these areas are unaware of potential adverse consequences.

#### *Active deficits in concern for others*

The high risk family environments identified above for surfeits in concern for others are also sometimes associated with deficits, both active and passive,<sup>44</sup> so work is needed to identify child characteristics that differentiate these three groups. Research on young children's high observed active disregard and low empathy and prosociality<sup>45,46,20</sup> predicts antisocial behaviour and psychopathic/callous-unemotional traits. Callous-unemotional traits predict severity and stability of conduct problems and delinquency.<sup>21</sup>

There is ample research on physiological correlates of active deficits and antisocial behaviour.<sup>47</sup> Measures of physiological underarousal are often associated with callous/psychopathic traits and

antisocial behaviour, though this is not invariable. Aggressive/disruptive toddlers show heightened rather than diminished physiological reactivity and they do not show lower concern for others.<sup>48</sup> Negative relations between concern for others and aggressive behaviour may develop over time, suggesting the value of early interventions since concern is still preserved in some young aggressive children.<sup>46</sup>

The salience of early development is highlighted in two recent studies of observed active disregard for others in the second and third years of life.<sup>49,25</sup> Early active disregard predicted antisocial behaviour in childhood and adolescence based on mother, teacher and child reports. Early language predicted less disregard and greater concern, suggesting the possible protective role and the importance of encouraging language from the first years of life. There is also substantial research on environmental contributions, including child-rearing and discipline practices, to active disregard and antisocial behaviour.<sup>50,51,46</sup>

Atypical empathy is present at the neural level in adolescents with conduct disorder and psychopathic traits.<sup>16</sup> Youths appeared to show no neural response deficits in pain-experiencing regions when viewing others in distress. However, those with conduct disorder showed less coupling compared to controls between the amygdala, a key region in emotion processing, and the ventromedial prefrontal cortex, a region thought to be involved in behavioural responses. This relative deficit in functional connectivity between these regions has been found for individuals with callous-disregard as well.<sup>52</sup> Another study with adolescents with conduct disorder also found structural neural deficits associated with lack of empathy.<sup>53</sup>

#### *Passive deficits in concern for others*

Laboratory research using structured distress probes documents deficits in empathy and prosocial behaviour in children on the autism spectrum<sup>54,55</sup> consistent with parent reports. It is not clear why these differences occur and whether they always reflect core deficits; greater emotional reactivity and sensitivity to environmental stressors as indexed by high levels of cortisol,<sup>56</sup> and lack of communicative skills associated with neurological deficits may blunt empathy in some children. Since language plays a role in empathy even in the first years of life,<sup>54,49</sup> the study of variations in language in autistic children may help to explain why empathy is relatively preserved in some of these children.<sup>27</sup> When autism was first identified as a disorder, cold, distant mothering, (a.k.a. 'refrigerator mothers') was claimed to create autism, including empathy deficits. These views were discredited, as the primacy of biological/genetic

underpinnings became known.

In humans, exposure to high levels of prenatal androgens may result in masculine behaviours and abilities. Simon Baron-Cohen has proposed an extreme male brain of autism whereby fetal testosterone, more common in males than females, creates a hyper-masculinized brain, associated with autism/Asperger's, difficulty in social relationships, and restricted interests.<sup>57</sup> This may also be true at a subsyndromal level. In typically developing 4 year-olds, fetal testosterone predicted problems in empathy, social relations and restricted interests, for both sexes.<sup>58</sup> Similar patterns were observed in other research, with fetal testosterone, showing an inverse relationship with empathy.<sup>59,43</sup> More male-typical behaviours and fewer female-typical behaviours, including empathy<sup>60</sup> are seen in females exposed to high prenatal testosterone due to a genetic disorder congenital adrenal hyperplasia (CAH) or because mothers were prescribed hormones during pregnancy.

Passive deficits occur on a continuum; low concern does not necessarily reflect psychopathology but can still create interpersonal problems. Physiological and gene re-lated effects have been identified. Low empathy in preschool children of depressed mothers is associated with right frontal EEG asymmetry.<sup>61</sup> The AVPR1A gene variant is associated with preschooler's lower altruistic behaviour.<sup>62</sup>

## **Research Gaps**

There are no well-established standardized tests or norms for identifying surfeits and deficits in concern for others. Mostly, extremes are inferred based on how they relate to or predict other measures that reflect risk and/or psychopathology. Often, extremes result from a combination of genetic and environmental factors, yet little is known about specific processes that interact to produce different developmental outcomes. Only some children show surfeits or deficits even in high-risk environments and some children show surfeits or deficits in apparently low-risk environments. Future research is needed to address these complexities. Also, rather than just dichotomizing children as extreme or not, it is important to study individual differences within categories of surfeits and deficits.

Little is known about intentions and motives that underlie surfeits and deficits in concern for others. Initial empathy-based acts of caring toward distressed caregivers may be taken over by anxiety, guilt and shame. Greater knowledge of children's emotions is needed. Some children,

who appear to be inexpressive, may in fact experience concern that we do not yet know how to tap. Some show multiple emotions associated with both concern and active disregard in the same context. What sets these children apart?

## **Conclusions**

Three extremes of empathy and prosociality, i.e. surfeits, active deficits, and passive deficits emerge in the first years of life. These extremes have been associated with different psychological and psychiatric problems later in development. Surfeits are more commonly associated with internalizing problems and deficits with externalizing problems and autism spectrum disorders. Comorbidity is also possible and requires further attention. Surfeits and deficits in empathy and prosociality are not invariably prodromal signs of later problems; hence it will be important to determine why only some young children go on to experience serious difficulties.

Knowledge about surfeits and deficits of concern for others has come mainly from three largely separate research domains. Conceptual and empirical work would benefit from studies that explore relations among them, e.g. recent work comparing multiple features of empathy in two different populations (autism spectrum and conduct disorders) who both show deficits<sup>63</sup> or examining concern and active disregard in the same populations.<sup>46</sup>

Both normatively<sup>64</sup> and at the extremes, girls show higher empathy and prosocial behaviour than boys and boys show more active and passive disregard than girls. This parallels sex differences in forms of psychopathology from childhood and adolescence through adulthood. Conduct problems and autism-spectrum problems show a marked male preponderance, while anxiety and depression show a marked female preponderance.<sup>65</sup> Empathy deficits in fact are symptoms that help to define male-preponderant problems and surfeits are correlates (possibly symptoms or causes) of female preponderant problems. Gender differences in concern and disregard, in conjunction with other known gender differences in child temperament,<sup>66</sup> may provide a better window into our understanding of etiologies of the different psychological and psychiatric problems considered here.<sup>67</sup>

## **Implications for Parents, Services, and Policy**

It is valuable for parents, teachers, and other caregivers to encourage children's social competence, including expressions of concern for others, and to begin early in life. Several

programs are available,<sup>68-73</sup> more often for older children than younger children, and there is considerable research to guide additional program development.<sup>74-79,32</sup> More work has been done with community samples than with troubled children. The extent to which intervention paradigms and findings from community samples will generalize to extremes in concern and lack of regard for others is not yet clear.

For children with surfeits of concern for others, interventions exist to improve social functioning by reducing children's sense of responsibility and empathic over-involvement for the problems of their parents.<sup>80,81</sup> Because parental distress is also associated with other extremes of aggression and avoidance,<sup>44</sup> i.e. deficits in concern for others, further interventions should be tailored to these child characteristics.

Recent classroom interventions with preschoolers and older children have focused on mindfulness and loving-kindness practices to increase attentional focus and self-regulation, heighten empathy, and reduce bullying and other forms of aggression.<sup>82,83</sup> Practices to increase mindfulness are now used with parents,<sup>84</sup> but not yet with children at the extremes. Such practices might help reduce both overly high and low concern for others, since one goal is to subdue overwhelming and stormy feelings, as well as create calm and caring for the self. While we've emphasized the need for environmental interventions, recent work on biological interventions, is also relevant to empathy.<sup>85,75,28</sup> Oxytocin, for example, plays a role in mediating low parental mood and child empathy.

Some extremes in concern and disregard for others and associated internalizing and externalizing problems are unlikely to be amenable to interventions, because they occur within the broader context of societal problems such as poverty and parental problems such as child maltreatment. Interventions directed solely toward the child may be of little consequence until the larger issues are addressed.

## References

1. Eisenberg N, Fabes RA, Spinrad TL. Prosocial development. In: Eisenberg N, Damon W, Lerner RM, eds. *Handbook of child psychology*. Hoboken, NJ: Wiley; 2006:646-718.
2. Zahn-Waxler C, Van Hulle C. Empathy, guilt and depression: When caring for others becomes costly to children. In: Oakley B, Knafo A, Madhavan G, Wilson DS, eds. *Pathological altruism*. New York, NY US: Oxford University Press; 2011:243-259.
3. Sigman MD, Kasari C, Kwon JH, Yirmiya N. Responses to the negative emotions of others by autistic, mentally retarded, and normal children. *Child Development*. 1992;63(4):796-807. doi: 10.2307/1131234.
4. Blair RJR. Responding to the emotions of others: dissociating forms of empathy through the study of typical and psychiatric populations. *Consciousness and Cognition*. 2005;14(4):698-718. doi: 10.1016/j.concog.2005.06.004.



5. Decety J, Moriguchi Y. The empathic brain and its dysfunction in psychiatric populations: implications for intervention across different clinical conditions. *BioPsychoSocial Medicine*. 2007;1(1):22. doi: 10.1186/1751-0759-1-22.
6. Davidov M, Zahn-Waxler C, Roth-Hanania R, Knafo A. Concern for others in the first year of life: theory, evidence, and avenues for research. *Child Development Perspectives*. 2013;7(2):126-131. doi: 10.1111/cdep.12028.
7. Zahn-Waxler C, Radke-Yarrow M, Wagner E, Chapman M. Development of concern for others. *Developmental Psychology*. 1992;28(1):126. doi: 10.1037/0012-1649.28.1.126.
8. Radke-Yarrow M, Zahn-Waxler C, Richardson DT, Susman A, Martinez P. Caring behavior in children of clinically depressed and well mothers. *Child Development*. 1994;65(5):1405-1414. doi: 10.1111/j.1467-8624.1994.tb00825.x
9. Zahn-Waxler C, Kochanska G, Krupnick J, McKnew D. Patterns of guilt in children of depressed and well mothers. *Developmental Psychology*. 1990;26(1):51. doi: 10.1037/0012-1649.26.1.51.
10. Cummings EM, Zahn-Waxler C, Radke-Yarrow M. Young children's responses to expressions of anger and affection by others in the family. *Child Development*. 1981;52(4):1274-1282. doi: 10.2307/1129516
11. O'Connor LE., Berry JW, Lewis TB, Stiver DJ. Empathy-based pathogenic guilt, pathological altruism, and psychopathology. In: Oakley B, Knafo A, Madhavan G, Wilson DS, eds. *Pathological altruism*. New York, NY US: Oxford University Press; 2011:10-30.
12. Mervis CB, Klein-Tasman BP. Williams syndrome: cognition, personality, and adaptive behavior. *Mental Retardation and Developmental Disabilities Research Reviews*. 2000;6(2):148-158. doi: 10.1002/1098-2779(2000)6:2<148::AID-MRDD10>3.0.CO;2-T.
13. Klimes-Dougan B, Kistner J. Physically abused preschoolers' responses to peers' distress. *Developmental Psychology*. 1990;26(4):599. doi: 10.1037/0012-1649.26.4.599.
14. Young S, Fox N, Zahn-Waxler C. Relations between temperament and empathy in two-year-olds. *Developmental Psychology*. 1999;35(5):1189-1197. doi: 10.1037/0012-1649.35.5.1189.
15. Schwartz CE, Snidman N, Kagan J. Adolescent social anxiety as an outcome of inhibited temperament in childhood. *Journal of the American Academy of Child & Adolescent Psychiatry*. 1999;38(8):1008-1015. doi: 10.1097/00004583-199908000-00017.
16. Decety J, Michalska KJ, Akitsuki Y, Lahey BB. Atypical empathic responses in adolescents with aggressive conduct disorder: a functional MRI investigation. *Biological Psychology*. 2009;80(2):203-211. doi: 10.1016/j.biopsycho.2008.09.004.
17. Kochanska G. *My Child, version 2: A preliminary manual*. Iowa City: University of Iowa, Department of Psychology; 1992.
18. Bryant BK. An index of empathy for children and adolescents. *Child development*. 1982;53(2):413-425. doi: 10.2307/1128984.
19. Davies PT, Coe JL, Martin MJ, Sturge-Apple ML, Cummings EM. The developmental costs and benefits of children's involvement in interparental conflict. *Developmental Psychology*. 2015;51(8):1026-1047. doi: 10.1037/dev0000024.
20. Shirtcliff EA, Vitacco MJ, Graf AR, Gostisha AJ, Merz JL, Zahn-Waxler C. Neurobiology of empathy and callousness: implications for the development of antisocial behavior. *Behavioral Sciences & the Law*. 2009;27(2):137-171. doi: 10.1002/bsl.862.
21. Frick PJ, Stickle TR, Dandreaux DM, Farrell JM, Kimonis ER. Callous-unemotional traits in predicting the severity and stability of conduct problems and delinquency. *Journal of Abnormal Child Psychology*. 2005;33(4):471-487. doi: 10.1007/s10648-005-5728-9
22. Zahn-Waxler C, Park JH, Usher B, Belouad F, Cole P, Gruber R. Young children's representations of conflict and distress: A longitudinal study of boys and girls with disruptive behavior problems. *Development and Psychopathology*. 2008;20(01):99-119. doi: 10.1017/S0954579408000059

23. Zahn-Waxler C, Radke-Yarrow M, King RA. Child rearing and children's prosocial initiations toward victims of distress. *Child Development*. 1979;50(2):319-330. doi: 10.2307/1129406.
24. Zahn-Waxler C, Robinson JL, Emde RN. The development of empathy in twins. *Developmental Psychology*. 1992;28(6):1038. doi: 10.1037/0012-1649.28.6.1038
25. Rhee SH, Friedman NP, Boeldt DL, Corley RP, Hewitt JK, Knafo A, Lahey BB, Robinson J, Van Hulle CA, Waldman ID, Young SE, Zahn-Waxler C. Early concern and disregard for others as predictors of antisocial behavior. *Journal of Child Psychology and Psychiatry, and Allied Disciplines*. 2013;54(2):157-166. doi:10.1111/j.1469-7610.2012.02574.x.
26. Van Hulle C, Zahn-Waxler C, Robinson JL, Rhee SH, Hastings PD, Knafo A. Autonomic correlates of children's concern and disregard for others. *Social Neuroscience*. 2013;8(4):275-290. doi: 10.1080/17470919.2013.791342.
27. Scheeren AM, Koot HM, Mundy PC, Mous L, Begeer S. Empathic responsiveness of children and adolescents with high-functioning autism spectrum disorder. *Autism Research*. 2013;6(5):362-371. doi: 10.1002/aur.1299.
28. Bakermans-Kranenburg MJ, van Ijzendoorn MH. Oxytocin receptor (OXTR) and serotonin transporter (5-HTT) genes associated with observed parenting. *Social Cognitive and Affective Neuroscience*. 2008;3(2):128-134. doi: 10.1093/scan/nsn004.
29. Harold GT, Rice F, Hay DF, Boivin J, van den Bree M, Thapar A. Familial transmission of depression and antisocial behavior symptoms: disentangling the contribution of inherited and environmental factors and testing the mediating role of parenting. *Psychological Medicine*. 2011;41(06):1175-1185. doi: 10.1017/S0033291710001753.
30. Knafo A, Israel S, Ebstein RP. Heritability of children's prosocial behavior and differential susceptibility to parenting by variation in the dopamine receptor D4 gene. *Development and Psychopathology*. 2011;23(01):53-67. doi: 10.1017/S0954579410000647
31. Klimes-Dougan B, Bolger A. Coping with maternal depressed affect and depression: Adolescent children of well and depressed mothers. *Journal of Youth and Adolescence*. 1998;27:1-15.
32. Zahn-Waxler C, Iannotti RJ, Cummings EM, Denham S. Antecedents of problem behaviors in children of depressed mothers. *Development and Psychopathology*. 1990;2(03):271-291. doi: 10.1017/S0954579400000778.
33. Goodman SH, Rouse MH, Connell AM, Broth MR, Hall CM, Heyward . Maternal depression and child psychopathology: A meta-analytic review. *Clinical Child and Family Psychology Review*. 2011;14(1):1-27. doi: 10.1007/s10567-010-0080-1.
34. Luby J, Belden A, Sullivan J, Hayen R, McCadney A, Spitznagel E. Shame and guilt in preschool depression: evidence for elevations in self-conscious emotions in depression as early as age 3. *Journal of Child Psychology and Psychiatry* . 2009;50(9):1156-1166. doi: 10.1111/j.1469-7610.2009.02077.x.
35. Carter AS, Briggs-Gowan MJ, Jones SM, Little TD. The infant-toddler social and emotional assessment (ITSEA): Factor structure, reliability, and validity. *Journal of Abnormal Child Psychology*. 2003;31(5):495-514. doi: 10.1023/A:1025449031360.
36. Denham SA, Zahn-Waxler C, Cummings EM, Iannotti RJ. Social competence in young children's peer relations: Patterns of development and change. *Child Psychiatry and Human Development*. 1991;22(1):29-44. doi: 10.1007/BF00706057.
37. Essex MJ, Kraemer HC, Armstrong JM, Boyce WT, Goldsmith HH, Klein MH, Woodward H, Kupfer DJ. Exploring risk factors for the emergence of children's mental health problems. *Archives of General Psychiatry*. 2006;63(11):1246-1256. doi: 10.1001/archpsyc.63.11.1246.
38. Smith RL, Rose AJ. The "cost of caring" in youths' friendships: Considering associations among social perspective taking, co-rumination, and empathetic distress. *Developmental Psychology*. 2011;47(6):1792. doi: 10.1037/a0025309.
39. Rudolph KD, Conley CS. The socioemotional costs and benefits of social-evaluative concerns: Do girls care too much? *Journal of Personality*. 2005;73(1):115-138. doi: 10.1111/j.1467-6494.2004.00306.x.

40. Davies PT, Coe JL, Martin MJ, Sturge-Apple ML, Cummings EM. The developmental costs and benefits of children's involvement in interparental conflict. *Developmental Psychology*. 2015;51(8):1026-1047. doi: 10.1037/dev0000024.
41. Tone EB, Tully EC. Empathy as a "risky strength": A multilevel examination of empathy and risk for internalizing disorders. *Development and Psychopathology*. 2014;26(4pt2):1547-1565. doi: 10.1017/S0954579414001199.
42. Christov-Moore L, Simpson EA, Coudé G, Grigaityte K, Iacoboni M, Ferrari PF. Empathy: Gender effects in brain and behavior. *Neuroscience & Biobehavioral Reviews*. 2014;46:604-627. doi: 10.1016/j.neubiorev.2014.09.001.
43. Chapman E, Baron-Cohen S, Auyeung B, Knickmeyer R, Taylor K, Hackett G. Fetal testosterone and empathy: Evidence from the empathy quotient (EQ) and the "reading the mind in the eyes" test. *Social Neuroscience*. 2006;1(2):135-148. doi: 10.1080/17470910600992239.
44. Solantaus-Simula T, Punamäki RL, Beardslee WR. Children's responses to low parental mood. I: Balancing between active empathy, overinvolvement, indifference, and avoidance. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2002;41(3):278-286. doi: 10.1097/00004583-200203000-00007.
45. Blair RJR. Responsiveness to distress cues in the child with psychopathic tendencies. *Personality and Individual Differences*. 1999;27(1):135-145. doi: 10.1016/S0191-8869(98)00231-1.
46. Hastings PD, Zahn-Waxler C, Robinson J, Usher B, Bridges D. The development of concern for others in children with behavior problems. *Developmental Psychology*. 2000;36(5):531. doi: 10.1037//0012-1649.36.5.531.
47. Ortiz J, Raine A. Heart rate level and antisocial behavior in children and adolescents: A meta-analysis. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2004;43(2):154-162. doi: 10.1097/00004583-200402000-00010.
48. Gill KL, Calkins SD. Do aggressive/destructive toddlers lack concern for others? Behavioral and physiological indicators of empathic responding in 2-year-old children. *Development and Psychopathology*. 2003;15(01):55-71. doi: 10.1017.S095457940300004X.
49. Rhee SH, Boeldt DL, Friedman NP, Corley RP, Hewitt JK, Young SE, Knafo A, Robinson J, Waldman ID, Van Hulle CA, Zahn-Waxler C. The role of language in concern and disregard for others in the first years of life. *Developmental Psychology*. 2013;49(2):197. doi: 10.1037/a0028318.
50. Waller R, Gardner F, Hyde LW. What are the associations between parenting, callous-unemotional traits, and antisocial behavior in youth? A systematic review of evidence. *Clinical Psychology Review*. 2013;33(4):593-608. doi: 10.1016/j.cpr.2013.03.001.
51. Wagner NJ, Mills-Koonce WR, Willoughby MT, Zvara B, Cox MJ. Parenting and children's representations of family predict disruptive and callous-unemotional behaviors. *Developmental Psychology*. 2015;51(7):935-948. doi: 10.1037/a0039353.
52. Marsh AA, Finger EC, Mitchell DG, Reid ME, Sims C, Kosson DS, Towbin KE, Leibenluft E, Pine DS, Blair RJ. Reduced amygdala response to fearful expressions in children and adolescents with callous-unemotional traits and disruptive behavior disorders. *American Journal of Psychiatry*. 2008;165(6):712-720. doi: 10.1176/appi.ajp.2007.07071145.
53. Sterzer P, Stadler C, Poustka F, Kleinschmidt A. A structural neural deficit in adolescents with conduct disorder and its association with lack of empathy. *Neuroimage*. 2007;37(1):335-342. doi: 10.1016/j.neuroimage.2007.04.043.
54. Hutman T, Rozga A, DeLaurentis A, Sigman M, Dapretto M. Infants' pre-empathic behaviors are associated with a skills. *Infant Behavior and Development*. 2012;35(3):561-569. doi: 10.1016/j.infbeh.2012.05.007.
55. Hobson JA, Harris R, García-Pérez R, Hobson RP. Anticipatory concern: A study in autism. *Developmental Science*. 2009;12(2):249-263. doi: 10.1111/j.1467-7687.2008.00762.x.
56. Putnam SK, Lopata C, Thomeer ML, Volker MA, Rodgers JD. Salivary Cortisol Levels and Diurnal Patterns in Children with Autism Spectrum Disorder. *Journal of Developmental and Physical Disabilities*. 2015;27(4):453-465. doi: 10.1007/s10882-015-9428-2.

57. Baron-Cohen S. The extreme male brain theory of autism. *Trends in cognitive sciences*. 2002;6(6):248-254. doi:10.1016/S1364-6613(02)01904-6.
58. Knickmeyer R, Baron-Cohen S, Raggatt P, Taylor, K. Foetal testosterone, social relationships, and restricted interests in children. *Journal of Child Psychology and Psychiatry*. 2005;46(2):198-210. doi: 10.1111/j.1469-7610.2004.00349.x.
59. Knickmeyer R, Baron-Cohen S, Raggatt P, Taylor K, Hackett G. Fetal testosterone and empathy. *Hormones and Behavior* . 2006;49(3):282-292. doi: 10.1016/j.yhbeh.2005.08.010.
60. Constantinescu M, Hines M. Relating prenatal testosterone exposure to postnatal behavior in typically developing children: Methods and findings. *Child Development Perspectives*. 2012;6(4):407-413. doi: 10.1111/j.1750-8606.2012.00257.x
61. Jones NA, Field T, Davalos M. Right frontal EEG asymmetry and lack of empathy in preschool children of depressed mothers. *Child Psychiatry and Human Development*. 2000;30(3):189-204. doi: 10.1023/A:1021399605526.
62. Avinun R, Israel S, Shalev I, Gritsenko I, Bornstein G, Ebstein RP, Knafo A. AVPR1A variant associated with preschoolers' lower altruistic behavior. *PLoS One*. 2011;6(9):e25274-e25274. doi: 10.1371/journal.pone.0025274.
63. Bons D, van den Broek E, Scheepers F, Herpers P, Rommelse N, Buitelaar JK. Motor, emotional, and cognitive empathy in children and adolescents with autism spectrum disorder and conduct disorder. *Journal of Abnormal Child Psychology*. 2013;41(3):425-443. doi: 10.1007/s10802-012-9689-5.
64. McClure EB. A meta-analytic review of sex differences in facial expression processing and their development in infants, children, and adolescents. *Psychological Bulletin*. 2000;126(3):424. doi: 10.1037/0033-2909.126.3.424.
65. Zahn-Waxler C, Shirtcliff EA, Marceau K. Disorders of childhood and adolescence: Gender and psychopathology. *Annual Review of Clinical Psychology*. 2008;4:275-303. doi: 10.1146/annurev.clinpsy.3.022806.0913.
66. Else-Quest NM, Hyde JS, Goldsmith HH, Van Hulle CA. Gender differences in temperament: a meta-analysis. *Psychological Bulletin*. 2006;132(1):33. doi: 10.1037/0033-2909.132.1.33.
67. Rutter M, Caspi A, Moffitt TE. Using sex differences in psychopathology to study causal mechanisms: unifying issues and research strategies. *Journal of Child Psychology and Psychiatry*. 2003;44(8):1092-1115. doi: 10.1111/1469-7610.00194.
68. Havighurst SS, Wilson KR, Harley AE, Prior MR, Kehoe C. Tuning in to kids: Improving emotion socialization practices in parents of preschool children—findings from a community trial. *Journal of Child Psychology and Psychiatry* . 2010;51(12):1342-1350. doi: 10.1111/j.1469-7610.2010.02303.x.
69. Dewar G. Teaching empathy: Evidence-based tips for fostering empathy in children. *Parenting Science* . 2009. Available at: <http://www.parentingscience.com/teaching-empathy-tips.html>. Accessed January 29, 2016.
70. Castillo R, Salguero JM, Fernández-Berrocal P, Balluerka N. Effects of an emotional intelligence intervention on aggression and empathy among adolescents. *Journal of Adolescence*. 2013;36(5):883-892. doi: 10.1016/j.adolescence.2013.07.001.
71. Hyde LW, Shaw DS, Gardner F, Cheong J, Dishion TJ, Wilson M. Dimensions of callousness in early childhood: Links to problem behavior and family intervention effectiveness. *Development and Psychopathology*. 2013;25(02):347-363. doi: 10.1017/S0954579412001101.
72. Wilson KR, Havighurst SS, Harley AE. Tuning in to kids: An effectiveness trial of a parenting program targeting emotion socialization of preschoolers. *Journal of Family Psychology*. 2012;26(1):56. doi: 10.1037/a0026480.
73. Schonert-Reichl KA, Smith V, Zaidman-Zait A, Hertzman C. Promoting children's prosocial behaviors in school: Impact of the "Roots of Empathy" program on the social and emotional competence of school-aged children. *School Mental Health* . 2012;4(1):1-21. doi: 10.1007/s12310-011-9064-7.
74. Vinik J, Almas A, Grusec J. Mothers' knowledge of what distresses and what comforts their children predicts children's coping, empathy, and prosocial behavior. *Parenting: Science and Practice*. 2011;11(1):56-71. doi: 10.1080/15295192.2011.539508.

75. Apter-Levy Y, Feldman M, Vakart A, Ebstein RP, Feldman R. Impact of maternal depression across the first 6 years of life on the child's mental health, social engagement, and empathy: The moderating role of oxytocin. *American Journal of Psychiatry*. 2014;170(10):1161-1168. doi: 10.1176/appi.ajp.2013.12121597.
76. Brownell CA, Svetlova M, Anderson R, Nichols SR, Drummond J. Socialization of early prosocial behavior: Parents' talk about emotions is associated with sharing and helping in toddlers. *Infancy*. 2013;18(1):91-119. doi: 10.1111/j.1532-7078.2012.00125.x.
77. Tong L, Shinohara R, Sugisawa Y, Tanaka E, Yato Y, Yamakawa N, Anme T. Early development of empathy in toddlers: Effects of daily parent-child interaction and home-rearing environment. *Journal of Applied Social Psychology*. 2012;42(10):2457-2478. doi: 10.1111/j.1559-1816.2012.00949.x.
78. Hastings PD, Utendale WT, Sullivan C. The socialization of prosocial development. In: Grusec JE, Hastings PD, eds. *Handbook of Socialization: Theory and Research*. New York and London: The Guilford Press; 2007:638-664.
79. Willoughby MT, Mills-Koonce WR, Gottfredson NC, Wagner NJ. Measuring callous unemotional behaviors in early childhood: factor structure and the prediction of stable aggression in middle childhood. *Journal of Psychopathology and Behavioral Assessment*. 2014;36(1):30-42. doi: 10.1007/s10862-013-9379-9.
80. Beardslee WR, Gladstone TR. Prevention of childhood depression: Recent findings and future prospects. *Biological Psychiatry*. 2001;49(12):1101-1110. doi: 10.1016/S0006-3223(01)01126-X.
81. Beardslee WR, Gladstone TR, Wright EJ, Cooper AB. A family-based approach to the prevention of depressive symptoms in children at risk: Evidence of parental and child change. *Pediatrics*. 2003;112(2):e119-e131. doi: 10.1542/peds.112.2.e119.
82. Flook L, Goldberg SB, Pinger L, Davidson RJ. Promoting prosocial behavior and self-regulatory skills in preschool children through a mindfulness-based kindness curriculum. *Developmental Psychology*. 2015;51(1):44-51. doi: 10.1037/a0038256.
83. Schonert-Reichl KA, Oberle E, Lawlor MS, Abbott D, Thomson K, Oberlander TF, Diamond A. Enhancing cognitive and social-emotional development through a simple-to-administer mindfulness-based school program for elementary school children: A randomized controlled trial. *Developmental Psychology*. 2015;51(1):52. doi: 10.1037/a0038454.
84. Coatsworth JD, Duncan LG, Nix RL, Greenberg MT, Gayles JG, Bamberger KT, Berrena E, Demi MA. Integrating mindfulness with parent training: Effects of the mindfulness-enhanced strengthening families program. *Developmental Psychology*. 2015;51(1):26-35. doi: 10.1037/a0038212.
85. Harris JC, Carter CS. Therapeutic interventions with oxytocin: current status and concerns. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2013;52(10):998-1000. doi: 10.1016/j.jaac.2013.08.001.