

TOBACCO AND PREGNANCY

Prenatal Cigarette Smoke Exposure: Effects on Offspring

Marie D. Cornelius, PhD, Natacha De Genna, PhD

University of Pittsburgh School of Medicine, USA

March 2011

Introduction

Over 20% of North American and Canadian adults smoke despite the fact that cigarette smoking is the leading cause of preventable morbidity and mortality.^{1,2} Tobacco is also the most commonly used substance during pregnancy;³ 13.8% of pregnant American women smoke⁴ and 10.5% of pregnant Canadian women smoke⁵ affecting more than 600,000 live births per year.

Subject and Research Context

The scientific literature is replete with findings of adverse prenatal cigarette smoke exposure (PCSE) effects on offspring. More recent studies have improved on earlier methodologies with prospective designs, biological measures and statistical control of confounders. Many of these studies also use more sophisticated measures including objective *neurobehavioural test batteries* , *neuroimaging techniques* and assessment of *genotypic variability*. This brief synopsis will focus on studies published since the previous report on this topic by Peter Fried.⁶

Recent Research Results

The evidence is clear about PCSE's causal effects on neonatal morbidity and mortality,⁷ and its

effects on neurobehaviour is a recent focus in the infancy literature. PCSE is associated with infant irritability, inattention, decreased response to inanimate auditory stimuli,⁸ *increased hypertonicity*⁹ and more problematic temperament.¹⁰ Key et al.¹¹ found that PCSE infants discriminated fewer syllables and processed them more slowly than non-exposed infants. Similarly, Golub and colleagues¹² demonstrated that exposed nonhuman primates showed less novelty preference in visual recognition.

Studies on the effects during childhood continue to show that PCSE is a consistent predictor of higher rates of problems such as language delay in preschoolers,¹³ externalizing and internalizing behavioural problems such as acting out or withdrawal in 2-year-old children,¹⁴ child aggression in 17- to 42-month olds,¹⁵ and externalizing in early childhood that persists to age 18.¹⁶ Studies have also considered the effects of both prenatal and postnatal tobacco smoke exposure and have demonstrated an independent effect of PCSE on childhood behaviour problems at ages 6¹⁷ and 10.^{18,19} In a large multinational study, Brion et al.² reported a direct relation between PCSE and offspring conduct and externalizing problems. In a clinical sample of children with ADHD, hyperactive-impulsive symptoms and conduct disorder symptoms were significantly higher among those with PCSE.²¹ Others have found a relation between PCSE and conduct disorder symptoms.^{22,23} Murray and colleagues²³ found that this relation between PCSE and conduct disorder symptoms in childhood extended to criminal behaviours when the offspring were ages 30-34. PCSE also has long-term effects on physical health in offspring. Johansson et al.²⁴ found timing effects in offspring wheezing, sleeping difficulties, excessive crying, and use of bronchodilating drugs depending on whether exposure was prenatal only, postnatal only, or both. Gilman et al.²⁵ examined over 52,000 children from birth to age 7 and found that low birth weight and a higher odds for being overweight were associated with PCSE. In two meta-analyses,^{26,27} PCSE was significantly related to offspring obesity and overweight, respectively. Rooney et al.²⁸ also found that PCSE was a significant predictor of obesity in a birth cohort during adolescence and adulthood.

Recent studies on the effects of PCSE on risk for smoking in offspring have replicated and extended earlier reports. Menezes et al.²⁹ found this relation in early adolescence. Agrawal and colleagues³⁰ demonstrated that PCSE was associated with earlier age of initiation as well as earlier regular smoking. O'Callaghan and colleagues³¹ reported this relation with nicotine dependence in young adulthood. Lotfipour and colleagues³² reported an interactive effect between in utero tobacco smoke exposure *polymorphism* in the *nicotinic acetylcholine* receptor

that influences smoking and other drug use. Underscoring these findings in an animal model, Slotkin et al.³³ noted that both prenatal and adolescent nicotine exposures resulted in permanent changes in synaptic function, and prenatal exposure sensitized females to the subsequent effects of nicotine.

With the development of faster genetic mapping techniques, more studies have incorporated genetics into their designs, documenting interaction effects of PCSE and genetic liability on outcomes. One such design found a dampened response to novelty among PCSE infants with genetic variability in the *DRD2* gene.³⁴ In another study,³⁵ 15-year-old males with PCSE who were *homozygous* for the DAT1 dopamine transporter gene had higher rates of hyperactivity and impulsivity than all other groups.

Similarly, Neuman³⁶ found that the odds of a *DSM-IV* Attention Deficit Hyperactivity Disorder (ADHD) diagnosis were 2.9 times greater in twins with the *DAT1 allele*, 2.6 times greater in those with the *DRD4* seven-repeat allele, and 9.0 for offspring with PCSE and both alleles. Wakschlag and colleagues³⁷ tested the effects of a polymorphism of the enzyme *monamine oxidase* (MAOA) and PCSE on antisocial behaviour in adolescents demonstrating that *MAOA genotype*, PCSE and gender interact to predict antisocial behaviour in exposed male offspring.

Brain imaging studies have also advanced our knowledge about the mechanisms that may drive PCSE effects. Rivkin³⁸ examined the brain volume in a *magnetic resonance imaging (MRI)* study of 10- to 14-year-olds. PCSE was associated with significant reduction in cortical gray matter and total *parenchymal* volumes and head circumference. In another MRI study, Toro et al.³⁹ found that orbitofrontal, middle frontal, and *parahippocampal* cortices were thinner in the PCSE- exposed adolescents. Jacobsen et al.⁴⁰ used MRI and *diffusion tensor imaging (DTI)* to examine effects of prenatal and adolescent tobacco exposure on structure of brain white matter and found that both white matter microstructure and auditory processing were affected indicating that nicotine-induced disruption of the auditory *corticofugal fibers* may lead to reduced efficiency in auditory processing. In another study by Jacobsen and colleagues,⁴¹ PCSE and exposure during adolescence was found to exert gender-specific deleterious effects on auditory and visual attention *fMRI (functional magnetic resonance imaging)*.

Research Gaps

Despite the mounting body of evidence linking PCSE to problems for offspring, several recent

studies have not found significant associations.^{42,43,44,45,46} These results suggest that it is important to consider factors that may account for the explained variance in outcomes, many of which are also related to smoking during pregnancy, such as lower education and income. Several review articles have synthesized PCSE effects,^{47,48,49} with some reviews including non-human data.^{50,51,52} One recent review⁵³ of the long-term consequences of fetal and neonatal nicotine exposure highlights concerns about the safety and utility of nicotine replacement therapy during pregnancy. Suggestions from reviews to address current gaps in the literature include: considering multiple factors in explaining the nature of attention hyperactivity disorder and behavioural problems; attaining an unbiased estimate of the magnitude of the association between exposure and outcome, and more comprehensive study designs that involve the gene-environment interplay.

Conclusions

The large majority of recent studies have built upon the evidence of the last five decades that smoking during pregnancy is deleterious for multiple offspring outcomes and that these effects can be detected from infancy through adulthood. However, based on some of the studies with negative findings, it is apparent that there is a need for adequate control of potential confounds that may also contribute to these outcomes. Establishing causal links requires replication of findings across large numbers of studies with varying study populations. Part of the difficulty in concluding causal effects is due to the inability to separate prenatal exposure effects from other confounding environmental and genetic factors. New research examining genetic liability suggests that PCSE may interact with genes to produce effects. Animal models offer added support for linkage by supporting biological plausibility for such relations. Imaging techniques provide visible effects of exposure on brain structure and function. Although the mechanisms of PCSE's effects on the developing brain are not completely understood, newer research informs us that the mechanisms are multifactorial, involving biological effects, genetic susceptibility, and environmental factors.

Implications for Parents, Services, Policy

Since 1957 a plethora of studies have implicated PCSE with multiple adverse outcomes in offspring from birth through adulthood.⁵⁴ It is encouraging that rates of smoking during pregnancy have decreased in the past decade, but the prevalence is still too high, especially for specific groups that are already at higher risk for fetal problems such as teenagers and women of low-

economic status. Public health efforts that have proved to be efficacious in reducing smoking should continue to receive support. With evidence of nicotine vulnerability among those prenatally exposed, stopping exposure in the first place, will not only prevent neurotoxicological effects in the offspring, but will prevent continuing smoking exposure in succeeding generations. There is now notable evidence that preventing PCSE may help prevent smoking uptake during adolescence in offspring. Based on what we do know, it is clear that women who are pregnant or who may become pregnant should abstain from smoking and nicotine exposure.

References

1. Centers for Disease Control and Prevention. Vital Signs: Current cigarette smoking among adults aged ≥ 18 years - United States, 2009. *MMWR*. 2010;59(35):1135-1140.
2. Statistics Canada. Smoking, 2009: Canadian Community Health Survey. Health Fact Sheets, Aug 16, 2010.
3. Centers for Disease Control and Prevention (CDC). Smoking during pregnancy - United States, 1990-2002. *MMWR*. 2004;53,911-915.
4. Tong V, Jones J, Dietz P, D'Angelo D, Bombard J. Trends in smoking before, during, and after pregnancy-Pregnancy risk assessment monitoring system (PRAMS), United States, 31 Sites, 2000-2005. *MMWR Surveill Summ*. 2009;58(SS04):1-29.
5. Al Sa-hab B, Saqib M, Hauser G, Tamim H. Prevalence of smoking during pregnancy and associated risk factors among Canadian women: a national survey. *BMC Pregnancy and Childbirth*. 2010;10:1-9.
6. Fried P. Tobacco consumption during pregnancy and its impact on child development. In: Tremblay RE, Barr RG, Peters RDeV, eds. *Encyclopedia on Early Childhood Development* [online]. Montreal, Quebec: Centre of Excellence for Early Childhood Development; 2002:1-5. Available at: <http://www.child-encyclopedia.com/documents/FriedANGxp.pdf>. Accessed March 14, 2011.
7. Dietz P, England L, Shapiro-Mendoza C, Tong V, Farr S, Callaghan W. Infant morbidity and mortality attributable to prenatal smoking in the U.S. *Am J Prev Med*. 2010;39:45-52.
8. Lotfipour S, Leonard G, Perron M, Pike B, Richer L, Seguin J, Toro R, Veillette S, Pausova Z, Paus T. Prenatal exposure to maternal cigarette smoking interacts with a polymorphism in the $\alpha 6$ nicotinic acetylcholine receptor gene to influence drug use and striatum volume in adolescence. *Mol Psychiatry*. 2010;15:6-8; doi: 10.1038/mp.2009.63.
9. Stroud L, Paster R, Goodwin M, Shenassa E, Buka S, Niaura R, Rosenblith J, Lipsitt L. Maternal smoking during pregnancy and neonatal behavior: A large-scale community study. *Pediatrics* 2009;23:e842 - e848.
10. Pickett K, Wood C, Adamson J, DeSouza L, Wakschlag L. Meaningful differences in maternal smoking behaviour during pregnancy: Implications for infant behavioural vulnerability. *J Epidemiol Community Health*. 2008;62:318-324.
11. Key A, Ferguson M, Mofese D, Peach K, Lehman C, Molfese V. Smoking during pregnancy affects speech-processing ability in newborn infants. *Environ Health Perspect*. 2007;115(4):623-629.
12. Golub M, Slotkin T, Tarantal A, Pinkerton K. Visual recognition memory and auditory brainstem response in infant rhesus monkeys exposed perinatally to environmental tobacco smoke. *Brain Res*. 2007;1151:102-106.
13. Lewis B, Kirchner H, Short E et al. Prenatal cocaine and tobacco effects on children's language trajectories. *Pediatrics* 2007;120:e78-e85.
14. Carter S, Paterson J, Gao W, Lusitini L. Maternal smoking during pregnancy and behaviour problems in a birth cohort of 2-year-old Pacific children in New Zealand. *Early Hum Dev*. 2008;84:59-66.

15. Huijbregts S, Seguin J, Zoccolillo M, Boivin M, Tremblay R. Role of maternal prenatal smoking, parental antisocial behavior and early childhood physical aggression. *Dev Psychopathol.* 2008;20:437-453.
16. Ashford J, van Lier P, Timmermans M, Cuijpers P, Koot H. Prenatal smoking and internalizing and externalizing problems in children studied from childhood to late adolescence. *J Am Acad Child Adolesc Psychiatry.* 2008;47:779-787.
17. Cornelius M, Goldschmidt L, De Genna N, Day N. Smoking during teenage pregnancies: Effects on Behavioral problems in offspring. *Nicotine Tob Res.* 2007;9:739-750.
18. Cornelius M, De Genna N, Leech S, Willford J. Effects of prenatal tobacco exposure on neurobehavioral outcomes of 10-year-old children of teenage mothers. *Neurotoxicol Teratol.* 2011;33:137-144. NIHMS234727; PMID: NTT6178.
19. Ruckinger S, Rzehak P, Chen C, Sausenthaler S, Koetzko S, Bauer C, Hoffmann U, Kramer U, Berdel D, von Berg A, Bayer O, Wichmann H, von Kries R, Heinrich J. Prenatal and postnatal tobacco exposure and behavioral problems in 10-year-old children: Results from the GINI-plus Prospective birth cohort study. *Environ Health Perspect.* 2010;118:150-154.
20. Brion M, Victora C, Matijasevich A, Horta B, Anselmi L, Menezes A, Lawlor D, Smith G. Maternal smoking and child psychological problems: disentangling causal and noncausal effects. *Pediatrics* 2010;126:e57-e65.
21. Langley K, Holmans P, Van den Bree M, Thapar A. Effects of low birthweight, maternal smoking during pregnancy and social class on the phenotypic manifestations of Attention Deficit Hyperactivity Disorder and associated antisocial behaviour: Investigation in a clinical sample. *BMC Psychiatry* 2007;7:e26.
22. Gatzke-Kopp L, Beauchaine T. Direct and passive prenatal nicotine exposure and the development of externalizing psychopathology. *Child Psychiatry Hum Dev.* 2007;38:255-269.
23. Murray J, Irving B, Farrington D, Colman I, Bloxson C. Very early predictors of conduct problems and crime: results from a national cohort study. *J Child Psychol Psychiatry.* 2010;51:1198-1207.
24. Johansson A, Ludvigsson J, Hermansson G. Adverse health effects related to tobacco smoke exposure in a cohort of three-year-olds. *Acta Paediatr.* 2008;97:354-357.
25. Gilman S, Gardener H, Buka S. Maternal smoking during pregnancy and children's cognitive and physical development: A causal risk factor? *Am J Epidemiol.* 2008;168:522-531.
26. Ino T. Maternal smoking during pregnancy and offspring obesity: Meta-analysis. *Pediatr Int.* 2010;52:94-99.
27. Oken E, Levitan E, Gillman M. Maternal smoking during pregnancy and child overweight: Systematic review and meta-analysis. *Int J Obest (Lond).* 2008;32:201-210.
28. Rooney B, Mathiason M, Schauburger C. Predictors of obesity in childhood, adolescence, and adulthood in a birth cohort. *Matern Child Health J.* 2010 DOI 10.1007/s10995-010-0689-1.
29. Menezes A, Goncalves H, Anselmi L, Hallal P, Araujo C. Smoking in early adolescence: Evidence from the 1993 Pelotas (Brazil) Birth Cohort Study. *J Adolesc Health.* 2006;39:669-677.
30. Agrawal A, Scherrer J, Grant J, Sartor C, Pergadia M, Duncan A, Madden P, Haber J, Jacob T, Bucholz K, Xian H. The effects of maternal smoking during pregnancy on offspring outcomes. *Prev Med.* 2010;50:13-18.
31. O'Callaghan F, Mamun A, O'Callaghan M, Alati R, Najman J, Williams G, Bor W. Maternal smoking during pregnancy predicts nicotine disorder (dependence or withdrawal) in young adults - a birth cohort study. *Aust N Z J Public Health.* 2009;33:371-377.
32. Lotfipour S, Leonard G, Perron M, Pike B, Richer L, Seguin J, Toro R, Veillette S, Pausova Z, Paus T. Prenatal exposure to maternal cigarette smoking interacts with a polymorphism in the $\alpha 6$ nicotinic acetylcholine receptor gene to influence drug use and striatum volume in adolescence. *Mol Psychiatry.* 2010;15:6-8; doi: 10.1038/mp.2009.63.
33. Slotkin T, MacKillop E, Rudder C, Ryde I, Tate C, Seidler F. Permanent, sex-selective effects of prenatal or adolescent nicotine, exposure, separately or sequentially in rat brain regions: indices of cholinergic and serotonergic synaptic function, cell signaling and neural cell number and size at six months of age. *Neuropsychopharmacology* 2007;32:1082-1097.

34. Wiebe S, Espy K, Stopp C, Respass J, Stewart P, Jameson T, Gilbert D, Huggenciv J. Gene-environment interactions across development: Exploring DRD2 genotype and prenatal smoking effects on self-regulation. *Dev Psychol.* 2009;45:31-44.
35. Becker K, El-Faddagh M, Schmidt M, Esser G, Laucht M. Interaction of dopamine transporter genotype with prenatal smoke exposure on ADHD symptoms. *J Pediatr.* 2008;152:263-269.
36. Neuman R, Lobos E, Reich W, Henderson C, Sun L, Todd R. Prenatal smoking exposure and dopaminergic genotypes interact to cause a severe ADHD subtype. *Biol Psychiatr.* 2007;61:1320-1328.
37. Wakschlag L, Kistner E, Pine D, Biesecker G, Pickett K, Skol A, Dukic V, Blair R, Leventhal B, Cox N, Burns J, Kasza K, Wright R, Cook E. Interaction of prenatal exposure to cigarettes and MAOA genotype in pathways to youth antisocial behavior. *Mol Psychiatry.* 2010;15:928-937.
38. Rivkin M, Davis P, Lemaster J, Cabral H, Warfield S, Mulkern R, Robson C, Rose-Jacobs R, Frank D. Volumetric MRI study of brain in children with intrauterine exposure to cocaine, alcohol, tobacco, and marijuana. *Pediatrics.* 2008;121:741-750.
39. Toro R, Leonard G, Lerner J, Lerner R, Perron M, Pike G, Richer L, Veillette S, Pausova Z, Paus T. Prenatal exposure to maternal cigarette smoking and the adolescent cerebral cortex. *Neuropsychopharmacology* 2008;33:1019-1027.
40. Jacobsen L, Picciotto M, Heath C, Frost S, Tsou K, Dwan R, Constable R, Menci W. Prenatal and adolescent exposure to tobacco smoke modulates the development of white matter microstructure. *J Neurosci.* 2007;27(49):13491-13498.
41. Jacobsen L, Slotkin T, Menci E, Frost S, Pugh K. Gender-specific effects of prenatal and adolescent exposure to tobacco smoke on auditory and visual attention. *Neuropsychopharmacology* 2007;32:2453-2464.
42. Ball S, Gilman S, Mick E, Fitzmaurice, Ganz M, Seidman L, Buka S. Revisiting the association between maternal smoking during pregnancy and ADHD. *J Psychiatr Res.* 2010;22:1058-1062.
43. Boutwell B, Beaver K. Maternal cigarette smoking during pregnancy and offspring externalizing behavioral problems: a propensity score matching analysis. *Int J Environ Res Public Health.* 2010;7:146-163.
44. D'Onofrio B, Van Hulle C, Waldman I, Rodgers, Harden K, Rathouz P, Lahey B. Smoking during pregnancy and offspring externalizing problems: An exploration of genetic and environmental confounds. *Dev Psychopathol.* 2008;20:139-164.
45. Lavigne J, Hopkins J, Gouze K, Bryant F, LeBailly S, Binns H, Lavigne P. Is smoking during pregnancy a risk factor for psychopathology in young children? A methodological caveat and report on preschoolers. *J Pediatr Psychol. AdvanceAccess* 2010 DOI:10.1093/jpepsy/jsqo44.
46. Roza S, Verhulst F, Jaddoe V, Steegers E, Mackenbach J, Hofman A, Tiemeier H. Maternal smoking during pregnancy and child behaviour problems: the Generation R study. *Int J Epidemiol.* 2008;1-10.
47. Button T, Maughan B, McGuffin P. The relationship of maternal smoking to psychological problems in the offspring. *Early Hum Dev.* 2007;83(11):727-732.
48. Cornelius M, Day N. Developmental consequences of prenatal tobacco exposure. *Curr Opin Neurol.* 2009; 22:121-125.
49. Pickett K, Wakschlag L. The short-term and long-term developmental consequences of maternal smoking during pregnancy. In: Preece P, Riley E, eds. *Drugs in Pregnancy: The Price for the Child: Exposure to Fetal Teratogens and Long Term Neurodevelopmental Outcomes.* London: Mac Keith Press; (In press).
50. Knopik V. Maternal smoking during pregnancy and child outcomes: real or spurious effects. *Dev Neuropsychol.* 2009;34:1-36.
51. Shea A & Steiner M. Cigarette smoking during pregnancy. *Nicotine Tob Res.* 2008;10(2):267-278.
52. Slotkin T. If nicotine is a developmental neurotoxicant in animal studies, dare we recommend nicotine replacement therapy in pregnant women and adolescents? *Neurotoxicol Teratol.* 2008;20:1-19.
53. Bruin J, Gerstein H, Holloway A. Long-term consequences of fetal and neonatal nicotine exposure: A critical review. *Toxicol Sci.* 2010;116:364-374.

54. Simpson W. A preliminary report of cigarette smoking and the incidence of prematurity. *Am J Obstet Gynecol.* 1957;73:808-815.